

World Journal of *Otorhinolaryngology*

World J Otorhinolaryngol 2023 February 28; 10(1): 1-3



LETTER TO THE EDITOR

- 1 Examining the life-cycle of the Coblator II device: Increases in paediatric post-tonsillectomy haemorrhage after six years of use

Winters R, Rodriguez KH

ABOUT COVER

Editorial Board Member of *World Journal of Otorhinolaryngology*, Mario Rigante, MD, PhD, Associate Professor, Department of Otorhinolaryngology, University of Perugia, Perugia 06132, Italy. mario.rigante@unipg.it

AIMS AND SCOPE

The primary aim of *World Journal of Otorhinolaryngology* (WJO, *World J Otorhinolaryngol*) is to provide scholars and readers from various fields of otorhinolaryngology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of otorhinolaryngology and covering a wide range of topics including adenoidectomy, audiology, auditory brain stem implantation, ciliary motility disorders, cochlear implantation, ear diseases, endolymphatic shunt, fenestration, labyrinth, laryngeal diseases, laryngectomy, laryngoplasty, laryngoscopy, middle ear ventilation, myringoplasty, nasal surgical procedures, neck dissection, nose diseases, ossicular replacement, otologic surgical procedures, otorhinolaryngologic diseases, otorhinolaryngologic neoplasms, otorhinolaryngologic surgical procedures, pharyngeal diseases, pharyngectomy, pharyngostomy, rhinoplasty, stapes surgery, tonsillectomy, tracheostomy, transtympanic micropressure treatment, and tympanoplasty.

INDEXING/ABSTRACTING

The WJO is now abstracted and indexed in Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xu Guo*, Editorial Office Director: *Yu-Jie Ma*.

NAME OF JOURNAL

World Journal of Otorhinolaryngology

ISSN

ISSN 2218-6247 (online)

LAUNCH DATE

December 28, 2011

FREQUENCY

Continuous Publication

EDITORS-IN-CHIEF

Amr El-Shazly

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2218-6247/editorialboard.htm>

PUBLICATION DATE

February 28, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Examining the life-cycle of the Coblator II device: Increases in paediatric post-tonsillectomy haemorrhage after six years of use

Ryan Winters, Kimsey H Rodriguez

Specialty type:

Otorhinolaryngology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model:

Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer:

Abubakar MS,
Nigeria; Shorrah AA, United Arab
Emirates

Received: November 28, 2022

Peer-review started: November 28, 2022

First decision: January 3, 2023

Revised: January 4, 2023

Accepted: February 10, 2023

Article in press: February 10, 2023

Published online: February 28, 2023



Ryan Winters, Department of Otorhinolaryngology-Head & Neck Surgery, NSW Health/John Hunter Hospital, New Lambton Heights 2305, NSW, Australia

Kimsey H Rodriguez, Department of Otolaryngology-Head & Neck Surgery, Ochsner Medical Center, New Orleans, LA 70121, United States

Corresponding author: Ryan Winters, MD, Associate Professor, Department of Otorhinolaryngology-Head & Neck Surgery, NSW Health/John Hunter Hospital, Lookout Road, New Lambton Heights 2305, NSW, Australia. ryan.winters@health.nsw.gov.au

Abstract

All paediatric tonsillectomies were examined from 2012 – 2019 at a single tertiary-referral institution, and all were performed by one of two paediatric otolaryngologists. One exclusively used the diathermy, the other exclusively used the Coblator II. Two Coblator units were purchased simultaneously in 2012 and not replaced. There was no significant difference in number of tonsillectomies performed (1298 *via* diathermy, 1221 *via* Coblator), nor in postoperative day of bleed, patient age, indication for procedure, and no patient had an underlying coagulopathy. The most common indication for tonsillectomy in both groups was sleep-disordered breathing. There was no significant difference in postoperative haemorrhage rates between groups for the first six years of the study (0%-1.4%/year). Years 7 and 8 saw the Coblator group haemorrhage rate significantly increase (0%-0.6% diathermy group *vs* 2%-3% Coblator group), though still fell within accepted rate of postoperative haemorrhage. The devices were then replaced, and the differences in haemorrhage disappeared. There appears to be a significant increase in paediatric post-tonsillectomy haemorrhage when using a Coblator device greater than six years old. This may suggest a useful lifespan for the Coblator II device.

Key Words: Tonsillectomy; Paediatric; Post-operative; Haemorrhage; Diathermy; Coblator

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: There appears to be a significant increase in paediatric post-tonsillectomy haemorrhage when using a coblator device greater than six years old.

Citation: Winters R, Rodriguez KH. Examining the life-cycle of the Coblator II device: Increases in paediatric post-tonsillectomy haemorrhage after six years of use. *World J Otorhinolaryngol* 2023; 10(1): 1-3

URL: <https://www.wjgnet.com/2218-6247/full/v10/i1/1.htm>

DOI: <https://dx.doi.org/10.5319/wjo.v10.i1.1>

TO THE EDITOR

Many techniques have been described for paediatric tonsillectomy, diathermy and Coblator each have long, successful, histories of safe outcomes[1,2]. As with any piece of technology or hardware, there is a potential useful life for the Coblator device after which the reliability becomes more questionable. Ochsner acquired two Coblator devices in January 2012 and used the same devices continuously (without replacement, with manufacturer-prescribed periodic preventative maintenance performed according to the manufacturer's protocol) continuously throughout the study period. We describe the experience of tonsillectomy bleeds by two paediatric otolaryngologists requiring return to theatre at a quaternary referral centre. One surgeon exclusively utilized diathermy, the other exclusively Coblator, all other practice parameters equivalent, from 2012–2019. For insurance reasons, all patients were seen within our health system for all elective and emergent complaints (Table 1).

2519 paediatric tonsillectomies were performed during the study period. 1298 by the diathermy surgeon, 1221 performed by the Coblator surgeon. There were no significant differences with respect to bleeding dyscrasias, medical co-morbidities, age, indications, day of bleed, or bleeds managed expectantly. No significant differences in postoperative haemorrhage were noted for the first five years (2012–2017: Diathermy surgeon: 0%-1.4%/year, Coblator surgeon: 0%-1.3%/year). Years six and seven revealed a significant ($P < 0.05$) increase in haemorrhage requiring return to theatre in the Coblator group (2018–2019 diathermy surgeon: 0%-0.6%/year, Coblator surgeon: 2%-3%/year). This resolved with replacement of the Coblator devices after 2019. As with any hardware device, there is presumably a useful lifespan. The device life for the Coblator II unit has not been previously reported in relation to paediatric tonsillectomy, though device maintenance for Coblator-based reusable wands have been discussed in the orthopaedic literature[3]. These orthopaedic reports are significantly different in that the complication was from the reusable wand, rather than the Coblator unit itself. The wand used in paediatric tonsillectomy is single-use, but each was connected to the same Coblator devices in our study. It appears that the rate of postoperative haemorrhage essentially doubles for Coblator tonsillectomy after six years of continued use. Even this elevated haemorrhage rate falls within the "acceptable" range for paediatric tonsillectomy, but is still significantly elevated when compared to younger devices. This could describe a "working life" for the Coblator device in paediatric tonsillectomy. Lower haemorrhage rates could potentially be maintained with regular device replacement[4].

Table 1 Postoperative haemorrhage rates by year and surgical technique

Year	Coblator actual (%)	Diathermy actual (%)	P value
2012	0-0	0-0	1.0
2013	1-0.7	2-1	0.77
2014	1-0.7	0-0	0.30
2015	2-1.4	1-0.5	0.38
2016	2-0.9	1-0.5	0.62
2017	1-0.6	2-1.3	0.51
2018	4-2	1-0.6	0.05
2019	5-3	0-0	0.03

FOOTNOTES

Author contributions: All authors participated in experimental design, patient care/protocol application, and manuscript preparation and review.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-

NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Australia

ORCID number: Ryan Winters 0000-0003-2909-5667.

S-Editor: Xing YX

L-Editor: A

P-Editor: Xing YX

REFERENCES

- 1 **Divi V**, Benninger M. Postoperative tonsillectomy bleed: coblation vs noncoblation. *Laryngoscope* 2005; **115**: 31-33 [PMID: 15630361 DOI: 10.1097/01.mlg.0000150682.62517.0e]
- 2 **Pynnonen M**, Brinkmeier JV, Thorne MC, Chong LY, Burton MJ. Coblation vs other surgical techniques for tonsillectomy. *Cochrane Database Syst Rev* 2017; **8**: CD004619 [PMID: 28828761 DOI: 10.1002/14651858.CD004619.pub3]
- 3 **Kim Y**, Kim HK, Yoon JR, Muzaffar N, Kim TS, Shin YS. Separation of the tip of a coblation wand within the knee joint: a complication of arthroscopic adhesiolysis. *Clin Orthop Surg* 2010; **2**: 125-127 [PMID: 20514271 DOI: 10.4055/cios.2010.2.2.125]
- 4 **Reusser NM**, Bender RW, Agrawal NA, Albright JT, Duncan NO, Edmonds JL. Post-tonsillectomy hemorrhage rates in children compared by surgical technique. *Ear Nose Throat J* 2017; **96**: E7-E11 [PMID: 28719712 DOI: 10.1177/014556131709600702]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>



World Journal of *Otorhinolaryngology*

World J Otorhinolaryngol 2023 May 9; 10(2): 4-35



REVIEW

- 4 Post-COVID-19 persistent olfactory, gustatory, and trigeminal chemosensory disorders: Definitions, mechanisms, and potential treatments

Hamed SA

CASE REPORT

- 23 Transoral robotic surgery for adult parapharyngeal lymphangioma: A case report

Dabas S, Menon NN, Ranjan R, Gurung B, Shukla H, Sharma AK, Tiwari S, Sinha A, Bhatti SS, Sangal R

- 30 Wet cupping (Al-hijamah) as a strange cause of ear trauma: A case report

Al-Ani RM

ABOUT COVER

Peer Reviewer of *World Journal of Otorhinolaryngology*, Kádima N Teixeira, PhD, Professor, Campus Toledo, Universidade Federal do Paraná, 3796 Max Planck Av, 85919-899, Toledo, PR Brazil. kadimateixeira@ufpr.br

AIMS AND SCOPE

The primary aim of *World Journal of Otorhinolaryngology* (WJO, *World J Otorhinolaryngol*) is to provide scholars and readers from various fields of otorhinolaryngology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of otorhinolaryngology and covering a wide range of topics including adenoidectomy, audiology, auditory brain stem implantation, ciliary motility disorders, cochlear implantation, ear diseases, endolymphatic shunt, fenestration, labyrinth, laryngeal diseases, laryngectomy, laryngoplasty, laryngoscopy, middle ear ventilation, myringoplasty, nasal surgical procedures, neck dissection, nose diseases, ossicular replacement, otologic surgical procedures, otorhinolaryngologic diseases, otorhinolaryngologic neoplasms, otorhinolaryngologic surgical procedures, pharyngeal diseases, pharyngectomy, pharyngostomy, rhinoplasty, stapes surgery, tonsillectomy, tracheostomy, transtympanic micropressure treatment, and tympanoplasty.

INDEXING/ABSTRACTING

The WJO is now abstracted and indexed in Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Si Zhao*; Production Department Director: *Xu Guo*; Editorial Office Director: *Yun-Jie Ma*.

NAME OF JOURNAL

World Journal of Otorhinolaryngology

ISSN

ISSN 2218-6247 (online)

LAUNCH DATE

December 28, 2011

FREQUENCY

Continuous Publication

EDITORS-IN-CHIEF

Amr El-Shazly

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2218-6247/editorialboard.htm>

PUBLICATION DATE

May 9, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Post-COVID-19 persistent olfactory, gustatory, and trigeminal chemosensory disorders: Definitions, mechanisms, and potential treatments

Sherifa Ahmed Hamed

Specialty type: Neurosciences

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Bordonaro M, United States; Teixeira KN, Brazil

Received: January 20, 2023

Peer-review started: January 20, 2023

First decision: March 28, 2023

Revised: March 28, 2023

Accepted: April 18, 2023

Article in press: April 18, 2023

Published online: May 9, 2023



Sherifa Ahmed Hamed, Department of Neurology and Psychiatry, Assiut University, Faculty of Medicine, Assiut 71516, Egypt

Corresponding author: Sherifa Ahmed Hamed, MD, Professor, Department of Neurology and Psychiatry, Assiut University, Faculty of Medicine, Assiut University Street, Assiut 71516, Egypt. hamedsherifa@aun.edu.eg

Abstract

The nose and the oral cavities are the main sites for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) entry into the body. Smell and taste deficits are the most common acute viral manifestations. Persistent smell disorders are the most common and bothersome complications after SARS-CoV-2 infection, lasting for months to years. The mechanisms and treatment of persistent post-coronavirus disease 2019 (COVID-19) smell and taste disorders are still challenges. Information sources for the review are PubMed, Centers for Disease Control and Prevention, Ovid Medline, Embase, Scopus, Web of Science, International Prospective Register of Systematic Reviews, Cumulative Index to Nursing and Allied Health Literature, Elton Bryson Stephens Company, Cochrane Effective Practice and Organization of Care, Cooperation in Science and Technology, International Clinical Trials Registry Platform, World Health Organization, Randomized Controlled Trial Number Registry, and MediFind. This review summarizes the up-to-date information about the prevalence, patterns at onset, and prognoses of post-COVID-19 smell and taste disorders, evidence for the neurotropism of SARS-CoV-2 and the overlap between SARS-CoV-1, Middle East respiratory syndrome coronavirus, and SARS-CoV-2 in structure, molecular biology, mode of replication, and host pathogenicity, the suggested cellular and molecular mechanisms for these post-COVID19 chemosensory disorders, and the applied pharmacotherapies and interventions as trials to treat these disorders, and the recommendations for future research to improve understanding of predictors and mechanisms of these disorders. These are crucial for hopeful proper treatment strategies.

Key Words: COVID-19; SARS-CoV-2; Coronaviruses; Olfactory and gustatory chemosensory disorders; Anosmia; Ageusia; Parosmia; Neuronal degeneration; Neurogenesis

Core Tip: Smell loss is the most frequent acute manifestation of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. Persistent smell disorders (deficits and distortions) are the most frequent viral complications. Taste and flavor disorders are also complications of SARS-CoV-2 infection but at lower frequencies compared to smell disorders. It has been found that SARS-CoV-2 has a 4 times more affinity to infect the olfactory epithelium compared to other human coronaviruses. The mechanisms of these disorders have been explored mainly based on animal models of anosmia due to SARS-CoV-2 infection. It has been suggested that the post-coronavirus disease 2019 transient smell loss might be due to viral infection to the olfactory non-neuronal epithelial cells (particularly the sustentacular cells) which are important for the health of olfactory sensory neurons (OSNs). These cells rapidly regenerate after injury, within 1-3 wk, and restore smell function. Persistent smell disorders have been suggested to be due to injury of OSNs, disorganization of the olfactory epithelium, altered expression of olfactory receptors, and impaired olfactory neurogenesis. These cells require ≥ 3 mo to regenerate and restore function depending on the severity and type of injury. Taste disorders have been suggested to be due to viral infection of taste buds, disruption of the activity of the salivary glands, inflammation of the gustatory epithelium, and injury to the taste sensory cells. Treatment of these disorders is a medical challenge, and none of the available pharmacotherapies or interventions which are used to treat similar disorders due to other causes, showed curative effect.

Citation: Hamed SA. Post-COVID-19 persistent olfactory, gustatory, and trigeminal chemosensory disorders: Definitions, mechanisms, and potential treatments. *World J Otorhinolaryngol* 2023; 10(2): 4-22

URL: <https://www.wjgnet.com/2218-6247/full/v10/i2/4.htm>

DOI: <https://dx.doi.org/10.5319/wjo.v10.i2.4>

INTRODUCTION

Increasing data and meta-analyses reported a high prevalence of smell loss (anosmia/hyposmia) as an acute manifestation of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection (40%-86%) [1]. It has been reported that smell loss (anosmia/hyposmia) could be the hallmark or an isolated symptom of SARS-CoV-2 infection without other viral manifestations in 25%-44% of patients [1,2]. True taste loss (ageusia/hypogeusia) has been estimated to occur in 10.2%-42% of patients due to SARS-CoV-2 infection [1-3]. True nasal trigeminal chemosensory loss has been estimated to occur in 20%-33% of patients due to SARS-CoV-2 infection [4].

Long-lasting or persistent smell disorders (deficits and distortion), lasting for ≥ 6 mo, are the most common complications of SARS-CoV-2 infection with an estimated prevalence of ~20%-40% [1,5,6]. Parosmia is the most frequent type of smell distortions, occurring in 65% of patients after an interval of a month or more from the onset of smell loss or resolution of other acute viral manifestations. A few studies reported phantosmia in 20% of patients [7]. Distortions of taste (dysgeusia) or flavor (aroma) perception have been reported in ~12% [3].

There have been no systematic studies which determined the course of smell and taste disorders after SARS-CoV-2 variants. Hintschich *et al* [8] reported that the wild type SARS-CoV-2 was associated with a higher prevalence of hyposmia (73%) according to patients' self-reports and psychophysical testing compared to the alpha (41%) or delta (48%) variants. Boscolo-Rizzo *et al* [9] observed that 30% of patients infected with the omicron variant had olfactory dysfunction. Vihta *et al* [10] reported smell loss in 13%-16% of patients during the dominant period of omicron variant infection compared to 44% when the delta variant pandemic was dominant. The authors suggested that the difference in the omicron spike protein might result in less effective cell membrane fusion and host olfactory cell entry. However, the increase in the prevalence of chemosensory disorders after omicron infection has been suggested to be due to the fact that the transmission of omicron variant is 4-fold higher compared to the delta variant [11].

Information sources for the review are PubMed (<https://www.ncbi.nlm.nih.gov/research/coronavirus/>), the Centers for Disease Control and Prevention, Ovid Medline, Embase, Scopus, Web of Science, International Prospective Register of Systematic Reviews, Cumulative Index to Nursing and Allied Health Literature, Elton Bryson Stephens Company, Cochrane Effective Practice and Organization of Care, Cooperation in Science and Technology, International Clinical Trials Registry Platform, World Health Organization, United States National library of Medicine, Cochrane ENT Trials Register, Randomized Controlled Trial Number Registry, and MediFind. A manual search was also done for additional information. Search words were SARS-CoV-2, coronavirus disease 2019 (COVID-19), coronaviruses (CoVs), β -CoVs, neurotropism, smell, taste, flavor, trigeminal chemosensation,

olfactory sensation, gustatory sensation, olfactory neuronal regeneration, post-COVID-19 syndrome, anosmia, ageusia, parosmia, dysgeusia, and long-lasting or persistent post-COVID-19 chemosensory disorders. In this review, we discuss the prevalence of post-COVID-19 chemosensory disorders, their patterns and prognoses, the cellular and molecular mechanisms underlying these disorders, and the potential pharmacotherapies and interventions for their correction. A brief background about the olfactory and gustatory pathways, perceptual processes, and neuronal degeneration/regeneration is provided to facilitate readers to understand the pathophysiology underlying these disorders secondary to viral infection.

BACKGROUND

Nasal chemosensation


Smell sensation is carried out by three chemosensory systems[12]: (1) The main olfactory system which carries out perception of volatile odorants; (2) the accessory olfactory system which carries out perception of non-volatile substances. These substances are not consciously perceived as odors. They include pheromones or hormone metabolites as sweat, urine, seminal fluid, and vaginal secretion; and (3) the trigeminal olfactory system that carries out perception of pungent sensations (*i.e.*, with irritant components).

There are two routes for volatile odor perception: The orthonasal and retronasal routes. In orthonasal perception, the odor molecules (also known as scents) enter the nasal cavity through the nostrils after sniffing and then to the olfactory sensory epithelium. In retronasal perception, the odor molecules of foods and drinks are transported back to the nasopharynx after swallowing or during respiratory exhalation and then to the olfactory sensory epithelium (Figure 1). Retronasal olfaction contributes to the flavor of foods or drinks (the aroma perception). It is associated with taste sensation[13].

The olfactory information is carried out from the olfactory neuroepithelium, nerves, bulb, and brain. The olfactory mucosa is composed of the olfactory epithelium and the underlying lamina propria. The olfactory epithelium has olfactory sensory neurons (OSNs) and non-neuronal cells. OSNs are bipolar cells. The bipolar cell has dendrites at one end with hair-like cilia extending to the mucus lining the nasal cavity and the other end has a single apical axon which projects directly to a glomerulus in the olfactory bulb. The olfactory receptors are odor binding G protein-coupled that are expressed in the membranes of the bipolar cell dendrites[14] (Figure 2A). The axons of the bipolar cells are the olfactory nerve fibers[15]. It has been observed that a single odorant molecule has specific chemical properties to either bind to a selective receptor or a number of receptors[14]. The olfactory epithelium is spatially organized into non-overlapping zones in which each is dominated by a certain set of odor receptors, meaning that a specific odor molecule activates a specific zone within the epithelium but not others. The normal olfactory function requires highly regulated projections from sensory neurons and regrowth of nascent axons *en masse* onto the olfactory bulb for proper integration of olfactory signals in the olfactory cortex[16]. Therefore, the same topographical mapping is maintained in the glomeruli and bulb since formed during development and throughout life (Figure 2A and B). The glomerular map is important as the glomeruli act as organizers for the odorant signals before sending them to the brain for odor detection, recognition, and discrimination[17]. Within the glomerulus, the axon of the bipolar neuron synapses with the mitral and tufted cells along with periglomerular cells (Figure 2A).

The olfactory non-neuronal cells include the sustentacular (also known as supporting cells), microvillar, Bowman's gland, and horizontal basal (stem) cells (Figure 2A). These cells, the nasal respiratory epithelial cells, and the vascular pericytes are important for the health of OSNs. Sustentacular cells surround and support OSNs, detoxify harmful molecules, and maintain epithelial salt and water balance[18]. The Bowman's glands secrete mucous[18]. The mucus contains large concentrations of small soluble proteins which keep odorant molecules dissolved in the mucus. The odorant-protein complexes bind to the receptors[15]. The microvillar cells (also known as brush cells) help in protection of OSNs by stimulating the sustentacular cells to metabolize and detoxify harmful molecules [19], and have a role in adjustment of ion gradients which is important for maintenance of the function of OSNs, and their basal surfaces are in contact with afferent nerve endings of the trigeminal nerve for transduction of nasal general sensation[18]. Basal cells are multipotent stem cells that undergo continuous division during life to produce new OSNs and non-neuronal epithelial cells, *i.e.*, endogenous regeneration process[18,20].

In the process of olfaction, the chemical information of the odorants converts into electrical information. The odorant molecules trigger a signaling cascade which includes activation of the cyclic nucleotide-gated ion channels followed by activation of adenylyl cyclase type 3 (ADCY3). ADCY3 is localized into the cilia of OSNs. ADCY3 activation is an essential component to convert adenine triphosphate to cyclic adenine monophosphate (cAMP) (*i.e.*, the olfactory cAMP signaling pathway) [21]. cAMP unfolds ion channels resulting in influx of calcium (Ca^{++}) and sodium (Na^{+}) inside the cell. This opens Ca^{++} -activated chloride (Cl^{-}) channels leading to outflow of Cl^{-} , resulting in neuronal membrane depolarization and propagation of action potential from the olfactory epithelium to the bulb and then the brain[22] (Figure 3). The primary olfactory cortical areas include the uncus, anterior



DOI: 10.5319/wjo.v10.i2.4 Copyright ©The Author(s) 2023.

Figure 1 Olfactory perception and its pathways. There are two routes for odor perception: The orthonasal and retronasal routes. The process of olfaction converts chemical information from odorants into electrical potentials extending from the olfactory epithelium to the bulb and then the brain. Brain areas for perception of olfaction sensation are divided into the primary and the secondary cortical olfactory areas. The primary areas include the uncus, anterior olfactory nucleus, olfactory tubercle, piriform cortex, lateral entorhinal cortex, and the cortical nucleus of the amygdala. The secondary areas include the hypothalamus, mediodorsal thalamic nucleus, nucleus basalis Meynert, hippocampus, the septal region, substantia innominata, mesencephalic reticular formation, and the orbitofrontal cortex.

olfactory nucleus, piriform cortex, lateral entorhinal cortex, olfactory tubercle, and cortical nucleus of the amygdala[23]. The primary olfactory cortex projects to the secondary olfactory cortical areas for further processing of olfactory impulses as odor perception, discrimination, attention, memory, and motivational and emotional aspects of smell. These areas are the hypothalamus, mediodorsal thalamic nucleus, nucleus basalis Meynert, hippocampus, septal region, substantia innominata, mesencephalic reticular system, and orbitofrontal cortex[24] (Figure 1).

The trigeminal chemosensation allows the perception of pungent, irritant, burning, stinging, tickling, and noxious chemicals odorants even in the absence of olfactory sensation[25] and triggers the protective respiratory reflexes (as sneezing and coughing). This system relies on free nerve endings and on the trigeminal-innervated solitary chemosensory cells (SCCs) in the nasal epithelium. The free nerve endings recognize the lipophilic irritant odors (as mints and ammonia) while the SCCs enhance the chemosensory capabilities of the trigeminal irritant-detection system through expression of certain receptors for detection of noxious chemicals[26].

It has been reported that the olfactory and trigeminal systems are intimately connected at peripheral (*i.e.*, nasal mucosa) and central (the piriform cortex, the ventral insula, and the frontal cortex) levels[27, 28].

Oral chemosensation

Till now, the five known chemical structures that taste sensing cells can detect are sweet (sugar), salty (salt), sour (lemon juice), bitter (unsweetened decaffeinated coffee), and umami or savory (proteins as meat and aged cheese). Everything else is known as flavor which is the function of retronasal olfaction [29,30]. All regions of the tongue that detect taste respond to all the five taste qualities; however, it has been observed that the sensations of sweet and salt are best detected in the tip of the tongue. The highest concentrations of bitter receptors are in the posterior tongue[31] (Figure 4A). Tastants are detected by specialized receptor cells within the taste buds of the tongue, soft palate, and epiglottis. In the tongue, taste buds are located within the tongue papillae[29]. Most papillae have 3-5 taste buds, but some types of papillae have thousands of taste buds. The taste bud has taste sensing cells and non-neuronal cells. The taste sensing cells express specific receptors[32]. The taste receptor cells are arranged within the bulb in a manner that their tips form a small pore in which the microvilli or hair-like projections extend to the saliva layer within the oral cavity. The saliva interacts with and protects the taste receptors in the mouth. For the transduction of taste stimuli, the molecules of food or drink must be dissolved in the saliva[30] (Figure 4B). A single taste bud has 50 to 100 taste cells, among which there are 10 and 50 taste




Figure 2 Structure of the olfactory mucosa and topographical mapping of the olfactory sensory neurons and their connections. A: The olfactory mucosa is composed of the olfactory epithelium and the lamina propria. The epithelium is composed of olfactory sensory neurons and non-neuronal cells (the sustentacular, microvillar, and Bowman's gland and basal cells). A bipolar cell has dendrites at one end extending to the mucus and a single axon which projects to the glomerulus. Within a glomerulus, the axon of the bipolar neuron synapses with the mitral and tufted cells along with periglomerular cells; B: The olfactory neurons which express the same type of receptors converge onto the same glomerulus maintaining a topographical mapping of odorant receptors where they synapse with sensory neurons of the olfactory tract. The same topographical mapping is maintained in the glomeruli and bulb.




Figure 3 The process of olfaction. The signaling cascade which converts chemical information of the odorant into electrical information includes activation of cyclic nucleotide-gated ion channel followed by activated ADCY3 to convert ATP to cyclic adenosine monophosphate (cAMP). cAMP unfolds Ca^{++} activated Cl^- channels, causing influx of Na^+ and Ca^{++} from outside to inside the olfactory sensory neurons, and this activates the Cl^- channels, causing outflow of Cl^- , neuronal membrane depolarization, and the action potential. The latter extends from the epithelium to the bulb and then to the olfactory cortex.

sensory cells. The non-neuronal cells include the supporting cells and the basal cells. The basal cells are stem cells which actively divide throughout life into taste sensory cells and accessory gustatory epithelial cells (Figure 4A and B).

In the process of gustation, the tastant molecules are connected to their specialized receptors and upon doing so, activation of nerve impulses occurs and the receptors release neurotransmitters which activate sensory neurons in the facial, glossopharyngeal, and vagus cranial nerves. The taste information




Figure 4 Distributional mapping of the five basic taste sensations on the tongue, the gustatory pathways, and the taste bud and receptors. A: The five basic taste qualities are sweet, salty, sour, bitter, and umami; B: Tastants are detected by the taste buds on the tongue, soft palate, and epiglottis. Activation of taste receptors is transmitted to the sensory neurons in the facial, glossopharyngeal, and vagus cranial nerves from which their axons carry information to the nucleus solitarius in the medulla oblongata, to the ipsilateral ventral posteromedial nucleus of the thalamus, and then to the primary gustatory cortex. The nuclei in the medulla also send projections to the hypothalamus and amygdalae to activate brainstem reflexes. Secondary fibers project from the insula to the posterolateral portion of the orbitofrontal cortex; C: The taste bud has taste sensory and supporting cells. The taste receptors have long microvilli or hairs which project through the taste pore to connect with substances in the food or drink dissolved into the saliva. VPM: Ventral posteromedial.

is carried out to the nucleus solitarius in the medulla oblongata, the ipsilateral ventral posteromedial nucleus of the thalamus, and then the primary gustatory cortex which includes the frontal operculum or anterior insular cortex located near the inferior margin of the post-central gyrus. The nuclei in the medulla also send projections to the hypothalamus and amygdalae, which are involved in autonomic reflexes such as gagging and salivation[32]. The primary gustatory cortex is important for discrimination of taste sensations. It also receives information about the smell and texture of food. The secondary fibers travel from the primary gustatory cortex to the posterolateral portion of the orbitofrontal cortex for integration of taste and smell sensations (Figure 4).

Studies found that the orally sourced odors share processing circuitry with taste and internal odors to produce flavor preferences. They found that inactivation of the insular gustatory cortex impaired selectively the expression of retronasal preferences[13]. Trigeminal chemosensation is also considered a part of flavor perception[32].

Endogenous neurogenesis processes of olfactory and gustatory cells

The olfactory and gustatory neurons in the human body are the only types of neurons that are regularly replaced within the nasal and oral epithelia throughout life by newly generated neurons as a part of the turnover process (*i.e.*, life-long olfactory and gustatory sensory neurogenesis)[33]. Replacement of olfactory neurons occurs at 4-12 wk intervals[16,33]. The gustatory receptor cells are short-lived and replaced every 10 d[34]. This process has been understood to happen because the olfactory mucosa can be easily harmed by airborne and toxic chemicals and viruses, and the gustatory mucosa can be easily damaged by the activities that occur in the mouth. A reconstitution of the sensory epithelium occurs from the stem cells with a subsequent restoration of olfactory function. This process requires a complex signaling system, neurotropic factors, neurotransmitters, *etc.* After injury, the regeneration and restoration of olfactory receptors and their connections depend on the degree and type of injury. In many circumstances, the olfactory system overcomes many obstacles encountered when rewiring the olfactory bulb after olfactory neuroepithelial injury, and can be able to maintain its capacity to regenerate new axon processes and reestablish functional connections within the olfactory bulb[16].

Investigators observed that when injury was limited to the olfactory neuroepithelium sparing the olfactory nerves, the regenerating new axons grew along the axons that were already in place, and the newly rewired connections converged onto the glomeruli in specific areas maintaining the glomerular topographical mapping of odorant receptors and re-established the functional connections with the olfactory bulb[35]. It has been discovered that stabilization of olfactory receptor expression and guidance and stabilization of the axonal growth of OSNs require ADCY3[36-38]. However, if injury involved the olfactory neuroepithelium and nerves, there was a competition among the regenerating axons to occupy synaptic sites within a glomerulus, with inability of the glomerulus to be dominated by a single odorant, resulting in impaired odor discrimination and anosmia and dysosmia[39-41] (Figure 5A and B). Those authors also observed that long-lasting anosmia can alter the integrity of the mitral and tufted cell dendrites. Murai *et al*[42] observed reduced connectivity of mitral/tufted cell dendrites and attenuated odor responses in these cells after olfactory neuronal injury.

PATTERNS AND PROGNOSSES OF SMELL AND TASTE DISORDERS WITH SARS-CoV-2 INFECTION

Studies reported two main patterns for smell loss (with or without taste or flavor loss) as an acute manifestation of SARS-CoV-2 infection: (1) Sudden smell loss (anosmia/hyposmia) in association with general, systemic, or other ear, nose, and throat manifestations. Anosmia is the term used to define the absence or loss of the sense of smell. Hyposmia is the decreased sense of smell. This is the most frequent pattern. It has been reported that smell loss occurs either at the same time with other viral manifestations (22.8%-88.0%)[1,2], after the recovery of other viral manifestations (26.6%-65.4%)[1,2], or before the onset of other manifestations (11.8%)[2]; and (2) Isolated smell loss without any other viral manifestations (~16%-19.4%)[43].

Clinical studies reported that the most frequent other acute nasal symptoms of SARS-CoV-2 infection are dryness, itching, nasal burning, and sneezing, occurring in 75%-87% of patients, while nasal congestion and blockage and anterior nasal discharge (rhinorrhea), which are the main symptoms due to other viral infections of the upper respiratory tract (*e.g.*, influenza), are rare with SARS-CoV-2 infection, occurring in only 12.9%-25%[2,44,45].

Some authors observed a mismatch between subjective and objective smell and taste manifestations at onset. The high frequency of subjective anosmia and ageusia compared to the low frequency of true ageusia detected by objective evaluation (40%-86% *vs* 10.2%-42%), is not surprising. The belief that taste will be lost with smell loss is based on the fact that the ability to perceive flavors is adversely affected by the concomitant presence of smell loss[46].

The majority of studies reported the presence of true ageusia exclusively in patients with anosmia, while ageusia without anosmia was rarely reported[3,47]. Loss of flavor of food and drinks (aroma perception) is also an infrequent manifestation of SARS-CoV-2 infection but has been found to be always dependent on the presence of ageusia but not anosmia[3]. Subjective sudden loss to pungent or irritant odors and tastants (*e.g.*, mint gum, black pepper, ginger, and spice) has been frequently reported with severe smell loss after SARS-CoV-2 infection[4,48]. However, true nasal trigeminal chemosensory loss due to SARS-CoV-2 infection, as detected by objective evaluation, has been reported in only 18%-20% and exclusively reported in association with anosmia[4]. True oral trigeminal chemosensory loss has been reported in 20%-33% and exclusively reported in association with ageusia[4].

Studies also reported two prognoses for these disorders. They are: (1) Transient deficits which resolve within days to weeks after onset (mean: ~20 d). This is the most frequent prognosis occurring in 60%-80% of patients[49,50]; and (2) Long-lasting/persistent disorders (deficits and distortions) lasting for months to years, occurring in 20%-40%[49]. Many studies reported that the diagnosis of mild COVID-19 at onset would predict the prognosis of persistent smell disorders. Lechien *et al*[2] reported that the persistent smell disorders were prevalent in patients with isolated conditions or if associated with mild COVID-19 manifestations (86%-95% *vs* 4.5%-10% with moderate/severe manifestations). However, the increasing number of published studies has shown that none of the demographics, acute manifestations, and severity at onset were predictors of the development of persistent disorders[1,5].

Dysosmia is a frequent manifestation (~75%-80%) in patients with persistent post-COVID smell loss. Dysosmia is a qualitative or distortion of perception of an odor. It is another smell or a strange sensation (pleasant or unpleasant) of inhaled odorants. Dysosmia often develops after a month or more from the onset of acute manifestations or smell loss[2,44]. Parosmia is the frequently encountered type of smell distortion after SARS-CoV-2 infection, occurring in 65% of patients. Parosmia is defined as perception of an odor in the presence of odorous source[7]. Some patients would develop an excessively unpleasant repulsive, unbearable, disgusting, or intolerable odor (cacosmia). A few studies reported the occurrence of phantosmia in patients with persistent post-COVID-19 anosmia (20%). Phantosmia is defined as distortion of smell perception in which there is a perception of an odor in the absence of odorous source [7]. Dysgeusia or distortion of taste and distortion of aroma[3] were only reported in patients with persistent post-COVID taste loss (~12%-20%). Common descriptions for parosmia or dysgeusia or distortion of flavor included burnt (like burnt hair or burnt leather), rotten (as spoiled or fermented




Figure 5 Mechanism of dysosmia after severe acute respiratory syndrome coronavirus 2 infection. A: The disorganized olfactory neuroepithelium after severe acute respiratory syndrome coronavirus 2 infection and its associated immunological response would result in a disturbance of rewiring of the glomeruli; B: A glomerulus is no longer dominated by a single type of odor and the newly growing axon after injury would go to a different brain spot other than the one before injury, resulting in a distorted sense of smell.

food), gasoline, fecal, smoke or metallic (like copper), chemical (like sulphur), etc. However, many described the unpleasant unknown odor which was difficult to describe (*i.e.*, new)[7].

Magnetic resonance imaging (MRI) had been used for evaluation of olfactory structures and connections after SARS-CoV-2 infection in patients with anosmia. However, findings were limited and conflicting. Some found olfactory bulb edema and/or olfactory cleft edema in patients with anosmia, suggesting local inflammation and a greater local immune response[51-56]. Some authors reported microbleeding or a break in the blood-brain barrier (BBB) of the olfactory bulbs and tracts[57,58]. Kandemirli *et al*[59] reported a high percentage of abnormal olfactory bulb signal intensity and contour changes in cases with persistent COVID-19 anosmia ($n = 23$). Yildirim *et al*[60] used MRI, digital tensor imaging, and olfactory functional MRI for evaluation of patients. They found decreased olfactory bulb volume, indicating bulb damage and decreased white matter tract integrity of the olfactory regions. The authors evaluated the olfactory bulb in patients with persistent post-COVID-19 olfactory disorders ($n = 31$, 100% were anosmic and had a mean interval of 1.5 mo between onset of olfactory disorder and imaging) and patients with other post-infectious olfactory disorders ($n = 97$, among which 18.6% were hyposmic and 81.4% were anosmic and had a mean interval of 6 mo between onset of olfactory disorder and imaging). Compared to patients with post-infectious olfactory disorders, the authors found significantly higher volumes of olfactory bulbs in patients with persistent post-COVID-19 olfactory disorders, deformed bulb morphology in 58.1% (*vs* 63.9%, with no significant difference), increased olfactory bulb signal intensity in 51.6% (*vs* 46.4%, with no significant difference), and a significantly higher rate of olfactory nerve clumping and anisotropy values at orbitofrontal and entorhinal regions but no significant difference had been identified between the two groups in the orbitofrontal and entorhinal activity apart of the robust trigemino-sensory activity in patients with persistent post-COVID-19 olfactory disorders but not in those with other post-infectious olfactory disorders.

NEUROTROPISM POTENTIALS OF COVS AND ROUTES FOR VIRAL SPREAD TO NERVOUS TISSUE WITH SPECIAL EMPHASIS ON SARS-COV-1, MIDDLE EAST RESPIRATORY SYNDROME-COV, AND SARS-COV-2

There are two known groups of human CoVs (HCoVs): (1) α -CoVs which include HCoV-229E (named after a student specimen coded 229E) and HCoV-NetherLand 63 (NL63)[61]; and (2) β -CoVs which include HCoV-Hong Kong University 1 (HKU1), HCoV-Organ Culture 43 (OC43), SARS-CoV-1, Middle East respiratory syndrome (MERS)-CoV, and SARS-CoV-2[61]. HCoV-229E, HCoV-NL63, HCoV-HKU1, and HCoV-OC43 induce mild and spontaneously resolved (*i.e.*, transient) respiratory symptoms such as rhinitis, sore throat, and dry cough similar to common cold or enteric and systemic manifestations[62]. SARS-CoV-1[63], MERS-CoV[64], and SARS-CoV-2[65] are dangerous CoVs. SARS-CoV-1 was firstly

identified in China in 2002-2003. MERS-CoV was firstly identified in Saudi Arabia in 2012. SARS-CoV-2 was firstly identified in Wuhan, China in December 2019[65]. They are the causes of severe and potentially fatal respiratory tract infections and the known three worldwide pandemics[61,66]. Human neurotropic CoVs are CoV-229E, CoV-OC43, SARS-CoV-1, MERS-CoV, and SARS-CoV-2[63,64].

In general, CoVs are frequently recognized as respiratory pathogens[67]; however, some have neuroinvasive and neurotropic properties[61]. In animal models, some authors detected that some CoVs spread to the brain using the neuronal pathway (*i.e.*, peripheral trigeminal or olfactory systems) or through disruption of the BBB. In the mouse model, Perlman *et al*[68] found that ablation of the olfactory pathway did not allow the neural spread of CoVs. Hara *et al*[69] found that the viral particles of hemagglutinating encephalomyelitis virus, a member of β -CoVs, bud into the endoplasmic reticulum-Golgi cells and form virion vesicles into Golgi cells, followed by secretion into the surroundings and infection of adjacent nerve cells[70]. Koyuncu *et al*[71] reported nasal mucociliary transport and disruption of the ciliary nasal epithelium with HCoV-229E[72]. Mori *et al*[40] found entry of CoVs to the central nervous system through the olfactory tract in earlier stages of viral infection. Dubé *et al*[73] detected HCoV-OC43 in the piriformis cortex, brainstem, and spinal cord by the fourth day after nasal viral inoculation and indicated that HCoV-OC43 can propagate from neuron to neuron through axonal transportation. Netland *et al*[74] detected SARSCoV1 in mice expressing human angiotensin-converting enzyme 2 (ACE2) in the olfactory bulb after 60 h of nasal viral inoculation. They also found the virus after four days of inoculation in the pyriform cortex, ventral pallidum, and lateral preoptic regions in the basal ganglia and dorsal nucleus of Rafe in the midbrain. However, the authors did not look for a potential infection of the olfactory mucosa for being the route of SARS-CoV-1 to the brain, *i.e.*, where these neurons project. Li *et al*[75] found MERS-CoV in the brain of mice expressing human CD26/dipeptidyl peptidase 4, particularly the thalamus and brainstem after nasal viral inoculation. Bryche *et al*[76] observed damage and desquamation of the olfactory sensory epithelium and loss of cilia in golden Syrian hamsters as early as 2 d after nasal inoculation of SARS-CoV-2. The authors observed infection of a large proportion of sustentacular cells but not OSNs or bulbs. Yinda *et al*[77] found SARS-CoV-2 in K18-hACE2 mice in the olfactory bulb and the connected brain areas after nasal viral inoculation. Those authors also found that the taste receptors of the gustatory sensory epithelium were targets for SARS-CoV-2 to enter and infect the body[78]. Hematogenous spread of the viruses to the brain has been found in influenza and other coronaviruses[71]. In a mouse model, Bleau *et al*[79] found that hepatitis coronavirus had the ability to cross the BBB and infect the brain. Some authors suggested that the observed brain hemorrhage or microbleeding in some patients after SARS-CoV-2 infection could be attributed to disruption and increase in the permeability of the BBB caused by the virus[80].

In autopsies of infected individuals, studies reported the presence of CoVs particles in neuronal and glial cells and brain endothelial capillary pericytes. Ding *et al*[81] found SARS-CoV-1 particles in the cerebral hemispheres. Steardo *et al*[82] detected SARS-CoV-2 antigens in the nucleus solitarius and the nucleus ambiguus in the brainstem. Coolen *et al*[83] detected SARS-CoV-2 particles in only the olfactory bulbs in early postmortem study of infected patients with anosmia. Solomon *et al*[84] used immunohistochemistry, electron microscopy, and real-time reverse transcription polymerase chain reaction (RT-PCR) in their investigations. The authors found SARS-CoV-2 particles in the olfactory epithelium and bulb. Lee *et al*[85] used high-field MRI to examine brain tissue of dead patients. The authors found microvascular damage consistent with endothelial activation and widespread vascular injury due to viral spread to the brain.

It has been found that SARS-CoV-1, MERS-CoV, and SARS-CoV-2 have close relation to each other and to animal neurotropic CoVs in structure, molecular biology, mode of replication, and host pathogenicity[86]. SARS-CoV-1 (26 kb), MERS-CoV (32 kb), and SARS-CoV-2 (30 kb) have the largest genome among RNA viruses[66]. SARS-CoV-2 has 79.5% identical RNA sequence to SARS-CoV-1 and 96% identical RNA sequence to bat CoV[62,65,66]. SARS-CoV-2 has a similar 3D structure and homology of the trimeric S proteins and their receptor binding domains as SARS-CoV-1[87]. SARS-CoV-1 [63,88,89], MERS-CoV[64], and SARS-CoV-2[76,87,90,91] enter the body and infect similar specific olfactory sensory, nasal respiratory, and tongue epithelial cells. However, apart from olfactory and gustatory chemosensory disorders after SARS-CoV-2 infection, several studies observed that SARS-CoV-2 infection did not directly infect the brain and did not follow the same pathway as SARS-CoV-1. They found that anosmia due to SARS-CoV-2 infection did not correlate with encephalopathies observed in some infected patients[80,92-94]. Also many studies did not detect SARS-CoV-2 RNA by RT-PCR in the cerebrospinal fluid from living patients who were infected with SARS-CoV-2 and had neuropsychiatric manifestations[94]. Investigators suggested that the non-specific acute symptoms such as disturbed consciousness, delirium, encephalopathy, lack of concentration, fatigue, headache, and generalized weakness could be attributed to the acute severe illness with respiratory and metabolic disturbances or as a non-specific cellular immune or antibody-mediated phenomenon (*i.e.*, peri- or postinfectious processes or a para-infectious cytokine storm)[94,95]. Some authors suggested that SARS-CoV-2-related encephalopathy could include interruption of the BBB and microbleeding, brain hemorrhage[80,96], or clotting abnormalities[92].

CELLULAR AND MOLECULAR MECHANISMS OF SARS-COV-2 INDUCED SMELL AND TASTE DISORDERS

Mechanism(s) of anosmia

Evidence from most studies indicates that the nose is the main site for SARS-CoV-2 entry to the body and the site for viral shedding during the incubation period. Virus shedding is the phase in which the virus reproduces and exports a large amount of copies[93]. It has been indicated that post-COVID-19 smell loss is not solely related to nasal inflammation associated with mucosal edema, and airflow obstruction as in other upper respiratory infections[44]. Experimental studies found that SARS-CoV-2 mainly infects the olfactory non-neuronal epithelial cells (sustentacular cells, microvillar, mucin secreting, and basal cells), nasal vascular pericytes, and the nasal respiratory epithelium[97]. Previous studies also reported that SARS-CoV-1 and MERS-CoV enter the body from the supporting olfactory sensory epithelial cells, the nasal respiratory epithelial cells particularly the sustentacular cells, and the pericytes of the olfactory bulb[61]. These cells, particularly the sustentacular cells, are overloaded by ACE2 receptors[76,90,93]. The virus infects these cells through interaction between its spike glycoprotein (S) and ACE2 protein on target cells. This interaction requires cleavage of the S protein by transmembrane serine protease 2. In a golden Syrian hamster model of SARS-CoV-2 infection, investigators found viral infection of a large amount of sustentacular cells within the first 48 h after nasal inoculation of the virus, followed by rapid and massive invasion by immune cells in the nasal mucosa. This resulted in disorganization followed by desquamation of the olfactory epithelium. The authors found the desquamated epithelium and the cilia of OSNs in the nasal cavity and a reduction of ~80% of the thickness of the olfactory epithelium. The authors did not detect the virus in OSNs or the bulb. They also observed complete absence of the virus from the nasal cavity after 7-14 d and restoration of ~50% of the olfactory epithelium[76,90]. Zhang *et al*[98] observed infection of mature and immature OSNs by the virus but to a lesser extent compared to sustentacular cells infection. Injury of the accessory epithelial cells should disturb the function of OSNs, resulting in smell deficits[99]. Injury of the sustentacular cells disturbs the support of OSNs and salt and water balance of the epithelium[99]. Injury of the Bowman's glands will disrupt nasal mucous production. The change in the mucous density and the breakdown of mucin barrier will diminish olfactory sensitivity through reduction of the number of odorant molecules that can adhere to the receptors[100], resulting in smell loss, nasal dryness[100], or distortion of smell sensation[101]. Injury to microvillar cells will disturb epithelial ion gradients[18] and impair transduction of nasal sensation[19]. It has been suggested that injury of supporting cells by the viral infection could explain the occurrence of transient anosmia at onset of viral infection in the majority of patients [47]; because these cells rapidly regenerate within 1-3 wk after degeneration[33]. This is also compatible with the observed rapid restoration, within 14 d, of $\geq 50\%$ of the injured olfactory epithelium in hamsters infected with SARS-CoV-2[76,90].

However, long-lasting post-COVID-19 anosmia and dysosmia are due to injury of OSNs (receptors), because these cells require at least 3 mo to regenerate after injury[16]. OSNs do not express ACE2 receptors. Therefore, it is plausible to suggest that long-lasting/persistent post-COVID-19 anosmia and dysosmia could be due to severe damage and disorganization of the olfactory epithelium by viral infection and its associated immunopathology which could also include the stem cells, resulting in delay or even lack of restoration of the lost epithelial cells and reconstitution of the injured epithelium[16].

Some aspects of the molecular mechanisms of SARS-CoV-2 infection to the olfactory system have been explored in experimental models and supported the notion that injury/dysfunction of OSNs is the cause of smell disorders. Thakur *et al*[102] used the UV-neutralized serum from hamsters infected with SARS-CoV-2 to investigate the molecular mechanisms of SARS-CoV-2 induced olfactory injury and dysfunction. They found rapid changes in the nuclear architecture of OSNs. Zazhytska *et al*[91] investigated the hamster's olfactory epithelia infected with SARS-CoV-2. The authors found: (1) Widespread, severe, and persistent down-regulation of olfactory receptor genes and ADCY3, and other signaling genes for odor perception; (2) A delay in the transcription of OSNs compared to sustentacular cells; (3) The disruption and down-regulation of olfactory receptors' layers occurred before the down-regulation of *LHX2* and *EBF* genes, suggesting that disruption of the olfactory receptor genome architecture may be the earliest insult in dysfunction of OSNs and SARS-CoV-2 induced anosmia. *LHX2* and *EBF* are the transcription factors which regulate the olfactory receptor choice; and (4) The persistence of down-regulation of *LHX2* and *EBF* even after the restoration of olfactory receptors. Investigators observed overlap in the molecular changes in the OSNs caused by different SARS-CoVs, which include: (1) Reduction in expression of genes involved in every step of odor detection including down-regulation in olfactory receptors and their signaling molecules (*e.g.*, *ADCY3*)[103] and reduction in olfactory receptor proteins[15]; (2) Reduction in olfactory receptor chaperones[104]; and (3) Reduction in ion channels generating odor-evoked axon potential[105].

Mechanism(s) of dysosmia

It has been suggested that the cause of dysosmia is the disorganization of the topographical mapping of the receptors or OSNs within the epithelium due to SARS-CoV-2 infection pathology (Figure 5A). Therefore, a glomerulus will not be dominated by a single type of odor and the newly growing axons

will run in different conduits and reach a different brain spot from that before injury[36-38] (Figure 5B). It has been reported that dysomia is a marker of neuronal degeneration or regeneration (*i.e.*, a step toward neuronal recovery or neuronal regeneration after degeneration) regardless of its cause (*e.g.*, after upper respiratory infections, head trauma, and nasal sinus disease)[42]. Also, it is indicated that dysomia never exists with complete anosmia, meaning that it requires a relatively intact sensory system for its expression[106] and disappears with complete recovery of smell sensation over time or after therapy[42].

Mechanisms of ageusia

The mechanisms of post-COVID-19 true taste disorders are less understood compared to smell disorders. SARS-CoV-2 was found in the saliva of patients with viral infection[107] and viral replication has been shown to occur in human taste buds[108]. In animal models, investigators observed that SARS-CoV-2 selectively infects the tongue compared to the non-lingual epithelium. They observed that ACE2 is mainly expressed in epithelial cells outside of taste papillae which contain the taste buds[78], the human salivary gland also expresses high amounts of ACE2[109], and the tongue taste buds express sialic acid. The taste buds also express abundant amounts of Toll-like receptors (TLRs) compared to the other components of the oral cavity[110]. Sialic acid is a salivary mucin component. Mucin conveys the molecules of tastants into taste pores to prevent their premature enzymatic degradation[111]. The main function of the TLRs is to recognize the common structural components of microorganisms and activate the endogenous inflammatory immune system. Previous studies found a plethora of cytokines and chemokines in the nasal cavity and paranasal sinuses in patients with chronic rhinosinusitis and rhinitis in response against microbes[112]. Studies suggested that SARS-CoV-2 potentially uses multiple entry oral receptors, which are the ACE2, sialic acid receptors, and TLRs[113]. Therefore, binding of SARS-CoV-2 to sialic acid receptors could interfere with glycoprotein mediated transport of tastants and accelerate the degradation of the gustatory particles, which contributes to loss of taste[114]. Previous studies also reported the ability of the MERS-CoV and SARS-CoV-1 to bind the sialic acid receptors on the taste buds[115]. The disruption in the activity of the salivary glands by viral infection could produce imbalance or disruption of saliva composition, impairment of salivary flow, and the continuous renewal of tongue epithelial cells, resulting in dry mouth, impairment of taste transduction, and ageusia[78,116,117]. Studies observed that SARS-CoV-2 binding with oral mucosal cells might trigger inflammation, abnormal cell turnover, and reduced taste bud sensitivity[78,118]. *In situ* models of direct binding of coronavirus spike protein with TLR1, 4, and 6 have supported the specific roles of these TLRs in CoV-2 entry and SARS-CoV-2 infection[119]. It has been suggested that inflammation could increase epithelial cell exfoliation and constitute potential sources of viral SARS-CoV-2 RNA in saliva[120,121]. Experimental studies provided evidence that inflammatory cytokines induced by viral infections could negatively affect the function of taste buds[110]. Previous findings also suggested that viral invasion to taste cells can lead to genetic material exchange between the virus and taste cell subpopulations and this could be the cause of virus-induced taste disorders[122,123].

Mechanism(s) of dysgeusia

The mechanisms of dysgeusia due to SARS-CoV-2 infection are less understood compared to parosmia. It is possible that dysgeusia could be due to the reduction of the salivary flow which normally causes continuous renewal of tongue epithelial cells[107]. We also suggest that dysgeusia could be a trial of regeneration of gustatory neuronal cells after its degeneration by viral pathology and the disorganization of taste receptors' layering within the taste buds.

Mechanism(s) of trigeminal chemosensory disorders

The trigeminal chemosensory loss after SARS-CoV-2 infection could be due to loss of smell and flavor. This is because of the intimate relation between the olfactory, trigeminal, and gustatory systems[19,27,28]. It could also be due to the damage of the nasal epithelium and its feeding blood vessels by viral infection.

APPLIED TREATMENT POTENTIALS FOR PERSISTENT POST-COVID-19 SMELL AND TASTE DISORDERS

Smell and taste disorders have important impacts not only at the physical level but also at the emotional and psychological wellbeing level. Smell is important for detection and discrimination of odors, enjoyment of the energizing aroma, judging the qualitative value of odors (pleasant or unpleasant), influencing social behaviors (*e.g.*, social interaction as feelings of attraction or disgust), evoking good memories, and warning against toxins, leaking gas, polluted air, smoke, *etc.* Taste and flavor are important for food choice, enjoyment of the depth and complexity of food and drinks, the general appetite and control of food intake, and the onset of satiation and warn against rotten or toxic foods and drinks.

Post-COVID-19 persistent smell and taste disorders affect the individual not only at the physical but also at the psychological level and can have an impact on the quality of life. Affected patients are often distressed as these impairments can hinder the enjoyment of food and create hygiene problems. Studies from Europe and Asia reported the adverse impact of long-lasting smell disorders on quality of life, personal-social functioning, and mental health. In a cross-sectional study by Bagheri *et al* [124] on 10069 participants responding to an online checklist that evaluated the sense of smell and taste, the results indicated a significant correlation between anosmia caused by SARS-CoV-2 infection, decreased taste sensation, and decreased quality of life.

Trials to treat persistent disorders include the following pharmacotherapies and interventions which are used in routine clinical practice to treat similar disorders from other etiologies [125,126], and they include: Topical intranasal or systemic steroids [126,127], trace elements (*e.g.*, zinc and selenium) [128,129], vitamins [*e.g.*, vitamin B complex (*e.g.*, B1, B2, B6, B9, and B12), vitamin D, vitamin C, and vitamin E] [130,131], herbals and other nutritional supplements (*e.g.*, omega-3, L-carnitine [131,132], and alpha-lipoic acid [133]), and olfactory and gustatory training [134].

Zinc is widely used by otolaryngologists to treat smell and taste disorders. Zinc has a role in antiviral immune responses [135,136]. Zinc supplementation has shown efficacy to improve taste disorders which may or may not be accompanied by acute hypozincemia [137]. Zinc also has been shown to inhibit coronavirus RNA polymerase activity *in vitro* [138]. The idea of using B vitamins [thiamine (B1), pyridoxine (B6), and cobalamin (B12)] is based on the fact that these vitamins have neurotropic properties. They can greatly help in improving injured nervous system from different etiologies. They support the development of the new nervous system cells, maintain neuronal viability, and protect nerves against damaging environmental factors. Vitamin B1 facilitates the use of carbohydrates for energy production and acts as a site-directed antioxidant. Vitamin B6 balances nerve metabolism. Vitamin B12 promotes nerve cell survival and maintains myelin sheath and remyelination. Vitamin D, vitamin C, and vitamin E enhance host immunity and have antioxidant properties [139]. Omega-3 contains both docosahexaenoic acid and eicosapentaenoic acid. Omega-3 fatty acid deficient mice demonstrate evidence of olfactory dysfunction and mice receiving omega-3 fatty acids have improved recovery after peripheral nerve injury, which has been linked to neuroprotective effects mediated through antioxidant and anti-inflammatory pathways. Omega-3 also improves immunity and enhances energy production in different tissues [132]. L-carnitine transports long-chain fatty acids into the mitochondria to be oxidized for energy production [131]. A recent study showed that the loss of mitochondrial membrane potential could be related to long-lasting smell loss after SARS-CoV-2 infection [140]. These findings suggest that interventions using antioxidants and microelements conducted early in the course of the disease (including neuroprotectors), along with olfactory training, could be options for treating post-COVID-19 anosmia. Pentoxifylline is a hemorrheologic agent which improves peripheral microvascular blood flow and causes vasodilatation of peripheral blood vessels and could be an option for treating post-COVID-19 anosmia [141].

Olfactory training has been shown to be effective to regain smell loss from different causes; however, its results vary between subjects but generally slow, occurring over months to years. The idea of olfactory training is to strengthen the olfactory nerves by routine smelling. Training often involves daily sniffing of four strong scents (such as lemon, rosemary, curry powder, lavender, pungent herbs, spices cloves, rose, and essential oils containing scents) at least twice a day, as follows: The patient has to choose one scent and smell it for 15-20 s and try to remember what it smelled like if the patient has complete smell loss, then takes a rest for 10 s, followed by smelling the next scent for 15-20 s, then rest for 10 s, and so on until all four scents have been sampled. After 3 mo, the patient has to switch to a new set of four scents and train with them as described before. For parosmia, scent training can also be used, which involves inhaling a particular scent and thinking about what that scent should smell like [134]. For gustatory training, trials include the following: (1) When eating, if the patient cannot taste the full range of flavors of a dish, he/she has to pay attention to the basic five tastants: Sweet, bitter, sour, salty, or umami, as well as to the food's texture and the sensation which is felt in the palate. This will help to focus on what still can be tasted, rather than on what the subject cannot taste; and (2) The patient has to choose meals with a variety of colors and textures using aromatic herbs and spices and to add substances of stronger flavors to the food, such as aged cheese, olive oil, and toasted nuts [134].

None of the above mentioned treatments or interventions provided significant efficacy and the treatment of persistent smell and taste disorders is still a medical challenge. An increasing number of registered trials have been emerged; however, only two were completed and showed that the use of steroids had no therapeutic efficacy [126].

CONCLUSION

The patterns of acute manifestations and the course of smell and taste disorders caused by SARS-CoV-2 infection indicate that there are different proportions of damage of the olfactory epithelial cells by viral infection. Therefore, it is impossible to predict the long-term functional outcome in patients infected with SARS-CoV-2 after the acute deficits. SARS-CoV-2 infects and damages the non-neuronal olfactory

epithelial cells which are important for the health of the OSNs. SARS-CoV-2 causes chemosensory neuronal degeneration, delayed or lack of regeneration, and molecular changes in the host genomic architecture resulting in reduced expression of olfactory sensory receptors. Therefore, the neurogenesis after viral pathology and axonal outgrowth, connections, and re-wiring and the role of neurotrophic factors to restore the olfactory function after SARS-CoV-2 infection are important topics for further investigations. Further experimental studies are also required to determine the cellular and molecular mechanisms of post-COVID-19 taste disorders as they are much understudied compared to smell disorders. There is also a need for studies to determine the different phenotypes and the extent of smell and taste disorders with different mutated SARS-CoV-2 viral infections and the prevalence of persistent smell and taste disorders due to SARS-CoV-2 variant forms (*i.e.*, wild type alpha or delta, omicron, *etc.*). The treatment of persistent smell disorders is still a challenge. None of pharmacotherapies and interventions which were used to treat similar disorders from other etiologies showed any beneficial effect. To my knowledge, there are currently 13 registered trials as shown in the web [WHO International Clinical Trials Registry Platform (ICTRP Search Portal-WHO; <https://trialsearch.who.int/>), Cochrane ENT Trials Register (<https://ent.cochrane.org>), Ovid Embase (<https://tools.ovid.com>), ClinicalTrials.gov, MedFind (<https://medifind.in>)] including the use of the following: Local steroids; Systemic steroids; Omega 3; Quadruple; Intranasal insulin; Intranasal zinc; Intranasal gabapentin; Ice cube stimulation; Aerosolized retinoic acid; Intranasal theophylline; A combination of oral vitamin A and intense aromatic chemosensory smell training by pulse aromatic stimulation; Platelet-rich plasma isolated from a patient's own blood and intramuscular cerebrolysin, a multimodal neurotropic factor. However, results were "Recruiting" or "Not started recruiting yet" or "Pending" except for two completed trials for the use of steroids but their results showed no efficacy. Further understanding of the unforeseen mechanisms of these disorders may provide insights for beneficial therapies.

FOOTNOTES

Author contributions: Hamed SA was the guarantor and designed the study, did information collection, designed the figures, did manuscript drafting, and revised the article critically for important intellectual content.

Conflict-of-interest statement: The authors declare no conflict of interests for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Egypt

ORCID number: Sherifa Ahmed Hamed 0000-0002-1441-3530.

S-Editor: Zhang H

L-Editor: Wang TQ

P-Editor: Zhao S

REFERENCES

- 1 **Koyama S**, Ueha R, Kondo K. Loss of Smell and Taste in Patients With Suspected COVID-19: Analyses of Patients' Reports on Social Media. *J Med Internet Res* 2021; **23**: e26459 [PMID: 33788699 DOI: 10.2196/26459]
- 2 **Lechien JR**, Chiesa-Estomba CM, Beckers E, Mustin V, Ducarme M, Journe F, Marchant A, Jouffe L, Barillari MR, Cammaroto G, Ciciu MP, Hans S, Saussez S. Prevalence and 6-month recovery of olfactory dysfunction: a multicentre study of 1363 COVID-19 patients. *J Intern Med* 2021; **290**: 451-461 [PMID: 33403772 DOI: 10.1111/joim.13209]
- 3 **Sheen F**, Tan V, Lim AJ, Haldar S, Sengupta S, Allen D, Somani J, Chen HY, Tambyah P, Forde CG. The COVOSMIA-19 trial: Preliminary application of the Singapore smell and taste test to objectively measure smell and taste function with COVID-19. *Food Qual Prefer* 2022; **97**: 104482 [PMID: 34848929 DOI: 10.1016/j.foodqual.2021.104482]
- 4 **Parma V**, Ohla K, Veldhuizen MG, Niv MY, Kelly CE, Bakke AJ, Cooper KW, Bouysset C, Pirastu N, Dibattista M, Kaur R, Liuzza MT, Pepino MY, Schöpf V, Pereda-Loth V, Olsson SB, Gerkin RC, Rohlf's Dominguez P, Albayay J, Farruggia MC, Bhutani S, Fjaeldstad AW, Kumar R, Menini A, Bensafi M, Sandell M, Konstantinidis I, Di Pizio A, Genovese F, Öztürk L, Thomas-Danguin T, Frasnelli J, Boesveldt S, Saatci Ö, Saraiva LR, Lin C, Golebiowski J, Hwang LD, Ozdener MH, Guàrdia MD, Laudamiel C, Ritchie M, Havlíček J, Pierron D, Roura E, Navarro M, Nolden AA, Lim J, Whitcroft KL, Colquitt LR, Ferdenzi C, Brindha EV, Altundag A, Macchi A, Nunez-Parra A, Patel ZM, Fiorucci S, Philpott CM, Smith BC, Lundström JN, Mucignat C, Parker JK, van den Brink M, Schmuker M, Fischmeister FPS, Heinbockel T, Shields VDC, Faraji F, Santamaría E, Fredborg WEA, Morini G, Olofsson JK, Jalessi M, Karni N, D'Errico A, Alizadeh R, Pellegrino R, Meyer P, Huat C, Chen B, Soler GM, Alwashahi MK, Welge-Lüssen A, Freiherr J, de Groot

- JHB, Klein H, Okamoto M, Singh PB, Hsieh JW; GCCR Group Author, Reed DR, Hummel T, Munger SD, Hayes JE. More Than Smell-COVID-19 Is Associated With Severe Impairment of Smell, Taste, and Chemesthesis. *Chem Senses* 2020; **45**: 609-622 [PMID: [32564071](#) DOI: [10.1093/chemse/bjaa041](#)]
- 5 **Niklassen AS**, Drafi J, Huart C, Hintschich C, Bockberger S, Trecca EMC, Klimek L, Le Bon SD, Altundag A, Hummel T. COVID-19: Recovery from Chemosensory Dysfunction. A Multicentre study on Smell and Taste. *Laryngoscope* 2021; **131**: 1095-1100 [PMID: [33404079](#) DOI: [10.1002/lary.29383](#)]
- 6 **Lechien JR**, Vaira LA, Saussez S. Prevalence and 24-month recovery of olfactory dysfunction in COVID-19 patients: A multicentre prospective study. *J Intern Med* 2023; **293**: 82-90 [PMID: [36000469](#) DOI: [10.1111/joim.13564](#)]
- 7 **Amin MA**, Nahin S, Ferdous J, Rozars MFK, Hawlader MDH. Short duration phantosmia changes in a post-COVID-19 patient in Bangladesh. *Clin Case Rep* 2022; **10**: e05616 [PMID: [35356165](#) DOI: [10.1002/ccr3.5616](#)]
- 8 **Hintschich CA**, Wenzel JJ, Hummel T, Hankir MK, Kühnel T, Vielsmeier V, Bohr C. Psychophysical tests reveal impaired olfaction but preserved gustation in COVID-19 patients. *Int Forum Allergy Rhinol* 2020; **10**: 1105-1107 [PMID: [32613712](#) DOI: [10.1002/alar.22655](#)]
- 9 **Boscolo-Rizzo P**, Hummel T, Hopkins C, Dibattista M, Menini A, Spinato G, Fabbri C, Emanuelli E, D'Alessandro A, Marzolino R, Zanelli E, Cancellieri E, Cargnelutti K, Fadda S, Borsetto D, Vaira LA, Gardenal N, Polesel J, Tirelli G. High prevalence of long-term olfactory, gustatory, and chemesthesis dysfunction in post-COVID-19 patients: a matched case-control study with one-year follow-up using a comprehensive psychophysical evaluation. *Rhinology* 2021; **59**: 517-527 [PMID: [34553706](#) DOI: [10.4193/Rhin21.249](#)]
- 10 **Vihita KD**, Pouwels KB, Peto TE, Pritchard E, House T, Studley R, Rourke E, Cook D, Diamond I, Crook D, Clifton DA, Matthews PC, Stoesser N, Eyre DW, Walker AS; COVID-19 Infection Survey team. Omicron-associated changes in SARS-CoV-2 symptoms in the United Kingdom. *Clin Infect Dis* 2022; **76**: e133-e141 [PMID: [35917440](#) DOI: [10.1093/cid/ciac613](#)]
- 11 **Meng B**, Abdullahi A, Ferreira IATM, Goonawardane N, Saito A, Kimura I, Yamasoba D, Gerber PP, Fathi S, Rathore S, Zepeda SK, Papa G, Kemp SA, Ikeda T, Toyoda M, Tan TS, Kuramochi J, Mitsunaga S, Ueno T, Shirakawa K, Takaori-Kondo A, Brevini T, Mallery DL, Charles OJ; CITIID-NIHR BioResource COVID-19 Collaboration; Genotype to Phenotype Japan (G2P-Japan) Consortium; Ecuador-COVID19 Consortium, Bowen JE, Joshi A, Walls AC, Jackson L, Martin D, Smith KGC, Bradley J, Briggs JAG, Choi J, Madissoon E, Meyer KB, Mlcochova P, Ceron-Gutierrez L, Doffinger R, Teichmann SA, Fisher AJ, Pizzuto MS, de Marco A, Corti D, Hosmillo M, Lee JH, James LC, Thukral L, Veleser D, Sigal A, Sampaziotis F, Goodfellow IG, Matheson NJ, Sato K, Gupta RK. Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts infectivity and fusogenicity. *Nature* 2022; **603**: 706-714 [PMID: [35104837](#) DOI: [10.1038/s41586-022-04474-x](#)]
- 12 **Boehm U**. Accessory Olfactory System. In: Binder MD, Hirokawa N, Windhorst U. Encyclopedia of Neuroscience. Springer: Berlin, 2008
- 13 **Blankenship ML**, Grigorova M, Katz DB, Maier JX. Retronasal Odor Perception Requires Taste Cortex, but Orthonasal Does Not. *Curr Biol* 2019; **29**: 62-69.e3 [PMID: [30581018](#) DOI: [10.1016/j.cub.2018.11.011](#)]
- 14 **Duchamp-Viret P**, Chaput MA, Duchamp A. Odor response properties of rat olfactory receptor neurons. *Science* 1999; **284**: 2171-2174 [PMID: [10381881](#) DOI: [10.1126/science.284.5423.2171](#)]
- 15 **Buck L**, Axel R. A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* 1991; **65**: 175-187 [PMID: [1840504](#) DOI: [10.1016/0092-8674\(91\)90418-x](#)]
- 16 **Iwema CL**, Fang H, Kurtz DB, Youngentob SL, Schwob JE. Odorant receptor expression patterns are restored in lesion-recovered rat olfactory epithelium. *J Neurosci* 2004; **24**: 356-369 [PMID: [14724234](#) DOI: [10.1523/JNEUROSCI.1219-03.2004](#)]
- 17 **Vassar R**, Chao SK, Sitcheran R, Nuñez JM, Vosshall LB, Axel R. Topographic organization of sensory projections to the olfactory bulb. *Cell* 1994; **79**: 981-991 [PMID: [8001145](#) DOI: [10.1016/0092-8674\(94\)90029-9](#)]
- 18 **Getchell ML**, Getchell TV. Fine structural aspects of secretion and extrinsic innervation in the olfactory mucosa. *Microsc Res Tech* 1992; **23**: 111-127 [PMID: [1421551](#) DOI: [10.1002/jemt.1070230203](#)]
- 19 **Genovese F**, Tizzano M. Microvillous cells in the olfactory epithelium express elements of the solitary chemosensory cell transduction signaling cascade. *PLoS One* 2018; **13**: e0202754 [PMID: [30212469](#) DOI: [10.1371/journal.pone.0202754](#)]
- 20 **Kondo K**, Suzukawa K, Sakamoto T, Watanabe K, Kanaya K, Ushio M, Yamaguchi T, Nibu K, Kaga K, Yamasoba T. Age-related changes in cell dynamics of the postnatal mouse olfactory neuroepithelium: cell proliferation, neuronal differentiation, and cell death. *J Comp Neurol* 2010; **518**: 1962-1975 [PMID: [20394053](#) DOI: [10.1002/cne.22316](#)]
- 21 **Beites CL**, Kawauchi S, Crocker CE, Calof AL. Identification and molecular regulation of neural stem cells in the olfactory epithelium. *Exp Cell Res* 2005; **306**: 309-316 [PMID: [15925585](#) DOI: [10.1016/j.yexcr.2005.03.027](#)]
- 22 **Challis RC**, Tian H, Yin W, Ma M. Genetic Ablation of Type III Adenylyl Cyclase Exerts Region-Specific Effects on Cilia Architecture in the Mouse Nose. *PLoS One* 2016; **11**: e0150638 [PMID: [26942602](#) DOI: [10.1371/journal.pone.0150638](#)]
- 23 **Haberly LB**, Price JL. Association and commissural fiber systems of the olfactory cortex of the rat. II. Systems originating in the olfactory peduncle. *J Comp Neurol* 1978; **181**: 781-807 [PMID: [690285](#) DOI: [10.1002/cne.901810407](#)]
- 24 **Sosulski DL**, Bloom ML, Cutforth T, Axel R, Datta SR. Distinct representations of olfactory information in different cortical centres. *Nature* 2011; **472**: 213-216 [PMID: [21451525](#) DOI: [10.1038/nature09868](#)]
- 25 **Laska M**. Perception of trigeminal chemosensory qualities in the elderly. *Chem Senses* 2001; **26**: 681-689 [PMID: [11473934](#) DOI: [10.1093/chemse/26.6.681](#)]
- 26 **Finger TE**, Böttger B, Hansen A, Anderson KT, Alimohammadi H, Silver WL. Solitary chemoreceptor cells in the nasal cavity serve as sentinels of respiration. *Proc Natl Acad Sci U S A* 2003; **100**: 8981-8986 [PMID: [12857948](#) DOI: [10.1073/pnas.1531172100](#)]
- 27 **Cain WS**, Murphy CL. Interaction between chemoreceptive modalities of odour and irritation. *Nature* 1980; **284**: 255-257 [PMID: [7360255](#) DOI: [10.1038/284255a0](#)]
- 28 **Livermore A**, Hummel T, Kobal G. Chemosensory event-related potentials in the investigation of interactions between the olfactory and the somatosensory (trigeminal) systems. *Electroencephalogr Clin Neurophysiol* 1992; **83**: 201-210 [PMID: [11473934](#) DOI: [10.1093/chemse/26.6.681](#)]

- 1381671 DOI: [10.1016/0013-4694\(92\)90145-8](https://doi.org/10.1016/0013-4694(92)90145-8)
- 29 **Small DM**, Prescott J. Odor/taste integration and the perception of flavor. *Exp Brain Res* 2005; **166**: 345-357 [PMID: [16028032](https://pubmed.ncbi.nlm.nih.gov/16028032/) DOI: [10.1007/s00221-005-2376-9](https://doi.org/10.1007/s00221-005-2376-9)]
- 30 **Honig V**, Procházka P, Obergruber M, Roubík H. Nutrient Effect on the Taste of Mineral Waters: Evidence from Europe. *Foods* 2020; **9** [PMID: [33339333](https://pubmed.ncbi.nlm.nih.gov/33339333/) DOI: [10.3390/foods9121875](https://doi.org/10.3390/foods9121875)]
- 31 **Spence C**. The tongue map and the spatial modulation of taste perception. *Curr Res Food Sci* 2022; **5**: 598-610 [PMID: [35345819](https://pubmed.ncbi.nlm.nih.gov/35345819/) DOI: [10.1016/j.crf.2022.02.004](https://doi.org/10.1016/j.crf.2022.02.004)]
- 32 **Chaudhari N**, Roper SD. The cell biology of taste. *J Cell Biol* 2010; **190**: 285-296 [PMID: [20696704](https://pubmed.ncbi.nlm.nih.gov/20696704/) DOI: [10.1083/jcb.201003144](https://doi.org/10.1083/jcb.201003144)]
- 33 **Whitman MC**, Greer CA. Adult neurogenesis and the olfactory system. *Prog Neurobiol* 2009; **89**: 162-175 [PMID: [19615423](https://pubmed.ncbi.nlm.nih.gov/19615423/) DOI: [10.1016/j.pneurobio.2009.07.003](https://doi.org/10.1016/j.pneurobio.2009.07.003)]
- 34 **Barlow LA**, Klein OD. Developing and regenerating a sense of taste. *Curr Top Dev Biol* 2015; **111**: 401-419 [PMID: [25662267](https://pubmed.ncbi.nlm.nih.gov/25662267/) DOI: [10.1016/bs.ctdb.2014.11.012](https://doi.org/10.1016/bs.ctdb.2014.11.012)]
- 35 **Schwob JE**, Jang W, Holbrook EH, Lin B, Herrick DB, Peterson JN, Hewitt Coleman J. Stem and progenitor cells of the mammalian olfactory epithelium: Taking poietic license. *J Comp Neurol* 2017; **525**: 1034-1054 [PMID: [27560601](https://pubmed.ncbi.nlm.nih.gov/27560601/) DOI: [10.1002/cne.24105](https://doi.org/10.1002/cne.24105)]
- 36 **Imai T**, Suzuki M, Sakano H. Odorant receptor-derived cAMP signals direct axonal targeting. *Science* 2006; **314**: 657-661 [PMID: [16990513](https://pubmed.ncbi.nlm.nih.gov/16990513/) DOI: [10.1126/science.1131794](https://doi.org/10.1126/science.1131794)]
- 37 **Zou DJ**, Chesler AT, Le Pichon CE, Kuznetsov A, Pei X, Hwang EL, Firestein S. Absence of adenylyl cyclase 3 perturbs peripheral olfactory projections in mice. *J Neurosci* 2007; **27**: 6675-6683 [PMID: [17581954](https://pubmed.ncbi.nlm.nih.gov/17581954/) DOI: [10.1523/JNEUROSCI.0699-07.2007](https://doi.org/10.1523/JNEUROSCI.0699-07.2007)]
- 38 **Lyons DB**, Allen WE, Goh T, Tsai L, Barnea G, Lomvardas S. An epigenetic trap stabilizes singular olfactory receptor expression. *Cell* 2013; **154**: 325-336 [PMID: [23870122](https://pubmed.ncbi.nlm.nih.gov/23870122/) DOI: [10.1016/j.cell.2013.06.039](https://doi.org/10.1016/j.cell.2013.06.039)]
- 39 **Yee KK**, Costanzo RM. Changes in odor quality discrimination following recovery from olfactory nerve transection. *Chem Senses* 1998; **23**: 513-519 [PMID: [9805635](https://pubmed.ncbi.nlm.nih.gov/9805635/) DOI: [10.1093/chemse/23.5.513](https://doi.org/10.1093/chemse/23.5.513)]
- 40 **Mori I**, Goshima F, Imai Y, Kohsaka S, Sugiyama T, Yoshida T, Yokochi T, Nishiyama Y, Kimura Y. Olfactory receptor neurons prevent dissemination of neurovirulent influenza A virus into the brain by undergoing virus-induced apoptosis. *J Gen Virol* 2002; **83**: 2109-2116 [PMID: [12185263](https://pubmed.ncbi.nlm.nih.gov/12185263/) DOI: [10.1099/0022-1317-83-9-2109](https://doi.org/10.1099/0022-1317-83-9-2109)]
- 41 **Bozza T**, Vassalli A, Fuss S, Zhang JJ, Weiland B, Pacifico R, Feinstein P, Mombaerts P. Mapping of class I and class II odorant receptors to glomerular domains by two distinct types of olfactory sensory neurons in the mouse. *Neuron* 2009; **61**: 220-233 [PMID: [19186165](https://pubmed.ncbi.nlm.nih.gov/19186165/) DOI: [10.1016/j.neuron.2008.11.010](https://doi.org/10.1016/j.neuron.2008.11.010)]
- 42 **Murai A**, Iwata R, Fujimoto S, Aihara S, Tsuboi A, Muroyama Y, Saito T, Nishizaki K, Imai T. Distorted Coarse Axon Targeting and Reduced Dendrite Connectivity Underlie Dysosmia after Olfactory Axon Injury. *eNeuro* 2016; **3** [PMID: [27785463](https://pubmed.ncbi.nlm.nih.gov/27785463/) DOI: [10.1523/ENEURO.0242-16.2016](https://doi.org/10.1523/ENEURO.0242-16.2016)]
- 43 Loss of sense of smell as marker of COVID-19 infection. 2020 Apr 6 [cited 28 September 2021]. In: ENT UK. Available from: <https://www.entuk.org/Loss-sense-smell-marker-covid-19-infection>
- 44 **Soler ZM**, Patel ZM, Turner JH, Holbrook EH. A primer on viral-associated olfactory loss in the era of COVID-19. *Int Forum Allergy Rhinol* 2020; **10**: 814-820 [PMID: [32271490](https://pubmed.ncbi.nlm.nih.gov/32271490/) DOI: [10.1002/alr.22578](https://doi.org/10.1002/alr.22578)]
- 45 **American Academy of Otolaryngology–Head and Neck Surgery (AAO-HNS)**. Anosmia Reporting tool. 2020 Mar [cited 28 September 2021]. Available from: <https://www.entnet.org/content/reporting-tool-patients-anosmia-related-covid-19>
- 46 **Landis BN**, Scheibe M, Weber C, Berger R, Brämerson A, Bende M, Nordin S, Hummel T. Chemosensory interaction: acquired olfactory impairment is associated with decreased taste function. *J Neurol* 2010; **257**: 1303-1308 [PMID: [20221768](https://pubmed.ncbi.nlm.nih.gov/20221768/) DOI: [10.1007/s00415-010-5513-8](https://doi.org/10.1007/s00415-010-5513-8)]
- 47 **von Bartheld CS**, Hagen MM, Butowt R. Prevalence of Chemosensory Dysfunction in COVID-19 Patients: A Systematic Review and Meta-analysis Reveals Significant Ethnic Differences. *ACS Chem Neurosci* 2020; **11**: 2944-2961 [PMID: [32870641](https://pubmed.ncbi.nlm.nih.gov/32870641/) DOI: [10.1021/acscchemneuro.0c00460](https://doi.org/10.1021/acscchemneuro.0c00460)]
- 48 **Ferrelli F**, Di Bari M, Gaino F, Albanese A, Politi LS, Spriano G, Mercante G. Trigeminal features in COVID-19 patients with smell impairment. *Int Forum Allergy Rhinol* 2021; **11**: 1253-1255 [PMID: [33728825](https://pubmed.ncbi.nlm.nih.gov/33728825/) DOI: [10.1002/alr.22796](https://doi.org/10.1002/alr.22796)]
- 49 **Estiri H**, Strasser ZH, Brat GA, Semenov YR; Consortium for Characterization of COVID-19 by EHR (4CE), Patel CJ, Murphy SN. Evolving Phenotypes of non-hospitalized Patients that Indicate Long Covid. *medRxiv* 2021 [PMID: [33948602](https://pubmed.ncbi.nlm.nih.gov/33948602/) DOI: [10.1101/2021.04.25.21255923](https://doi.org/10.1101/2021.04.25.21255923)]
- 50 **Chapurin N**, Totten DJ, Chaballout B, Brennan J, Dennis S, Lubner R, Chowdhury NI, Turner JH, Trone T, Chandra RK. Differential olfactory outcomes in COVID-19: A large healthcare system population study. *Int Forum Allergy Rhinol* 2022; **12**: 108-111 [PMID: [34289262](https://pubmed.ncbi.nlm.nih.gov/34289262/) DOI: [10.1002/alr.22870](https://doi.org/10.1002/alr.22870)]
- 51 **Galougahi MK**, Ghorbani J, Bakhshayeshkaram M, Naeini AS, Haseli S. Olfactory Bulb Magnetic Resonance Imaging in SARS-CoV-2-Induced Anosmia: The First Report. *Acad Radiol* 2020; **27**: 892-893 [PMID: [32295727](https://pubmed.ncbi.nlm.nih.gov/32295727/) DOI: [10.1016/j.acra.2020.04.002](https://doi.org/10.1016/j.acra.2020.04.002)]
- 52 **Eliezer M**, Hautefort C, Hamel AL, Verillaud B, Herman P, Houdart E, Eloit C. Sudden and Complete Olfactory Loss of Function as a Possible Symptom of COVID-19. *JAMA Otolaryngol Head Neck Surg* 2020; **146**: 674-675 [PMID: [32267483](https://pubmed.ncbi.nlm.nih.gov/32267483/) DOI: [10.1001/jamaoto.2020.0832](https://doi.org/10.1001/jamaoto.2020.0832)]
- 53 **Laurendon T**, Radulesco T, Mugnier J, Gérault M, Chagnaud C, El Ahmadi AA, Varoquaux A. Bilateral transient olfactory bulb edema during COVID-19-related anosmia. *Neurology* 2020; **95**: 224-225 [PMID: [32444492](https://pubmed.ncbi.nlm.nih.gov/32444492/) DOI: [10.1212/WNL.0000000000009850](https://doi.org/10.1212/WNL.0000000000009850)]
- 54 **Strauss SB**, Lantos JE, Heier LA, Shatzkes DR, Phillips CD. Olfactory Bulb Signal Abnormality in Patients with COVID-19 Who Present with Neurologic Symptoms. *AJNR Am J Neuroradiol* 2020; **41**: 1882-1887 [PMID: [32855190](https://pubmed.ncbi.nlm.nih.gov/32855190/) DOI: [10.3174/ajnr.A6751](https://doi.org/10.3174/ajnr.A6751)]
- 55 **Rombaux P**, Duprez T, Hummel T. Olfactory bulb volume in the clinical assessment of olfactory dysfunction. *Rhinology* 2009; **47**: 3-9 [PMID: [19382487](https://pubmed.ncbi.nlm.nih.gov/19382487/)]

- 56 **Corrêa DG**, Hygino da Cruz LC Jr, Lopes FCR, Rangel CC, de Araújo Henriques Tessorollo AL, Coelho KCG, Brandão RZ, Novis RAF, Novis SSC, Silveira R, Carvalho VT, Nascimento OJM. Magnetic resonance imaging features of COVID-19-related cranial nerve lesions. *J Neurovirol* 2021; **27**: 171-177 [PMID: [33460014](#) DOI: [10.1007/s13365-020-00934-0](#)]
- 57 **Aragão MFVV**, Leal MC, Cartaxo Filho OQ, Fonseca TM, Valença MM. Anosmia in COVID-19 Associated with Injury to the Olfactory Bulbs Evident on MRI. *AJNR Am J Neuroradiol* 2020; **41**: 1703-1706 [PMID: [32586960](#) DOI: [10.3174/ajnr.A6675](#)]
- 58 **Hatipoglu N**, Yazici ZM, Palabiyik F, Gulustan F, Sayin I. Olfactory bulb magnetic resonance imaging in SARS-CoV-2-induced anosmia in pediatric cases. *Int J Pediatr Otorhinolaryngol* 2020; **139**: 110469 [PMID: [33120100](#) DOI: [10.1016/j.ijporl.2020.110469](#)]
- 59 **Kandemirli SG**, Altundag A, Yildirim D, Tekcan Sanli DE, Saatci O. Olfactory Bulb MRI and Paranasal Sinus CT Findings in Persistent COVID-19 Anosmia. *Acad Radiol* 2021; **28**: 28-35 [PMID: [33132007](#) DOI: [10.1016/j.acra.2020.10.006](#)]
- 60 **Yildirim D**, Kandemirli SG, Tekcan Sanli DE, Akinci O, Altundag A. A Comparative Olfactory MRI, DTI and fMRI Study of COVID-19 Related Anosmia and Post Viral Olfactory Dysfunction. *Acad Radiol* 2022; **29**: 31-41 [PMID: [34810059](#) DOI: [10.1016/j.acra.2021.10.019](#)]
- 61 **Desforges M**, Favreau DJ, Brison É, Desjardins J, Meessen-Pinard M, Jacomy H, Talbot PJ. Human coronavirus: respiratory pathogens revisited as infectious neuroinvasive, neurotropic, and neurovirulent agents. In: Singh SK, Daniel Ruzek D. Neuroviral infections: RNA viruses and retroviruses. Boca Raton: CRC Press, 2013
- 62 **Khaitovich AB**, Sataieva TP, Sheyko EA, Zukow W. Khaitovich AB, Sataieva TP, Sheyko EA, Zukow W. The analysis of biological diversity of coronaviruses contributes in the early awareness of their zoonotic spreading. *Ecological Questions* 2021; **32**: 67-75 [DOI: [10.12775/eq.2021.025](#)]
- 63 **Sasinka M**, Plenta I, Pavlovic M, Izakovic V, Kaiserová E, Páleníková O, Stubna J. [Complications of immunosuppressive therapy in chronic nephropathies in children]. *Cesk Pediatr* 1976; **31**: 122-126 [PMID: [1269009](#)]
- 64 **Zaki AM**, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012; **367**: 1814-1820 [PMID: [23075143](#) DOI: [10.1056/NEJMoa1211721](#)]
- 65 **Wu A**, Peng Y, Huang B, Ding X, Wang X, Niu P, Meng J, Zhu Z, Zhang Z, Wang J, Sheng J, Quan L, Xia Z, Tan W, Cheng G, Jiang T. Genome Composition and Divergence of the Novel Coronavirus (2019-nCoV) Originating in China. *Cell Host Microbe* 2020; **27**: 325-328 [PMID: [32035028](#) DOI: [10.1016/j.chom.2020.02.001](#)]
- 66 **Paraskevis D**, Kostaki EG, Magiorkinis G, Panayiotakopoulos G, Sourvinos G, Tsiodras S. Full-genome evolutionary analysis of the novel corona virus (2019-nCoV) rejects the hypothesis of emergence as a result of a recent recombination event. *Infect Genet Evol* 2020; **79**: 104212 [PMID: [32004758](#) DOI: [10.1016/j.meegid.2020.104212](#)]
- 67 **Miura TA**, Holmes KV. Host-pathogen interactions during coronavirus infection of primary alveolar epithelial cells. *J Leukoc Biol* 2009; **86**: 1145-1151 [PMID: [19638499](#) DOI: [10.1189/jlb.0209078](#)]
- 68 **Perlman S**, Jacobsen G, Afifi A. Spread of a neurotropic murine coronavirus into the CNS via the trigeminal and olfactory nerves. *Virology* 1989; **170**: 556-560 [PMID: [2543129](#) DOI: [10.1016/0042-6822\(89\)90446-7](#)]
- 69 **Hara Y**, Hasebe R, Sunden Y, Ochiai K, Honda E, Sakoda Y, Umemura T. Propagation of swine hemagglutinating encephalomyelitis virus and pseudorabies virus in dorsal root ganglia cells. *J Vet Med Sci* 2009; **71**: 595-601 [PMID: [19498285](#) DOI: [10.1292/jvms.71.595](#)]
- 70 **Andries K**, Pensaert MB. Immunofluorescence studies on the pathogenesis of hemagglutinating encephalomyelitis virus infection in pigs after oronasal inoculation. *Am J Vet Res* 1980; **41**: 1372-1378 [PMID: [6255837](#)]
- 71 **Koyuncu OO**, Hogue IB, Enquist LW. Virus infections in the nervous system. *Cell Host Microbe* 2013; **13**: 379-393 [PMID: [23601101](#) DOI: [10.1016/j.chom.2013.03.010](#)]
- 72 **Pedersen M**, Sakakura Y, Winther B, Brofeldt S, Mygind N. Nasal mucociliary transport, number of ciliated cells, and beating pattern in naturally acquired common colds. *Eur J Respir Dis Suppl* 1983; **128** (Pt 1): 355-365 [PMID: [6578083](#)]
- 73 **Dubé M**, Le Coupanec A, Wong AHM, Rini JM, Desforges M, Talbot PJ. Axonal Transport Enables Neuron-to-Neuron Propagation of Human Coronavirus OC43. *J Virol* 2018; **92** [PMID: [29925652](#) DOI: [10.1128/JVI.00404-18](#)]
- 74 **Netland J**, Meyerholz DK, Moore S, Cassell M, Perlman S. Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol* 2008; **82**: 7264-7275 [PMID: [18495771](#) DOI: [10.1128/JVI.00737-08](#)]
- 75 **Li K**, Wohlford-Lenane C, Perlman S, Zhao J, Jewell AK, Reznikov LR, Gibson-Corley KN, Meyerholz DK, McCray PB Jr. Middle East Respiratory Syndrome Coronavirus Causes Multiple Organ Damage and Lethal Disease in Mice Transgenic for Human Dipeptidyl Peptidase 4. *J Infect Dis* 2016; **213**: 712-722 [PMID: [26486634](#) DOI: [10.1093/infdis/jiv499](#)]
- 76 **Bryche B**, St Albin A, Murri S, Lacôte S, Pulido C, Ar Gouilh M, Lesellier S, Servat A, Wasniewski M, Picard-Meyer E, Monchatre-Leroy E, Volmer R, Rampin O, Le Goffic R, Marianneau P, Meunier N. Massive transient damage of the olfactory epithelium associated with infection of sustentacular cells by SARS-CoV-2 in golden Syrian hamsters. *Brain Behav Immun* 2020; **89**: 579-586 [PMID: [32629042](#) DOI: [10.1016/j.bbi.2020.06.032](#)]
- 77 **Yinda CK**, Port JR, Bushmaker T, Offei Owusu I, Purushotham JN, Avanzato VA, Fischer RJ, Schulz JE, Holbrook MG, Hebner MJ, Rosenke R, Thomas T, Marzi A, Best SM, de Wit E, Shaia C, van Doremalen N, Munster VJ. K18-hACE2 mice develop respiratory disease resembling severe COVID-19. *PLoS Pathog* 2021; **17**: e1009195 [PMID: [33465158](#) DOI: [10.1371/journal.ppat.1009195](#)]
- 78 **Xu H**, Zhong L, Deng J, Peng J, Dan H, Zeng X, Li T, Chen Q. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci* 2020; **12**: 8 [PMID: [32094336](#) DOI: [10.1038/s41368-020-0074-x](#)]
- 79 **Bleau C**, Filliol A, Samson M, Lamontagne L. Brain Invasion by Mouse Hepatitis Virus Depends on Impairment of Tight Junctions and Beta Interferon Production in Brain Microvascular Endothelial Cells. *J Virol* 2015; **89**: 9896-9908 [PMID: [26202229](#) DOI: [10.1128/JVI.01501-15](#)]
- 80 **Alexopoulos H**, Magira E, Bitzogli K, Kafasi N, Vlachoyiannopoulos P, Tzioufas A, Kotanidou A, Dalakas MC. Anti-SARS-CoV-2 antibodies in the CSF, blood-brain barrier dysfunction, and neurological outcome: Studies in 8 stuporous and comatose patients. *Neurol Neuroimmunol Neuroinflamm* 2020; **7** [PMID: [32978291](#) DOI: [10.1212/NXI.0000000000000500](#)]

- 10.1212/NXI.0000000000000893]
- 81 **Ding Y**, He L, Zhang Q, Huang Z, Che X, Hou J, Wang H, Shen H, Qiu L, Li Z, Geng J, Cai J, Han H, Li X, Kang W, Weng D, Liang P, Jiang S. Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus transmission pathways. *J Pathol* 2004; **203**: 622-630 [PMID: [15141376](#) DOI: [10.1002/path.1560](#)]
- 82 **Steardo L**, Steardo L Jr, Zorec R, Verkhatsky A. Neuroinfection may contribute to pathophysiology and clinical manifestations of COVID-19. *Acta Physiol (Oxf)* 2020; **229**: e13473 [PMID: [32223077](#) DOI: [10.1111/apha.13473](#)]
- 83 **Coolen T**, Lolli V, Sadeghi N, Rovai A, Trotta N, Taccone FS, Creteur J, Henrard S, Goffard JC, Dewitte O, Naeije G, Goldman S, De Tiège X. Early postmortem brain MRI findings in COVID-19 non-survivors. *Neurology* 2020; **95**: e2016-e2027 [PMID: [32546654](#) DOI: [10.1212/WNL.00000000000010116](#)]
- 84 **Solomon IH**, Normandin E, Bhattacharyya S, Mukerji SS, Keller K, Ali AS, Adams G, Hornick JL, Padera RF Jr, Sabeti P. Neuropathological Features of Covid-19. *N Engl J Med* 2020; **383**: 989-992 [PMID: [32530583](#) DOI: [10.1056/NEJMc2019373](#)]
- 85 **Lee MH**, Perl DP, Nair G, Li W, Maric D, Murray H, Dodd SJ, Koretsky AP, Watts JA, Cheung V, Masliah E, Horkayne-Szakaly I, Jones R, Stram MN, Moncur J, Hefti M, Folkner RD, Nath A. Microvascular Injury in the Brains of Patients with Covid-19. *N Engl J Med* 2021; **384**: 481-483 [PMID: [33378608](#) DOI: [10.1056/NEJMc2033369](#)]
- 86 **Brian DA**, Baric RS. Coronavirus genome structure and replication. *Curr Top Microbiol Immunol* 2005; **287**: 1-30 [PMID: [15609507](#) DOI: [10.1007/3-540-26765-4_1](#)]
- 87 **Ziegler CGK**, Allon SJ, Nyquist SK, Mbano IM, Miao VN, Tzouanas CN, Cao Y, Yousif AS, Bals J, Hauser BM, Feldman J, Muus C, Wadsworth MH 2nd, Kazer SW, Hughes TK, Doran B, Gatter GJ, Vukovic M, Taliaferro F, Mead BE, Guo Z, Wang JP, Gras D, Plaisant M, Ansari M, Angelidis I, Adler H, Sucre JMS, Taylor CJ, Lin B, Waghray A, Mitsialis V, Dwyer DF, Buchheit KM, Boyce JA, Barrett NA, Laidlaw TM, Carroll SL, Colonna L, Tkachev V, Peterson CW, Yu A, Zheng HB, Gideon HP, Winchell CG, Lin PL, Bingle CD, Snapper SB, Kropski JA, Theis FJ, Schiller HB, Zaragosi LE, Barbry P, Leslie A, Kiem HP, Flynn JL, Fortune SM, Berger B, Finberg RW, Kean LS, Garber M, Schmidt AG, Lingwood D, Shalek AK, Ordoñez-Montanes J; HCA Lung Biological Network. Lung Biological Network. SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell* 2020; **181**: 1016-1035.e19 [PMID: [32413319](#) DOI: [10.1016/j.cell.2020.04.035](#)]
- 88 **Hamming I**, Timens W, Bulthuis ML, Lely AT, Navis G, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol* 2004; **203**: 631-637 [PMID: [15141377](#) DOI: [10.1002/path.1570](#)]
- 89 **Li W**, Zhang C, Sui J, Kuhn JH, Moore MJ, Luo S, Wong SK, Huang IC, Xu K, Vasilieva N, Murakami A, He Y, Marasco WA, Guan Y, Choe H, Farzan M. Receptor and viral determinants of SARS-coronavirus adaptation to human ACE2. *EMBO J* 2005; **24**: 1634-1643 [PMID: [15791205](#) DOI: [10.1038/sj.emboj.7600640](#)]
- 90 **Sia SF**, Yan LM, Chin AWH, Fung K, Choy KT, Wong AYL, Kaewpreedee P, Perera RAPM, Poon LLM, Nicholls JM, Peiris M, Yen HL. Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature* 2020; **583**: 834-838 [PMID: [32408338](#) DOI: [10.1038/s41586-020-2342-5](#)]
- 91 **Zazhytska M**, Kodra A, Hoagland DA, Frere J, Fullard JF, Shayya H, McArthur NG, Moeller R, Uhl S, Omer AD, Gottesman ME, Firestein S, Gong Q, Canoll PD, Goldman JE, Roussos P, tenOever BR, Jonathan B Overdevest, Lomvardas S. Non-cell-autonomous disruption of nuclear architecture as a potential cause of COVID-19-induced anosmia. *Cell* 2022; **185**: 1052-1064.e12 [PMID: [35180380](#) DOI: [10.1016/j.cell.2022.01.024](#)]
- 92 **Al-Sarraj S**, Troakes C, Hanley B, Osborn M, Richardson MP, Hotopf M, Bullmore E, Everall IP. Invited Review: The spectrum of neuropathology in COVID-19. *Neuropathol Appl Neurobiol* 2021; **47**: 3-16 [PMID: [32935873](#) DOI: [10.1111/nan.12667](#)]
- 93 **Romoli M**, Jelcic I, Bernard-Valnet R, García Azorín D, Mancinelli L, Akhvlediani T, Monaco S, Taba P, Sellner J; Infectious Disease Panel of the European Academy of Neurology. A systematic review of neurological manifestations of SARS-CoV-2 infection: the devil is hidden in the details. *Eur J Neurol* 2020; **27**: 1712-1726 [PMID: [32503088](#) DOI: [10.1111/ene.14382](#)]
- 94 **Spudich S**, Nath A. Nervous system consequences of COVID-19. *Science* 2022; **375**: 267-269 [PMID: [35050660](#) DOI: [10.1126/science.abm2052](#)]
- 95 **Mehta P**, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ; HLH Across Speciality Collaboration, UK. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020; **395**: 1033-1034 [PMID: [32192578](#) DOI: [10.1016/S0140-6736\(20\)30628-0](#)]
- 96 **Sharifi-Razavi A**, Karimi N, Rouhani N. COVID-19 and intracerebral haemorrhage: causative or coincidental? *New Microbes New Infect* 2020; **35**: 100669 [PMID: [32322398](#) DOI: [10.1016/j.nmni.2020.100669](#)]
- 97 **Letko M**, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol* 2020; **5**: 562-569 [PMID: [32094589](#) DOI: [10.1038/s41564-020-0688-y](#)]
- 98 **Zhang H**, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med* 2020; **46**: 586-590 [PMID: [32125455](#) DOI: [10.1007/s00134-020-05985-9](#)]
- 99 **Bilinska K**, Jakubowska P, Von Bartheld CS, Butowt R. Expression of the SARS-CoV-2 Entry Proteins, ACE2 and TMPRSS2, in Cells of the Olfactory Epithelium: Identification of Cell Types and Trends with Age. *ACS Chem Neurosci* 2020; **11**: 1555-1562 [PMID: [32379417](#) DOI: [10.1021/acscchemneuro.0c00210](#)]
- 100 **Hummel T**, Whitcroft KL, Andrews P, Altundag A, Cinghi C, Costanzo RM, Damm M, Frasnelli J, Gudziol H, Gupta N, Haehner A, Holbrook E, Hong SC, Hornung D, Hüttenbrink KB, Kamel R, Kobayashi M, Konstantinidis I, Landis BN, Leopold DA, Macchi A, Miwa T, Moesges R, Mullol J, Mueller CA, Ottaviano G, Passali GC, Philpott C, Pinto JM, Ramakrishnan VJ, Rombaux P, Roth Y, Schlosser RA, Shu B, Soler G, Stjärne P, Stuck BA, Vodicka J, Welge-Lüssen A. Position paper on olfactory dysfunction. *Rhinology* 2016; **56**: 1-30 [PMID: [28623665](#) DOI: [10.4193/Rhino16.248](#)]
- 101 **Hildenbrand T**, Weber RK, Brehmer D. Rhinitis sicca, dry nose and atrophic rhinitis: a review of the literature. *Eur Arch Otorhinolaryngol* 2011; **268**: 17-26 [PMID: [20878413](#) DOI: [10.1007/s00405-010-1391-z](#)]

- 102 **Thakur KT**, Miller EH, Glendinning MD, Al-Dalahmah O, Banu MA, Boehme AK, Boubour AL, Bruce SS, Chong AM, Claassen J, Faust PL, Hargus G, Hickman RA, Jambawalikar S, Khandji AG, Kim CY, Klein RS, Lignelli-Dipple A, Lin CC, Liu Y, Miller ML, Moonis G, Nordvig AS, Overdeest JB, Prust ML, Przedborski S, Roth WH, Soung A, Tanji K, Teich AF, Agalliu D, Uhlemann AC, Goldman JE, Canoll P. COVID-19 neuropathology at Columbia University Irving Medical Center/New York Presbyterian Hospital. *Brain* 2021; **144**: 2696-2708 [PMID: [33856027](#) DOI: [10.1093/brain/awab148](#)]
- 103 **Wong ST**, Trinh K, Hacker B, Chan GC, Lowe G, Gaggari A, Xia Z, Gold GH, Storm DR. Disruption of the type III adenylyl cyclase gene leads to peripheral and behavioral anosmia in transgenic mice. *Neuron* 2000; **27**: 487-497 [PMID: [11055432](#) DOI: [10.1016/s0896-6273\(00\)00060-x](#)]
- 104 **Saito H**, Kubota M, Roberts RW, Chi Q, Matsunami H. RTP family members induce functional expression of mammalian odorant receptors. *Cell* 2004; **119**: 679-691 [PMID: [15550249](#) DOI: [10.1016/j.cell.2004.11.021](#)]
- 105 **Brunet LJ**, Gold GH, Ngai J. General anosmia caused by a targeted disruption of the mouse olfactory cyclic nucleotide-gated cation channel. *Neuron* 1996; **17**: 681-693 [PMID: [8893025](#) DOI: [10.1016/s0896-6273\(00\)80200-7](#)]
- 106 **Sanz G**, Thomas-Danguin T, Hamdani el H, Le Poupon C, Briand L, Pernollet JC, Guichard E, Tromelin A. Relationships between molecular structure and perceived odor quality of ligands for a human olfactory receptor. *Chem Senses* 2008; **33**: 639-653 [PMID: [18603653](#) DOI: [10.1093/chemse/bjn032](#)]
- 107 **Chen L**, Zhao J, Peng J, Li X, Deng X, Geng Z, Shen Z, Guo F, Zhang Q, Jin Y, Wang L, Wang S. Detection of SARS-CoV-2 in saliva and characterization of oral symptoms in COVID-19 patients. *Cell Prolif* 2020; **53**: e12923 [PMID: [33073910](#) DOI: [10.1111/cpr.12923](#)]
- 108 **Doyle ME**, Appleton A, Liu QR, Yao Q, Mazucanti CH, Egan JM. Human Taste Cells Express ACE2: a Portal for SARS-CoV-2 Infection. *bioRxiv* 2021 [PMID: [33907747](#) DOI: [10.1101/2021.04.21.440680](#)]
- 109 **Zhu F**, Zhong Y, Ji H, Ge R, Guo L, Song H, Wu H, Jiao P, Li S, Wang C, Du H. ACE2 and TMPRSS2 in human saliva can adsorb to the oral mucosal epithelium. *J Anat* 2022; **240**: 398-409 [PMID: [34590312](#) DOI: [10.1111/joa.13560](#)]
- 110 **Wang H**, Zhou M, Brand J, Huang L. Inflammation activates the interferon signaling pathways in taste bud cells. *J Neurosci* 2007; **27**: 10703-10713 [PMID: [17913904](#) DOI: [10.1523/JNEUROSCI.3102-07.2007](#)]
- 111 **Pushpass RG**, Pellicciotta N, Kelly C, Proctor G, Carpenter GH. Reduced Salivary Mucin Binding and Glycosylation in Older Adults Influences Taste in an In Vitro Cell Model. *Nutrients* 2019; **11** [PMID: [31554163](#) DOI: [10.3390/nu11102280](#)]
- 112 **van Tongeren J**, Golebski K, Van Egmond D, de Groot EJ, Fokkens WJ, van Drunen CM. Synergy between TLR-2 and TLR-3 signaling in primary human nasal epithelial cells. *Immunobiology* 2015; **220**: 445-451 [PMID: [25532794](#) DOI: [10.1016/j.imbio.2014.11.004](#)]
- 113 **Gadanec LK**, McSweeney KR, Qaradakh T, Ali B, Zulli A, Apostolopoulos V. Can SARS-CoV-2 Virus Use Multiple Receptors to Enter Host Cells? *Int J Mol Sci* 2021; **22** [PMID: [33498183](#) DOI: [10.3390/ijms22030992](#)]
- 114 **Milanetti E**, Miotto M, Di Rienzo L, Nagaraj M, Monti M, Golbek TW, Gosti G, Roeters SJ, Weidner T, Otzen DE, Ruocco G. In-Silico Evidence for a Two Receptor Based Strategy of SARS-CoV-2. *Front Mol Biosci* 2021; **8**: 690655 [PMID: [34179095](#) DOI: [10.3389/fmolb.2021.690655](#)]
- 115 **Li W**, Hulswit RJG, Widjaja I, Raj VS, McBride R, Peng W, Widagdo W, Tortorici MA, van Dieren B, Lang Y, van Lent JWM, Paulson JC, de Haan CAM, de Groot RJ, van Kuppeveld FJM, Haagmans BL, Bosch BJ. Identification of sialic acid-binding function for the Middle East respiratory syndrome coronavirus spike glycoprotein. *Proc Natl Acad Sci U S A* 2017; **114**: E8508-E8517 [PMID: [28923942](#) DOI: [10.1073/pnas.1712592114](#)]
- 116 **Fantozzi PJ**, Pampena E, Di Vanna D, Pellegrino E, Corbi D, Mammucari S, Alessi F, Pampena R, Bertazzoni G, Minisola S, Mastroianni CM, Polimeni A, Romeo U, Villa A. Xerostomia, gustatory and olfactory dysfunctions in patients with COVID-19. *Am J Otolaryngol* 2020; **41**: 102721 [PMID: [32977063](#) DOI: [10.1016/j.amjoto.2020.102721](#)]
- 117 **Fathi Y**, Hoseini EG, Atoof F, Mottaghi R. Xerostomia (dry mouth) in patients with COVID-19: a case series. *Future Virol* 2021; **16** [DOI: [10.2217/fvl-2020-0334](#)]
- 118 **Lozada-Nur F**, Chainani-Wu N, Fortuna G, Sroussi H. Dysgeusia in COVID-19: Possible Mechanisms and Implications. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2020; **130**: 344-346 [PMID: [32703719](#) DOI: [10.1016/j.oooo.2020.06.016](#)]
- 119 **Choudhury A**, Mukherjee S. In silico studies on the comparative characterization of the interactions of SARS-CoV-2 spike glycoprotein with ACE-2 receptor homologs and human TLRs. *J Med Virol* 2020; **92**: 2105-2113 [PMID: [32383269](#) DOI: [10.1002/jmv.25987](#)]
- 120 **Park YJ**, Choe YJ, Park O, Park SY, Kim YM, Kim J, Kweon S, Woo Y, Gwack J, Kim SS, Lee J, Hyun J, Ryu B, Jang YS, Kim H, Shin SH, Yi S, Lee S, Kim HK, Lee H, Jin Y, Park E, Choi SW, Kim M, Song J, Kim D, Jeon BH, Yoo H, Jeong EK; COVID-19 National Emergency Response Center, Epidemiology and Case Management Team. Contact Tracing during Coronavirus Disease Outbreak, South Korea, 2020. *Emerg Infect Dis* 2020; **26**: 2465-2468 [PMID: [32673193](#) DOI: [10.3201/eid2610.201315](#)]
- 121 **Yang X**, Yu Y, Xu J, Shu H, Xia J, Liu H, Wu Y, Zhang L, Yu Z, Fang M, Yu T, Wang Y, Pan S, Zou X, Yuan S, Shang Y. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020; **8**: 475-481 [PMID: [32105632](#) DOI: [10.1016/S2213-2600\(20\)30079-5](#)]
- 122 **Stewart RE**, DeSimone JA, Hill DL. New perspectives in a gustatory physiology: transduction, development, and plasticity. *Am J Physiol* 1997; **272**: C1-26 [PMID: [9038806](#) DOI: [10.1152/ajpcell.1997.272.1.C1](#)]
- 123 **Smith RL**, Traul DL, Schaack J, Clayton GH, Staley KJ, Wilcox CL. Characterization of promoter function and cell-type-specific expression from viral vectors in the nervous system. *J Virol* 2000; **74**: 11254-11261 [PMID: [11070024](#) DOI: [10.1128/jvi.74.23.11254-11261.2000](#)]
- 124 **Bagheri S**, Ghobadimoghadam S. Safety and Health Protection of Health Care Workers during the COVID-19 Pandemic. *Int J Community Based Nurs Midwifery* 2020; **8**: 362-363 [PMID: [33178860](#) DOI: [10.30476/IJCBNM.2020.86066.1319](#)]
- 125 **Khani E**, Khiali S, Beheshtirouy S, Entezari-Maleki T. Potential pharmacologic treatments for COVID-19 smell and taste loss: A comprehensive review. *Eur J Pharmacol* 2021; **912**: 174582 [PMID: [34678243](#) DOI: [10.1016/j.ejphar.2021.174582](#)]

- 126 **Webster KE**, O'Byrne L, MacKeith S, Philpott C, Hopkins C, Burton MJ. Interventions for the prevention of persistent post-COVID-19 olfactory dysfunction. *Cochrane Database Syst Rev* 2021; **7**: CD013877 [PMID: [34291812](#) DOI: [10.1002/14651858.CD013877.pub2](#)]
- 127 **Le Bon SD**, Konopnicki D, Pisarski N, Prunier L, Lechien JR, Horoi M. Efficacy and safety of oral corticosteroids and olfactory training in the management of COVID-19-related loss of smell. *Eur Arch Otorhinolaryngol* 2021; **278**: 3113-3117 [PMID: [33423106](#) DOI: [10.1007/s00405-020-06520-8](#)]
- 128 **Ivory K**, Prieto E, Spinks C, Armah CN, Goldson AJ, Dainty JR, Nicoletti C. Selenium supplementation has beneficial and detrimental effects on immunity to influenza vaccine in older adults. *Clin Nutr* 2017; **36**: 407-415 [PMID: [26803169](#) DOI: [10.1016/j.clnu.2015.12.003](#)]
- 129 **Iovino L**, Mazziotta F, Carulli G, Guerrini F, Morganti R, Mazzotti V, Maggi F, Macera L, Orciuolo E, Buda G, Benedetti E, Caracciolo F, Galimberti S, Pistello M, Petrini M. High-dose zinc oral supplementation after stem cell transplantation causes an increase of TRECs and CD4+ naïve lymphocytes and prevents TTV reactivation. *Leuk Res* 2018; **70**: 20-24 [PMID: [29747074](#) DOI: [10.1016/j.leukres.2018.04.016](#)]
- 130 **Wintergerst ES**, Maggini S, Hornig DH. Contribution of selected vitamins and trace elements to immune function. *Ann Nutr Metab* 2007; **51**: 301-323 [PMID: [17726308](#) DOI: [10.1159/000107673](#)]
- 131 **Patel A**, Jernigan DB; 2019-nCoV CDC Response Team. Initial Public Health Response and Interim Clinical Guidance for the 2019 Novel Coronavirus Outbreak - United States, December 31, 2019-February 4, 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 140-146 [PMID: [32027631](#) DOI: [10.15585/mmwr.mm6905e1](#)]
- 132 **Mousa HA**. Prevention and Treatment of Influenza, Influenza-Like Illness, and Common Cold by Herbal, Complementary, and Natural Therapies. *J Evid Based Complementary Altern Med* 2017; **22**: 166-174 [PMID: [27055821](#) DOI: [10.1177/2156587216641831](#)]
- 133 **Seden N**. Loss of Smell and Taste as Clinical Onset of COVID-19. In: Baddour MM. Fighting the COVID-19 Pandemic. London: IntechOpen Limited, 2021 [DOI: [10.5772/intechopen.99087](#)]
- 134 **Yaylaci A**, Azak E, Önal A, Aktürk DR, Karadenizli A. Effects of classical olfactory training in patients with COVID-19-related persistent loss of smell. *Eur Arch Otorhinolaryngol* 2023; **280**: 757-763 [PMID: [35904631](#) DOI: [10.1007/s00405-022-07570-w](#)]
- 135 **Hemilä H**. Zinc lozenges may shorten the duration of colds: a systematic review. *Open Respir Med J* 2011; **5**: 51-58 [PMID: [21769305](#) DOI: [10.2174/1874306401105010051](#)]
- 136 **Read SA**, Obeid S, Ahlenstiel C, Ahlenstiel G. The Role of Zinc in Antiviral Immunity. *Adv Nutr* 2019; **10**: 696-710 [PMID: [31305906](#) DOI: [10.1093/advances/nmz013](#)]
- 137 **Heckmann SM**, Hujoel P, Habiger S, Friess W, Wichmann M, Heckmann JG, Hummel T. Zinc gluconate in the treatment of dysgeusia--a randomized clinical trial. *J Dent Res* 2005; **84**: 35-38 [PMID: [15615872](#) DOI: [10.1177/154405910508400105](#)]
- 138 **te Velthuis AJ**, van den Worm SH, Sims AC, Baric RS, Snijder EJ, van Hemert MJ. Zn(2+) inhibits coronavirus and arterivirus RNA polymerase activity in vitro and zinc ionophores block the replication of these viruses in cell culture. *PLoS Pathog* 2010; **6**: e1001176 [PMID: [21079686](#) DOI: [10.1371/journal.ppat.1001176](#)]
- 139 **Beigmohammadi MT**, Bitarafan S, Hoseindokht A, Abdollahi A, Amoozadeh L, Soltani D. The effect of supplementation with vitamins A, B, C, D, and E on disease severity and inflammatory responses in patients with COVID-19: a randomized clinical trial. *Trials* 2021; **22**: 802 [PMID: [34776002](#) DOI: [10.1186/s13063-021-05795-4](#)]
- 140 **Díaz-Resendiz KJG**, Benitez-Trinidad AB, Covantes-Rosales CE, Toledo-Ibarra GA, Ortiz-Lazareno PC, Girón-Pérez DA, Bueno-Durán AY, Pérez-Díaz DA, Barcelos-García RG, Girón-Pérez MI. Loss of mitochondrial membrane potential ($\Delta\Psi(m)$) in leucocytes as post-COVID-19 sequelae. *J Leukoc Biol* 2022; **112**: 23-29 [PMID: [35355308](#) DOI: [10.1002/JLB.3MA0322-279RRR](#)]
- 141 **Gudziol V**, Hummel T. Effects of pentoxifylline on olfactory sensitivity: a postmarketing surveillance study. *Arch Otolaryngol Head Neck Surg* 2009; **135**: 291-295 [PMID: [19289709](#) DOI: [10.1001/archoto.2008.524](#)]



Transoral robotic surgery for adult parapharyngeal lymphangioma: A case report

Surender Dabas, Nandini N Menon, Reetesh Ranjan, Bikas Gurung, Himanshu Shukla, Ashwani K Sharma, Sukirti Tiwari, Ajit Sinha, Sahibinder Singh Bhatti, Rishu Sangal

Specialty type:

Otorhinolaryngology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model:

Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: He YF, China; Maurea S, Italy

Received: October 13, 2022

Peer-review started: October 13, 2022

First decision: December 13, 2022

Revised: December 17, 2022

Accepted: February 9, 2023

Article in press: February 9, 2023

Published online: May 9, 2023



Surender Dabas, Nandini N Menon, Reetesh Ranjan, Bikas Gurung, Himanshu Shukla, Ashwani K Sharma, Sukirti Tiwari, Ajit Sinha, Department of Surgical Oncology, BLK MAX Superspeciality Hospital, Delhi 110007, India

Sahibinder Singh Bhatti, Department of Pathology, BLK MAX Superspeciality Hospital, Delhi 110007, India

Rishu Sangal, Department of Radiology, BLK MAX Superspeciality Hospital, Delhi 110007, India

Corresponding author: Nandini N Menon, MBBS, MS, Attending Doctor, Department of Surgical oncology, BLK MAX Superspeciality Hospital, Karol Bagh, Delhi 110007, India. menonnandini1602@gmail.com

Abstract

BACKGROUND

Lymphangiomas are a group of benign swellings which are commonly seen in children. The most common sites of presentation is the head and neck region, less commonly seen in axilla, chest, liver, spleen, etc. The ideal modality of treatment has always been surgical excision irrespective of the site and age group. But with the advent of minimally invasive surgical techniques, it is now possible to perform excision of parapharyngeal space lesions with minimal morbidity and good clearance.

CASE SUMMARY

A 42-year-old male patient who presented with difficulty in swallowing and had undergone surgery twice outside, where Transcervical approach was attempted to remove the parapharyngeal mass, but failed. Magnetic resonance imaging scan demonstrated a 6 cm x 5 cm x 4 cm left parapharyngeal mass. He underwent transoral robotic surgery for the excision of the parapharyngeal lesion and had an uneventful post-operative recovery.

CONCLUSION

Lymphangiomas are hamartomatous swellings which are benign in nature. The symptoms of the patient with large parapharyngeal mass include dysphagia, dyspnoea and neck swelling. Clinoradiological evaluation is of utmost importance to determine the adjacent vital structures and the approach to the tumor. With the advent of robotics in oncology, transoral robotic excision is one of the

best approaches to perform such a surgery.

Key Words: Lymphangioma; Parapharyngeal space lesions; Transoral robotic surgery for parapharyngeal space mass; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This report describes a rare case of adult parapharyngeal space lymphangioma. It also emphasizes the role of minimally invasive techniques like transoral robotic surgery for the excision of parapharyngeal space tumors which has resulted in less morbidity in the patients and gives good clearance of tumor.

Citation: Dabas S, Menon NN, Ranjan R, Gurung B, Shukla H, Sharma AK, Tiwari S, Sinha A, Bhatti SS, Sangal R. Transoral robotic surgery for adult parapharyngeal lymphangioma: A case report. *World J Otorhinolaryngol* 2023; 10(2): 23-29

URL: <https://www.wjgnet.com/2218-6247/full/v10/i2/23.htm>

DOI: <https://dx.doi.org/10.5319/wjo.v10.i2.23>

INTRODUCTION

Cystic hygroma, also known as cystic lymphangioma, jugular lymphatic obstructive sequence and hygromaccollicysticum, was first described in 1828 by Redenbacher[1]. Only 6% of benign lumps in patients aged less than 20 years are attributable to lymphangiomas[2]. Parapharyngeal space is an extremely rare subsite for lymphangiomas in the head and neck. Other sites include axilla, mediastinum, groin and abdominal cavity[3].

Different etiological and pathogenic factors have been considered causal, including traumatic, infectious and chronic inflammatory factors, as well as alterations in embryological development[4]. In adults, late proliferation of cellular nests of the lymphatic system may appear due to stimulation caused by trauma or infection[5].

Lymphangiomas in the head and neck region commonly develop as an asymptomatic neck mass around 2 years of age.

The imaging modality preferred for lymphangiomas include ultrasound (USG) supplemented by computerized tomography (CT) and magnetic resonance imaging (MRI) scans for obtaining more information about the structural features, internal as well as peripheral enhancement patterns[6].

The most common treatment modality includes surgical excision of the mass. The other treatment modalities include aspiration, sclerotherapy, steroids, laser and radiotherapy.

We here describe a rare case of adult parapharyngeal space lymphangioma which was managed with transoral robotic surgery.

CASE PRESENTATION

Chief complaints

A 42-year-old male patient presented to our outpatient department with chief complaint of difficulty in swallowing both solids and liquids for the past 2 months.

History of present illness

The patient had dysphagia in the past 2 months with no associated odynophagia, respiratory difficulty or neck mass. He had a history of transcervical surgery twice outside for the same reason, which resulted in incomplete removal of the concerned pathology. No surgical records of the previous surgeries were available at the time he presented to us.

History of past illness

The patient had no history of any significant comorbidities.

Physical examination

Vital signs recorded were as follows: temperature 36.5 °C, pulse rate 88/ min, blood pressure 130/80 mmHg and respiratory rate 18 /min. On clinical examination, a submucosal bulge was noted in the left

tonsillar area. No swelling was however palpable in the neck.

Laboratory examinations

Results of routine blood tests as part of pre-operative work-up were within normal limits.

Imaging examinations

Fibreoptic laryngoscopy (FOL) showed a bulge in the region of the left lateral oropharyngeal wall with significant narrowing of the airway. MRI showed a 6 cm x 5 cm x 4 cm lobulated heterogeneous lesion involving the left lateral oropharyngeal wall, parapharyngeal space and peritonsillar area (**Figure 1**). The lesion compressed and displaced the left palatine tonsil anteroinferiorly, laterally reached the deep lobe of parotid gland, abuts medial pterygoid muscle anterolaterally and indents longus coli muscle posteriorly and compressed the external carotid artery. Postoperative changes were noted in the region of the left submandibular gland and anterior to the sternocleidomastoid muscle.

FINAL DIAGNOSIS

The main differential diagnosis was vascular malformation and lymphoreticular malformation. The final diagnosis was established after histopathological examination, after surgery.

TREATMENT

The patient was planned for upfront transoral robotic surgery (TORS) after obtaining anesthesia clearance. A written and informed consent was obtained from the patient before the procedure. The procedure was performed under general anesthesia. Feyh-Kastenbauer retractor was placed and da Vinci Xi robotic system was docked. Three arms of the robot were used for performing the surgery which contained the camera, bipolar and monopolar cautery. After making incision in the peritonsillar area, layer by layer dissection of the mucosa, submucosa and constrictor muscles with adequate hemostasis at each step was performed. A well encapsulated lesion was identified in the left parapharyngeal space which was excised (**Figure 2**) *in toto* and specimen was sent for histopathological examination. There were no intra- or peri-operative complications. The final histopathological report showed a multilobulated lesion with lymph containing sac and the diagnosis was lymphangioma (**Figure 3**).

OUTCOME AND FOLLOW-UP

Postoperatively, the patient stayed in the hospital for 3 days and was administered antibiotics and analgesics. For the initial 1 week, the patient was on Ryle's tube feed. Postoperatively, the patient followed up at the outpatient department on day 7, and fibreoptic laryngoscopy (FOL) showed a healed wound. The patient's speech and swallowing were evaluated. The speech of the patient was normal and his swallowing was also unimpaired. He was then started on liquids followed by soft diet. Normal diet was started on day 15 after surgery. The patient has been on regular follow-up since then for a period of 6 months and has no signs of residual disease, recurrence or any other post-operative complications.

DISCUSSION

Lymphangiomas are defined as benign, congenital and hamartomatous lesions which develop due to abnormalities in the lymphatic system. Their incidence was reported to be 1.2 -2.8 *per* 1000 new births [4]. First 5 years of life show the highest incidence of lymphangiomas (80%-90%) [7].

Histologically, lymphangiomas are classified as capillary, cavernous and cystic hygromas [8]. Based on size, it is divided into microcystic (< 2 cm) and macrocystic types (> 2 cm) [3]. Bhayya *et al* [9] classified them as lymphangioma simplex, cavernous lymphangioma, cystic lymphangioma and benign lymphangioendothelioma [9]. Based on anatomical location, in 1995, de Serres *et al* [10] proposed a classification system as follows: Class I, unilateral infrahyoid lesion; Class II, unilateral suprahyoid lesion; Class III, unilateral combined supra and infrahyoid lesion; Class IV, bilateral suprahyoid lesion, and Class V, bilateral combined supra and infrahyoid lesion [10].

It is reported that 80% of lymphangiomas develop in the neck and 85% are unilateral [4]. Most common presentations include swelling in the neck, respiratory distress, difficulty in swallowing and pain secondary to infection or haemorrhage [7]. Although lymphangiomas are benign lesions with no malignant potential, occasionally it can enlarge to compress and invade the adjacent structures. The




Figure 1 Magnetic resonance images (with contrast and plain) depicting a heterogeneous well enhancing lobulated lesion in the left parapharyngeal space as marked with a red arrow. A: Axial; B: Coronal.





Figure 2 Partly excised parapharyngeal mass visualized with a 30-degree endoscope indicated with a yellow arrow.

latter is a feature of poor prognosis in advanced cases[2]. Clinically, they usually appear as painless and cystic lumps, which are transilluminant in nature. They are associated with pain only when they are complicated with infection or haemorrhage[3].

Lymphangiomas are associated with various congenital syndromes which include trisomies of 13, 18, 21, Ullrich Turner syndrome, Noonan-Ehmke syndrome, aortic coarctation, hypoplastic left heart syndrome, Cantrell Haller Ravitsch syndrome, Apert syndrome, Brachmann-de Lange syndrome, foetal alcohol spectrum disorders, Fryns syndrome, Escobar syndrome, limb hypertrophy, fetal akinesia deformation sequence and hydrops[8].

Imaging is an important diagnostic modality to understand the exact origin, location and extent of lymphangioma. On ultrasound done during the antepartum period, lymphangiomas appear as thin walled multiloculated cysts. CT scan is better compared to USG for imaging lymphangiomas. In CT scans, they appear as low attenuation masses with regular margins and an enhancing rim on intravenous contrast. On T1 weighted images, they appear with low signal compared to muscles and on T2 weighted images, they appear hyperintense compared to fat[4,6].

In addition, 50% of parapharyngeal space tumors have a salivary cause, 20% have a neurogenic root, and 30% are benign and malignant lymphoreticular lesions, metastatic lesions, carotid body tumors and lipomas. Amongst neurogenic tumors, Schwannomas arising from the vagus nerve, presenting as post styloid compartment lesions are the most common[11]. On imaging, homogenous lesions seen displacing the internal carotid or common carotid artery anteriorly is suggestive of schwannoma arising from vagus nerve or sympathetic trunk[12]. Lipomas appear as hyperintense lesions on T1- and T2-weighted images with internal septations[13]. Vascular tumors are generally more conspicuous on T2-weighted images. Pleomorphic adenomas are usually characterized by the smooth and regular, non-



DOI: 10.5319/wjo.v10.i2.23 Copyright ©The Author(s) 2023.

Figure 3 Hematoxylin-eosin staining. A: Hematoxylin-eosin (HE), 40 ×, section examined shows a benign vascular tumour in the soft tissue and skeletal muscle fibres with morphological feature consistent with a lymphangioma; B: HE, 100 ×, section shows vascular spaces lined by a single layer of benign endothelial cells.

infiltrating margins with cystic or necrotic areas[14]. Paragangliomas have a characteristic “salt and pepper” appearance on MRI scan[15]. Dermoid and epidermoid cysts are differentiated from other similar lesions as they often appear as unilocular masses with homogeneous hypoattenuating fluid on CT scan[16].

The selection criteria employed for treatment of lymphangiomas include large size, disfigurement, recurrent infections, dysphagia or respiratory distress. Literature shows spontaneous regression in 15% cases[8]. Surgery forms the mainstay of treatment[17]. Other treatment options include laser excision, radiotherapy, sclerosant injection, unroofing, aspiration or steroids[4].

The decision on choosing the type of approach for surgery depends on the site and nature of the lesion. Traditionally, the transcervical transmandibular approach was observed to give good exposure of the parapharyngeal space[3]. The main disadvantages of this approach include the dysphagia and persistent respiratory difficulties, which occur secondary to the post-operative mucosal edema and loss of pharyngeal innervation[7].

Traditionally, surgeons believed that in case of lesions extending to parapharyngeal space, complete surgical excision was difficult to obtain without damaging nervous and vascular structures and thus they used to resort to alternatives like drainage and sclerotherapy with different substances such as tetracycline, bleomycin and triamcinolone and radiofrequency ablation[18]. Complications and associated morbidity can include infections, tissue necrosis, cranial nerve lesions, vascular thrombosis and even endocrine disorders[19]. But now, with the advent of robotic surgery, these disadvantages have been overcome and it is possible to obtain 100% clearance of lesions in parapharyngeal space without injuring any neural or vascular structure[20].

A surgical algorithm was proposed by Chu *et al*[21] in 2017 for access to parapharyngeal space (Figure 4).

TORS should be preferred for well circumscribed lesions in the pre- or post-styloid compartment[21] as suggested by Chu *et al*[21]. Transoral robotic surgery was opted in our case as it gives a magnified and high resolution view, along with motion scaling and tremor filtration which allows for delicate dissection around the tumor capsule. This approach is also less morbid compared to the other traditional approaches and reduces the chances of complications like swallowing problems, trismus, auriculotemporal syndrome, first bite syndrome, marginal mandibular and hypoglossal nerve palsy. Cosmesis achieved with this approach is also excellent compared to the transcervical approaches. The other advantages of using robotic surgery in this case include minimal blood loss, shorter hospital stay and early initiation of oral diet for the patient. Postoperatively our patient had no infections or wound dehiscence, and we were able to start orally by day 7.

Recurrence is usually noted in the first year following incomplete excision of lymphangiomas due to regeneration of the lymph vessels[22]. Besides completeness of excision, recurrence rate is based on the type and complexity of the mass. Simple lesions which are completely excised have 0% recurrence rate whereas complex lesions with partial excision have a recurrence rate of 10%–27 % and in those with complete excision, it is 50%–100%[4,7].

CONCLUSION

Lymphangiomas are benign hamartomatous lesions which rarely present in the parapharyngeal space. Large size, dysphagia, respiratory distress, pain due to recurrent infection and hemorrhage are the main indications warranting treatment. MRI is the ideal imaging modality. Transoral robotic surgery is one of the best approaches to parapharyngeal space lesions with favorable anatomy, with minimal morbidity




Figure 4 Surgical algorithm for approach to parapharyngeal tumors including transcervical approach, transoral robotic surgery, transcervical approach with parotidectomy, transcervical approach with styloid excision, transmandibular approach. PPS: Parapharyngeal space; TC: Transcervical approach; TORS: Transoral robotic surgery; TCP: Transcervical approach with parotidectomy; TCS: Transcervical approach with styloid excision; TCM: Transmandibular approach.

and excellent clearance.

ACKNOWLEDGEMENTS

I would like to acknowledge contributions from the department of surgical oncology and robotic surgery, radiology and pathology of BLK- MAX Superspeciality hospital and thank them for the same.

FOOTNOTES

Author contributions: Dabas S, Menon NN, Shukla H, Ranjan R contributed to study concepts; Menon NN, Bhatti SS, Sangal R contributed to study design; Menon NN, Bhatti SS contributed to data acquisition; Menon NN, Shukla H, Ranjan R contributed to quality control of data; Dabas S, Menon NN, Ranjan R, Gurung B, Shukla H, Sharma AK, Tiwari S, Sinha A, Bhatti SS, Sangal R contributed to manuscript preparation, manuscript editing, manuscript reviewing.

Informed consent statement: All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrolment.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: India

ORCID number: Nandini N Menon 0000-0001-6639-7563.

S-Editor: Liu GL

L-Editor: Ma JY

P-Editor: Liu GL

REFERENCES

- 1 **Mehta MR.** Cystic hygroma: Presentation of two casfes with a review of the literature. *Indian J Otolaryngol Head Neck Surg* 2000; **52**: 319-322 [PMID: [23119712](#) DOI: [10.1007/BF03006220](#)]
- 2 **Hartl DM,** Roger G, Denoyelle F, Nicollas R, Triglia JM, Garabedian EN. Extensive lymphangioma presenting with upper airway obstruction. *Arch Otolaryngol Head Neck Surg* 2000; **126**: 1378-1382 [PMID: [11074837](#) DOI: [10.1001/archotol.126.11.1378](#)]
- 3 **Balineni P,** C H VPK, Reddy E D, Reddy JS. Extensive parapharyngeal lymphangioma in a adult-Case report and review of literature. *Int J Surg Case Rep* 2020; **75**: 85-88 [PMID: [32919336](#) DOI: [10.1016/j.ijscr.2020.08.051](#)]
- 4 **Jaiswal AA,** Garg AK, Ravindranath M, Sarkar J, Mohanty MK. 'A huge congenital cervical lymphangioma'-case report with review of literature. *Egyptian Journal of Ear, Nose, Throat and Allied Sciences* 2015; **16**: 283-290 [DOI: [10.1016/j.ejenta.2015.07.005](#)]
- 5 **Aneeshkumar MK,** Kale S, Kabbani M, David VC. Cystic lymphangioma in adults: can trauma be the trigger? *Eur Arch Otorhinolaryngol* 2005; **262**: 335-337 [PMID: [15841412](#) DOI: [10.1007/s00405-004-0780-6](#)]
- 6 **Romeo V,** Maurea S, Guarino S, Sirignano C, Mainenti PP, Picardi M, Salvatore M. A case of lower-neck cystic lymphangioma: correlative US, CT and MR imaging findings. *Quant Imaging Med Surg* 2013; **3**: 224-227 [PMID: [24040619](#) DOI: [10.3978/j.issn.2223-4292.2013.08.03](#)]
- 7 **Grasso DL,** Pelizzo G, Zocconi E, Schleef J. Lymphangiomas of the head and neck in children. *Acta Otorhinolaryngol Ital* 2008; **28**: 17-20 [PMID: [18533550](#)]
- 8 **Sharma JVP,** Kazi FN. Cystic lymphangioma in adult-a rare case scenario or a misdiagnosis? *J Surg Case Rep* 2021; **2021**: rjab062 [PMID: [33747431](#) DOI: [10.1093/jscr/rjab062](#)]
- 9 **Bhayya H,** Pavani D, Avinash Tejasvi ML, Geetha P. Oral lymphangioma: A rare case report. *Contemp Clin Dent* 2015; **6**: 584-587 [PMID: [26681873](#) DOI: [10.4103/0976-237X.169851](#)]
- 10 **de Serres LM,** Sie KC, Richardson MA. Lymphatic malformations of the head and neck. A proposal for staging. *Arch Otolaryngol Head Neck Surg* 1995; **121**: 577-582 [PMID: [7727093](#) DOI: [10.1001/archotol.1995.01890050065012](#)]
- 11 **Giraddi G,** Vanaki SS, Puranik RS. Schwannoma of parapharyngeal space. *J Maxillofac Oral Surg* 2010; **9**: 182-185 [PMID: [22190783](#) DOI: [10.1007/s12663-010-0059-y](#)]
- 12 **Atkinson JC,** Gaze MN, Wilson JA. Benign neck disease. In: Stell and Maran's head and neck surgery. Oxford, Butterworth Heinemann 2000; **4**: 193-194
- 13 **Lee HSH,** Ngai CM. Parapharyngeal space lipoma: a case report. *Hong Kong Med J* 2020; **26**: 438-440 [PMID: [33089789](#) DOI: [10.12809/hkmj198214](#)]
- 14 **Jain SN,** Kumar A, Dhongade HS, Varma R, Hathiram B, Khattar V. Imaging of parapharyngeal space and infratemporal fossa. *Int J Otorhinolaryngol Clin* 2012; **4**: 113-121 [DOI: [10.5005/jp-journals-10003-1096](#)]
- 15 **Riffat F,** Dwivedi RC, Palme C, Fish B, Jani P. A systematic review of 1143 parapharyngeal space tumors reported over 20 years. *Oral Oncol* 2014; **50**: 421-430 [PMID: [24589290](#) DOI: [10.1016/j.oraloncology.2014.02.007](#)]
- 16 **Koeller KK,** Alamo L, Adair CF, Smirniotopoulos JG. Congenital cystic masses of the neck: radiologic-pathologic correlation. *Radiographics* 1999; **19**: 121-46; quiz 152 [PMID: [9925396](#) DOI: [10.1148/radiographics.19.1.g99ja06121](#)]
- 17 **Wang J,** Yang Y, Guo J, Yao Y, Dong L, Mou Y, Zhang Y, Song X. Cervical lymphangioma in adults: A report of seven cases and review of the literature. *Laryngoscope Investig Otolaryngol* 2022; **7**: 751-756 [PMID: [35734075](#) DOI: [10.1002/lto2.801](#)]
- 18 **Colbert SD,** Seager L, Haider F, Evans BT, Anand R, Brennan PA. Lymphatic malformations of the head and neck-current concepts in management. *Br J Oral Maxillofac Surg* 2013; **51**: 98-102 [PMID: [22360972](#) DOI: [10.1016/j.bjoms.2011.12.016](#)]
- 19 **Gowa L,** Rampat R, Khanb A, Mihaimeed F. Adult-onset cystic hygroma: a case report and review of management. *Grand Rounds* 2011; **11**: 5-11 [DOI: [10.1102/1470-5206.2011.0002](#)]
- 20 **Chan JY,** Tsang RK, Eisele DW, Richmon JD. Transoral robotic surgery of the parapharyngeal space: a case series and systematic review. *Head Neck* 2015; **37**: 293-298 [PMID: [24288351](#) DOI: [10.1002/hed.23557](#)]
- 21 **Chu F,** Tagliabue M, Giugliano G, Calabrese L, Preda L, Ansarin M. From transmandibular to transoral robotic approach for parapharyngeal space tumors. *Am J Otolaryngol* 2017; **38**: 375-379 [PMID: [28390805](#) DOI: [10.1016/j.amjoto.2017.03.004](#)]
- 22 **Jakubikova J.** Retropharyngeal lymphangioma. *Bratisl Lek Listy* 2006; **107**: 439-441 [PMID: [17425162](#)]



Wet cupping (Al-hijamah) as a strange cause of ear trauma: A case report

Raid M Al-Ani

Specialty type:

Otorhinolaryngology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model:

Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C, C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Mahmoud MZ, Saudi Arabia; Mirsalehi M, Iran

Received: December 31, 2022

Peer-review started: December 31, 2022

First decision: April 13, 2023

Revised: April 14, 2023

Accepted: April 27, 2023

Article in press: April 27, 2023

Published online: May 9, 2023



Raid M Al-Ani, Department of Surgery/Otolaryngology, University of Anbar College of Medicine, Ramadi City 31001, Anbar, Iraq

Corresponding author: Raid M Al-Ani, MBChB, Academic Editor, Consultant Physician-Scientist, Full Professor, Researcher, Science Editor, Department of Surgery/Otolaryngology, University of Anbar College of Medicine, Al-Andulus, Ramadi City 31001, Anbar, Iraq.
med.raed.alani2003@uoanbar.edu.iq

Abstract

BACKGROUND

Cupping therapy is used across the globe since ancient Egypt. It is used as a complementary or alternative to daily clinical practice. Cupping therapy could be dry or wet (Al-hijamah) type. It is considered a relatively safe procedure for a variety of clinical problems like lower backache, shoulder pain, neck pain, headache, and migraine. Application of cupping therapy over body orifices is contraindicated.

CASE SUMMARY

A 27-year-old gentleman presented with sudden left-sided hearing loss and tinnitus for 7 d. He gave a history of upper respiratory tract infection 14 d before his complaints. He received Al-hijamah over his left ear 3 d before seeking advice from the Otolaryngology clinic, but without benefit. Physical examination showed a dull-looking left ear drum, hemotympanum in the posterior part of the tympanic cavity, two blood clots over the eardrum, and multiple bleeding points over the external ear canal and tympanic membrane. A pure tone audiogram and tympanogram confirmed the diagnosis of middle ear effusion. Conservative treatment (avoidance of water entry, antibiotics, and local and systemic decongestants) was given for 5 d. The presenting symptoms as well as ear injuries resolved completely two weeks after the follow-up.

CONCLUSION

We do not recommend using Al-hijamah directly over the ear as a treatment option for hearing loss to avoid unwanted ear trauma as well as there is no benefit from this kind of treatment.

Key Words: Al-hijamah; Ear trauma; Hemotympanum; Secretary otitis media; Cupping therapy; Hearing loss; Case report

Core Tip: Al-hijamah is a useful treatment for a variety of local and systemic diseases. However, Al-hijamah is contraindicated over the body orifices. A 27-year-old gentleman presented with left-sided hearing loss and tinnitus was treated with a Al-hijamah over the left ear. The patient condition persist as well as there were signs of bleeding (fresh blood, clots, and hemotympanum). Secretary otitis media following upper respiratory tract infection was established on clinical background and audiological tests. He was treated successfully by avoidance of water entry, antibiotics, and decongestant. We don't recommend the usage of Al-hijamah over the ear to treat hearing loss.

Citation: Al-Ani RM. Wet cupping (Al-hijamah) as a strange cause of ear trauma: A case report. *World J Otorhinolaryngol* 2023; 10(2): 30-35

URL: <https://www.wjgnet.com/2218-6247/full/v10/i2/30.htm>

DOI: <https://dx.doi.org/10.5319/wjo.v10.i2.30>

INTRODUCTION

Wet cupping or Al-hijamah is an old traditional therapeutic procedure that is performed in many countries and referred to as the Unani part of medicine. The procedure involves bloodletting from acupoints on a subject's skin to achieve a satisfactory result[1]. Cupping therapy has many indications, both for local (lower back pain, neck and shoulder pain, and headache) and systemic diseases like diabetes mellitus[2]. Cupping therapy should be avoided in patients with organ failure, cancer, cardiovascular diseases, increased risk of cardiovascular disorders (like high serum cholesterol level), hemophilia or other similar blood disorders, and pacemaker. It should not be applied directly over veins, arteries, nerves, skin abnormalities, eyes, body orifices, lymph nodes, and dermatological inflammation[3].

Cupping therapy, like any kind of treatment modality, has adverse effects, especially in the hands of unqualified therapists[4,5]. A recent study reported a beneficial effect of wet cupping therapy as a complementary to steroids in the treatment of a 48-year-old woman with sudden sensorineural hearing loss[6]. However, there is no other study that investigated the use of cupping therapy in the treatment of hearing loss. We reported a 27-year-old male presented with left-sided conductive hearing loss who was treated (by an inexperienced) by the application of an Al-hijamah cup directly over the left ear. Unfortunately, this treatment modality was not useful in resolving the patient condition and resulted in a trauma to the external and middle ears.

CASE PRESENTATION

Chief complaints

Left-sided hearing loss and tinnitus.

History of present illness

A 27-year-old gentleman presented with sudden onset left-sided hearing loss and subjective tinnitus 7 d ago. He gave a history of upper respiratory tract infection 2 wk before his complaints. He was subjected to Al-hijamah directly over the left ear by a therapist to get rid of his presenting features. In addition to the bleeding from the affected ear following Al-hijamah, his clinical symptoms persist. There was no bruising over the affected auricle or postauricular region. The patient denied any history of otorrhea or vertigo.

History of past illness

The patient was healthy with no history of chronic medical diseases.

Personal and family history

The patient had no specific personal or family history.

Physical examination

The patient visited an Otolaryngology clinic. Otoendoscopy revealed several bleeding points covering the left external ear canal, two blood clots, dull looking eardrum, an absent cone of light, and hemotympanum in the posterior part of the left tympanic cavity (Figure 1).




Figure 1 Shows multiple bleeding points (blue arrows) over the left external ear canal, two blood clots over the eardrum (black arrows), hemotympanum in the posterior part of the tympanic cavity (white arrow), absent cone of light, and dull-looking tympanic membrane (brown arrow). A: A little bit far away from the tympanic membrane; B: Closer to the eardrum.

The right ear was completely normal. Tuning fork tests revealed a negative Rinne test on the left side and a positive on the right side, while the Weber test was lateralized to the left ear. Nasal endoscopy revealed clear both nasal cavities and nasopharynx apart from the congestion of the left nasopharyngeal opening of the Eustachian tube.

Audiological tests

Audiological tests in form of pure tone audiogram, which revealed conductive hearing loss in the left ear with an air-bone gap of 37 dB (Figure 2), and a tympanogram that showed a type B curve on the left ear and a type A curve on the right side (Figure 3).

FINAL DIAGNOSIS

The diagnosis was left secretory otitis media superadded by ear trauma from the Al-hijamah procedure.

TREATMENT

We advised the patient to avoid water entry to his left ear, antibiotic cover (amoxicillin 500 mg three times daily for 5 d), nasal decongestant drops (0.05% xylometazoline nasal drops 3 drops for each nostril three time daily for five days), and a combination of systemic decongestant and analgesia tablets (one tablet per day for five days).


OUTCOME AND FOLLOW-UP

At two weeks follow-up, there was a full recovery of all the presenting symptoms, and the left ear canal and ear drum returned to normal.

DISCUSSION

Cupping therapy is used as a complementary or alternative to the usual practice of medicine. It was practiced since ancient Egypt across the globe. It is useful for the management of a wide range of local or systemic clinical problems[7]. The exact therapeutic mechanism of action is not yet described even though an increasing number of investigations include a meta-analysis[8].

Cupping therapy is considered a safe procedure and is part of complementary and alternative medicine. However, frequent cupping treatment over a short period may result in generalized weakness, unsteadiness or drowsiness, and light headiness[9]. Other adverse effects might lead to bruising and skin discoloration, ecchymosis, bleeding, anemia, skin pigmentation, blisters, wound infection, and abscesses, particularly, if cupping therapy is performed by an unqualified or inexperienced person[4,5]. However, ear trauma due to Al-hijamah, as in our case, is not reported.



DOI: 10.5319/wjo.v10.i2.30 Copyright ©The Author(s) 2023.

Figure 2 Pure tone audiogram of the 27-year-old man shows conductive hearing loss of 37 dB in the left ear. Hz: Hertz; dB: Decibel; Circle: Air conduction hearing threshold at a specific frequency of the right ear; X: Air conduction hearing threshold at a specific frequency of the left ear; <: Bone conduction hearing threshold at a specific frequency of the right ear with masking of the left ear;]: Bone conduction hearing threshold at a specific frequency of the left ear.

Ear trauma usually occurs in young adult men owing to the high outdoor activities. It is usually caused during personal assaults, sports, or following personal accidents and road traffic accidents. Ear injury results in various clinical problems depending on the velocity and mechanism of the injury. These include; hematoma auris, lacerations or wounds, partial or complete avulsion of the auricle, hemotympanum, eardrum perforation, dislocation of the ossicles, Otic barotrauma, facial palsy, and inner ear injury[10]. Although, Al-hijamah is widely used in Iraq due to cultural and religious beliefs and resulted in the improvement of many conditions like diabetes mellitus, hypertension, and hyperlipidemia, treatment for hearing loss is not used before[11]. A recent case report study from Qatar reported that Al-hijamah is used for the treatment of sudden sensorineural hearing loss as a complementary treatment with steroids in a 48-year-old woman with satisfactory results[6]. To our best knowledge, there is no other study in the world that used cupping therapy as a treatment modality for hearing loss. Unfortunately, in the presenting case, cupping therapy was applied directly on the auricle resulting in hemotympanum and bleeding from the ear without improvement of the patient complaints (hearing loss and tinnitus). We do not recommend using Al-hijamah directly over the ear to avoid unwanted ear injuries as well as there is no benefit from its use. Furthermore, this study supports that cupping therapy is contraindicated over body orifices[3].

CONCLUSION

The direct application of cupping therapy on the external meatus results in bleeding from the ear canal and haemotympanum as well as there was no improvement of the patients' hearing loss. Therefore, we do not recommend using Al-hijamah directly on the ear.




Figure 3 Tympanogram of the 27-year-old man shows type A on the right ear and type B (flat curve) on the left ear. The left external canal volume is 0.99 mL.

FOOTNOTES

Author contributions: Al-Ani RM has the responsibility for the design of the study, writing the abstract, core tip, introduction, case presentation, discussion, conclusion, and references according to the journal style, and edited the draft and preparing the final version of the manuscript; Al-Ani RM revised and approved the final version of the manuscript.

Informed consent statement: Informed consent was obtained from the patient for publication of the case with its related images.

Conflict-of-interest statement: The author declares that there is no conflict of interest.

CARE Checklist (2016) statement: The author have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Iraq

ORCID number: Raid M Al-Ani 0000-0003-4263-9630.

S-Editor: Fan JR

L-Editor: A

P-Editor: Zhao S

REFERENCES

- 1 **Almaiman AA.** Proteomic effects of wet cupping (Al-hijamah). *Saudi Med J* 2018; **39**: 10-16 [PMID: [29332103](#) DOI: [10.15537/smj.2018.1.21212](#)]
- 2 **Aboushanab TS, AlSanad S.** Cupping Therapy: An Overview from a Modern Medicine Perspective. *J Acupunct Meridian Stud* 2018; **11**: 83-87 [PMID: [29436369](#) DOI: [10.1016/j.jams.2018.02.001](#)]
- 3 **Furhad S, Bokhari AA.** Cupping Therapy. 2023 Jan 2. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan- [PMID: [30855841](#)]
- 4 **Refaat B, El-Shemi AG, Ebid AA, Ashshi A, BaSalamah MA.** Islamic wet cupping and risk factors of cardiovascular diseases: effects on blood pressure, metabolic profile and serum electrolytes in healthy young adult men. *Altern Integ Med* 2014; **3**: 151 [DOI: [10.4172/2327-5162.1000151](#)]
- 5 **Vaccaro M, Coppola M, Ceccarelli M, Montopoli M, Guarneri C.** The good and the bad of cupping therapy: case report and review of the literature. *Eur Rev Med Pharmacol Sci* 2021; **25**: 2327-2330 [PMID: [33755970](#) DOI: [10.26355/eurev.202103.25266](#)]
- 6 **Almusleh ZA, El Ansari W.** Integrating Cupping Therapy in the Management of Sudden Sensorineural Hearing Loss: A Case Report. *Cureus* 2020; **12**: e7063 [PMID: [32226665](#) DOI: [10.7759/cureus.7063](#)]
- 7 **Choi TY, Ang L, Ku B, Jun JH, Lee MS.** Evidence Map of Cupping Therapy. *J Clin Med* 2021; **10** [PMID: [33920643](#) DOI: [10.3390/jcm10081750](#)]
- 8 **Al-Bedah AMN, Elsubai IS, Qureshi NA, Aboushanab TS, Ali GIM, El-Olemy AT, Khalil AAH, Khalil MKM, Alqaed MS.** The medical perspective of cupping therapy: Effects and mechanisms of action. *J Tradit Complement Med* 2019; **9**: 90-97 [PMID: [30963043](#) DOI: [10.1016/j.jtcme.2018.03.003](#)]
- 9 **Kim KH, Kim TH, Hwangbo M, Yang GY.** Anaemia and skin pigmentation after excessive cupping therapy by an unqualified therapist in Korea: a case report. *Acupunct Med* 2012; **30**: 227-228 [PMID: [22738805](#) DOI: [10.1136/acupmed-2012-010185](#)]
- 10 **Eagles K, Fralich L, Stevenson JH.** Ear trauma. *Clin Sports Med* 2013; **32**: 303-316 [PMID: [23522511](#) DOI: [10.1016/j.csm.2012.12.011](#)]
- 11 **Rahman HS, Ahmad GA, Mustapha B, Al-Rawi HA, Hussein RH, Amin K, Othman HH, Abdullah R.** Wet cupping therapy ameliorates pain in patients with hyperlipidemia, hypertension, and diabetes: A controlled clinical study. *Int J Surg Open* 2020; **26**: 10-15 [DOI: [10.1016/j.ijso.2020.07.003](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

