

# World Journal of *Gastrointestinal Pathophysiology*

*World J Gastrointest Pathophysiol* 2017 February 15; 8(1): 1-10





## Editorial Board

2016-2019

The *World Journal of Gastrointestinal Pathophysiology* Editorial Board consists of 503 members, representing a team of worldwide experts in gastrointestinal pathophysiology. They are from 45 countries, including Argentina (2), Australia (13), Austria (2), Belgium (7), Brazil (9), Brunei Darussalam (1), Canada (21), China (30), Croatia (1), Czech Republic (2), Denmark (3), Egypt (1), Estonia (1), Finland (1), France (8), Germany (24), Greece (7), Hungary (5), India (12), Indonesia (1), Iran (2), Ireland (2), Israel (8), Italy (38), Japan (45), Lebanon (3), Malaysia (1), Mexico (2), Netherlands (7), Norway (1), Poland (4), Portugal (1), Romania (1), Russia (1), Singapore (4), South Korea (12), Spain (24), Sweden (11), Switzerland (3), Thailand (2), Turkey (7), Ukraine (1), United Kingdom (10), United States (161) and Venezuela (1).

### EDITOR-IN-CHIEF

Thomas Y Ma, *Albuquerque*

### ASSOCIATE EDITOR

Naohisa Yoshida, *Kyoto*

### GUEST EDITORIAL BOARD MEMBERS

Jia-Ming Chang, *Taipei*  
Wai-Keung Chow, *Taichung*  
Chien-Wei Hsu, *Kaohsiung*  
Ming-Tsan Lin, *Taipei*  
Bor-Shyang Sheu, *Tainan*  
Jin-Town Wang, *Taipei*

### MEMBERS OF THE EDITORIAL BOARD



#### Argentina

Bernabé M Quesada, *Buenos Aires*  
Marcelo G Roma, *Rosario*



#### Australia

Chris R Abbiss, *Joondalup*  
Guy D Eslick, *Sydney*  
Montri Gururatsakul, *South Australia*  
Chandana B Herath, *Victoria*  
Michael Horowitz, *Adelaide*  
Mustafa Khasraw, *Victoria*  
Shu-Chuen Li, *New South Wales*  
Michael HJ Maes, *Geelong*  
Nam Q Nguyen, *South Australia*  
Kulmira Nurgali, *Victoria*  
Nicholas J Spencer, *South Australia*

Deborah J Verran, *NSW*  
Shufeng Zhou, *Victoria*



#### Austria

Cord Langner, *Graz*  
Dietmar Ofner-Velano, *Salzburg*



#### Belgium

Kathleen Blondeau, *Leuven*  
Ilse M Hoffman, *Leuven*  
Theodoor A Niewold, *Heverlee*  
Xavier Sagaert, *Leuven*  
Jean-Marie Vanderwinden, *Brussels*  
Kristin Verbeke, *Belgium*  
Mathieu Vinken, *Brussels*



#### Brazil

Uilian Andreis, *Botucatu*  
Everson LDA Artifon, *Sao Paulo*  
Joao B Calixto, *Trindade*  
Niels OS Camara, *Cidade Universitária*  
Júlio MF Chebli, *Juiz de Fora*  
Clélia A Hiruma-Lima, *Botucatu*  
Marcel CC Machado, *Sao Paulo*  
Juarez AS Quaresma, *Belem*  
Wagner Vilegas, *Araraquara*



#### Brunei Darussalam

Vui H Chong, *Bandar Seri Begawan*



#### Canada

Jean-Francois Beaulieu, *Sherbrooke*  
Francois Boudreau, *Sherbrooke*  
George A Bubenik, *Guelph*  
Wangxue Chen, *Ottawa*  
Fernando Fornari, *Montréal*  
Jan D Huizinga, *Puslinch*  
Ruben Hummelen, *Dundas*  
Kusum K Kharbanda, *Omaha*  
Wolfgang A Kunze, *Hamilton*  
Jianjun Li, *Ottawa*  
Roderick J MacLeod, *Kingston*  
Michele Molinari, *Halifax*  
Nathalie Rivard, *Sherbrooke*  
Nathalie Rivard, *Sherbrooke*  
Kirill Rosen, *Halifax*  
Manuela M Santos, *Montreal*  
Caroline Saucier, *Québec*  
Jean Sévigny, *Québec*  
Eldon A Shaffer, *Calgary*  
Alan BR Thomson, *Edmonton*  
Pierre H Vachon, *Sherbrooke*



#### China

Kai-Xing Ai, *Shanghai*  
Zhao-Xiang Bian, *Hong Kong*  
Minhu Chen, *Guangzhou*  
Chi-Hin Cho, *Hong Kong*  
Zhong-Hong Gao, *Wuhan*  
Jun-Ming Guo, *Ningbo*  
Jing-Yan Han, *Beijing*  
Jian-Dong Huang, *Hong Kong*  
Jia-Fu Ji, *Beijing*

Zhan-Ju Liu, *Shanghai*  
 Shi Liu, *Wuhan*  
 Zhen-Ning Wang, *Shenyang*  
 Xiao-Hong Wang, *Beijing*  
 Wei Wei, *Hefei*  
 Dong-Ping Xie, *Shanghai*  
 Wen-Xie Xu, *Shanghai*  
 Xiao Yang, *Beijing*  
 Hua Yang, *Chongqing*  
 Wei-Zhen Zhang, *Beijing*  
 Hua-Chuan Zheng, *Shenyang*  
 Min-Sheng Zhu, *Nanjing*  
 Jinxia Zhu, *Beijing*  
 Da-Ling Zhu, *Harbin*  
 Yong-Liang Zhu, *Hangzhou*



**Croatia**

Alen Protic, *Rijeka*



**Czech Republic**

Pavel Hladik, *Semily*  
 Martin Vokurka, *Prague*



**Denmark**

Lars Arendt-Nielsen, *Aalborg*  
 Frank V Schiodt, *Copenhagen*  
 Jonas Worsoe, *Aarhus*



**Egypt**

Mahmoud Aboelneen Khattab, *Minia*



**Estonia**

Enn Seppet, *Tartu*



**Finland**

Pauli Antero Puolakkainen, *Turku*



**France**

Bruno Bonaz, *Grenoble*  
 Pierre M Dechelotte, *Rouen*  
 Jean-Paul Lallès, *Saint-Gilles*  
 Charles-Henri Malbert, *Saint-Gilles*  
 Thierry Piche, *Nice*  
 Pascale Plaisancié, *Lyon*  
 Michelina Plateroti, *Lyon*  
 Véronique Vitton, *Marseille*



**Germany**

Hans G Beger, *Ulm*  
 Carsten Bergmann, *Ingelheim*  
 Elke Cario, *Essen*  
 Arno J Dormann, *Koln*  
 Nikolaus Gassler, *Aachen*  
 Andreas Geier, *Wuerzburg*

Werner Hartwig, *Heidelberg*  
 Marion Hewicker-Trautwein, *Hannover*  
 Jens Hoepfner, *Freiburg*  
 Tobias Keck, *Freiburg*  
 Jorg Kleeff, *Munich*  
 Peter Malfertheimer, *Magdeburg*  
 Oliver Mann, *Hamburg*  
 Christoph W Michalski, *Munich*  
 Andreas K Nussler, *Munich*  
 Christian Pehl, *Vilsbiburg*  
 Peter Schemmer, *Heidelberg*  
 Manuel A Silva, *Penzberg*  
 Marc Stemmler, *Freiburg*  
 Frank Tacke, *Aachen*  
 Sya N Ukena, *Braunschweig*  
 Brigitte Vollmar, *Rostock*  
 Thomas M Wex, *Magdeburg*  
 Margot Zoller, *Heidelberg*



**Greece**

Stelios F Assimakopoulos, *Patras*  
 George N Dalekos, *Larissa*  
 Alkiviadis Efthymiou, *thessaloniki*  
 Maria Gazouli, *Athens*  
 Ioannis E Koutroubakis, *Heraklion*  
 Gerassimos J Mantzaris, *Athens*  
 George V Papatheodoridis, *Athens*



**Hungary**

Mária Bagyánszki, *Szeged*  
 Mihály Boros, *Szeged*  
 Laszlo Czako, *Szeged*  
 Pal Miheller, *Budapest*  
 Zoltan Rakonczay, *Szeged*



**India**

Anil K Agarwal, *Delhi*  
 Uday Bandyopadhyay, *Kolkata*  
 Sriparna Basu, *Varanasi*  
 Chandra K Chakraborti, *Rourkela*  
 Nilesh M Dagia, *Mumbai*  
 Rajeev Garg, *Punjab*  
 Akhtar Mahmood, *Chandigarh*  
 Veerareddy P Reddy, *Andhra Pradesh*  
 Chandra P Sharma, *Thiruvananthapuram*  
 Shailesh V Shrikhande, *Mumbai*  
 Virendra Singh, *Chandigarh*  
 Nicholas J Skill, *Indianapolis*



**Indonesia**

Laurentius A Lesmana, *Jakarta*



**Iran**

Gholamreza Roshandel, *Gorgan*  
 Shahram Shahabi, *Urmia*



**Ireland**

Billy Bourke, *Dublin*  
 Stephen J Keely, *Dublin*



**Israel**

Yosefa Avraham, *Jerusalem*  
 Yaron Bar-Dayana, *Holon*  
 Shomron Ben-Horin, *Tel-Hashomer*  
 Boris Kirshtein, *Beer Sheva*  
 Stephen Malnick, *Rehovot*  
 Yaakov Maor, *Tel-Hashomer*  
 Rifaat Safadi, *Jerusalem*  
 Nachum Vaisman, *Tel Aviv*



**Italy**

Rosaria Acquaviva, *Catania*  
 Alessandro Antonelli, *Pisa*  
 Giacosa Attilio, *Genova*  
 Salvatore Auricchio, *Naples*  
 Guido Basilisco, *Milan*  
 Antonio Basoli, *Rome*  
 Massimo Bellini, *Pisa*  
 Luigi Bonavina, *Milano*  
 Alfio Brogna, *Catania*  
 Giuseppe Calamita, *Bari*  
 Raffaele Capasso, *Naples*  
 Ignazio Castagliuolo, *Padova*  
 Francesco Cresi, *Torino*  
 Rosario Cuomo, *Napoli*  
 Salvatore Cuzzocrea, *Gazzi*  
 Mario M D'Elios, *Florence*  
 Cinzia Domeneghini, *Milan*  
 Luca Elli, *Milano*  
 Walter Fries, *Messina*  
 Eugenio Gaudio, *Rome*  
 Marco Gobetti, *Bari*  
 Fabio Grizzi, *Rozzano*  
 Enzo Grossi, *Milanese*  
 Enzo Ierardi, *Foggia*  
 Pietro Invernizzi, *Monza*  
 Angelo A Izzo, *Naples*  
 Anna Kohn, *Rome*  
 Giovanni Latella, *L'Aquila*  
 Massimo Marignani, *Rome*  
 Sergio Morini, *Rome*  
 Raffaele Pezzilli, *Bologna*  
 Cristiano Rumio, *Milan*  
 Giovanni Sarnelli, *Naples*  
 Edoardo V Savarino, *Padua*  
 Pierpaolo Sileri, *Roma*  
 Annamaria Staiano, *Naples*  
 Giacomo C Sturniolo, *Padua*  
 Claudio Tiribelli, *Trieste*



**Japan**

Hirohata Akiho, *Fukuoka*  
 Akihiro Asakawa, *Kagoshima*  
 Hisashi Aso, *Sendai*  
 Yasu-Taka Azuma, *Osaka*  
 Shotaro Enomoto, *Wakayama*  
 Mikihiro Fujiya, *Hokkaido*  
 Takahisa Furuta, *Hamamatsu*  
 Akira Hokama, *Nishihara*  
 Ryota Hokari, *Saitama*  
 Yuichi Hori, *Hyogo*  
 Masahiro Iizuka, *Akita*  
 Shoji Ikuo, *Kobe*  
 Motohiro Imano, *Osaka*



Hajime Isomoto, *Nagasaki*  
 Tatehiro Kagawa, *Isehara*  
 Haruki Kitazawa, *Sendai*  
 Shigeru BH Ko, *Tokyo*  
 Xiao-Kang Li, *Tokyo*  
 Noriaki Manabe, *Okayama*  
 Atsushi Masamune, *Sendai*  
 Hiroyuki Matsubayashi, *Suntogun*  
 Kazuyuki Matsushita, *Inohana Chuo-ku*  
 Reiko Miyazawa, *Gunma*  
 Kazunari Murakami, *Oita*  
 Hikaru Nagahara, *Tokyo*  
 Yuji Naito, *Kyoto*  
 Atsushi Nakajima, *Yokohama*  
 Shoji Natsugoe, *Kagoshima City*  
 Tsutomu Nishida, *Suita*  
 Koji Nomoto, *Tokyo*  
 Shouji Shimoyama, *Tokyo*  
 Goshi Shiota, *Yonago*  
 Hidekazu Suzuki, *Tokyo*  
 Hitoshi Takagi, *Gunma*  
 Toru Takahashi, *Okayama*  
 Yoshihisa Takahashi, *Tokyo*  
 Kan Uchiyama, *Chiba*  
 Yoshiyuki Ueno, *Sendai*  
 Takato Ueno, *Fukuoka*  
 Hisayuki Uneyama, *Kawasaki*  
 Mitsunori Yamakawa, *Yamagata*  
 Takayuki Yamamoto, *Mie*  
 Yutaka Yata, *Maebashi-city*  
 Naohisa Yoshida, *Kyoto*  
 Hitoshi Yoshiji, *Kashihara*



#### **Lebanon**

Costantine F Daher, *Byblos*  
 Assaad M Soweid, *Beirut*  
 Julnar Usta, *Beirut*



#### **Malaysia**

Andrew CS Boon, *Ipoh*



#### **Mexico**

José M de la Roca-Chiapas, *Leon*  
 Maria-Raquel Huerta-Franco, *Leon*



#### **Netherlands**

Wouter J de Jonge, *Amsterdam*  
 Aldo Grefhorst, *Groningen*  
 Daniel Keszthelyi, *Maastricht*  
 Cornelis FM Sier, *ZA Leiden*  
 Pieter J Tanis, *AZ Amsterdam*  
 Luc JW van der Laan, *Rotterdam*  
 Sander van der Marel, *Leiden*



#### **Norway**

Anne M Bakke, *Oslo*



#### **Poland**

Stanisław A Hac, *Gdansk*

Stanisław J Kowalski, *Krakow*  
 Agata Mulak, *Wroclaw*  
 Napoleon Waszkiewicz, *Choroszcz*



#### **Portugal**

Ricardo Marcos, *Porto*



#### **Romania**

Mihai Ciocirlan, *Bucharest*



#### **Russia**

Ludmila Filaretova, *Petersburg*



#### **Singapore**

Madhav Bhatia, *Singapore*  
 Brian K Goh, *Singapore*  
 Khek-Yu Ho, *Singapore*  
 Cliff KS Ong, *Singapore*



#### **South Korea**

Jae Hee Cheon, *Seoul*  
 Myung-Haing Cho, *Seoul*  
 Ki-Baik Hahm, *Seongnam*  
 HO Jae Han, *Gwangju*  
 Chang-Duk Jun, *Gwangju*  
 Sang Geon Kim, *Seoul*  
 Hong Joo Kim, *Seoul*  
 Jin-Kyung Kim, *Gyeongseong-si*  
 Won-Jae Lee, *Seoul*  
 Sung-Joo Park, *Iksan*  
 Seung Ha Park, *Gangwon Do*  
 Kwan-Kyu Park, *Daegu*



#### **Spain**

Raquel Abalo, *Madrid*  
 Juan G Abalades, *Barcelona*  
 Dario Acuna-Castroviejo, *Armilla*  
 Agustin Albillos, *Madrid*  
 Maria-Angeles Aller, *Madrid*  
 Fernando Azpiroz, *Barcelona*  
 Ramón Bataller, *Barcelona*  
 Marco Bustamante-Balén, *Valencia*  
 Andres Cardenas, *Barcelona*  
 Joan Claria, *Barcelona*  
 Pere Clavé, *Barcelona*  
 Manuel B de Acosta, *Santiago*  
 Manuel Giner, *Madrid*  
 Borja Hernandez-Breijo, *Madrid*  
 Angel I Lanas, *Zaragoza*  
 Maite T Martin, *Barcelona*  
 Vicente Martinez, *Bellaterra*  
 Jose M Mates, *Malaga*  
 Julio Mayol, *Madrid*  
 Marçal Pastor-Anglada, *Barcelona*  
 María E Sáez, *Seville*  
 Yolanda Sanz, *Burjassot*  
 Carlos Taxonera, *Madrid*

Maria D Yago, *Granada*



#### **Sweden**

Marco Del Chiaro, *Stockholm*  
 Frida Fak, *Gothenburg*  
 Gunnar FA Flemstrom, *Uppsala*  
 Evangelos Kalaitzakis, *Gothenburg*  
 Kristina Lamas, *Umea*  
 Bob R Olsson, *Goteborg*  
 Sara M Regné, *Malmö*  
 Peter T Schmidt, *Stockholm*  
 Xiao-Feng Sun, *Linköping*  
 Henrik Thorlacius, *Malmö*  
 Curt Tysk, *Orebro*



#### **Switzerland**

Jyrki J Eloranta, *Zurich*  
 Remy Meier, *Liestal*  
 Catherine M Pastor, *Geneva*



#### **Thailand**

Thawatchai Akaraviputh, *Bangkok*  
 Weekitt Kittisupamongkol, *Bangkok*



#### **Turkey**

Mehmet Bektas, *Ankara*  
 Ugur Duman, *Bursa*  
 Mukaddes Esrefoglu, *Istanbul*  
 Ahmet Guven, *Ankara*  
 Muammer Karadeniz, *Bornova-Izmir*  
 Elvan Ozbek, *Sakarya*  
 Ilhami Yuksel, *Ankara*



#### **Ukraine**

Oksana S Zayavhivska, *Lviv*



#### **United Kingdom**

Geoffrey Burnstock, *London*  
 Janice E Drew, *Aberdeen*  
 Girish L Gupte, *Birmingham*  
 David C Hay, *Edinburgh*  
 Nusrat Husain, *Manchester*  
 Michael L Lucas, *Glasgow*  
 Jamie Murphy, *London*  
 Vadim Sumbayev, *Chatham Maritime*  
 Wing-Kin Syn, *Birmingham*  
 Andrea Varro, *Liverpool*



#### **United States**

Sami R Achem, *Florida*  
 Tauseef Ali, *Oklahoma City*  
 Shrikant Anant, *Oklahoma*  
 Mohammed S Anwer, *North Grafton*  
 Andrew Aronsohn, *Chicago*  
 Toms Augustin, *Sayre*

Gyorgy Baffy, *Boston*  
 Michael T Bailey, *Columbus*  
 Kim E Barrett, *San Diego*  
 Marc D Basson, *Grand Forks*  
 Robert L Bell, *Berkeley Heights*  
 David H Berger, *Texas*  
 Urs A Boelsterli, *Storrs*  
 Michael W Bradbury, *Erie*  
 Qiang Cai, *Atlanta*  
 Weibiao Cao, *Providence*  
 Subhash C Chauhan, *Sioux Falls*  
 Tao-Sheng Chen, *Memphis*  
 Jian-De Chen, *Galveston*  
 Chiang YL Chiang, *Rootstown*  
 Mashkoor A Choudhry, *Maywood*  
 Parimal Chowdhury, *Little Rock*  
 Eric Cohen, *Boston*  
 Robert T Cormier, *Duluth*  
 Edwin A Deitch, *Newark*  
 Sharon DeMorrow, *Texas*  
 Dan A Dixon, *Columbia*  
 James P Dolan, *Portland*  
 Henry H Dong, *Pittsburgh*  
 Hui Dong, *La Jolla*  
 Ashkan Farhadi, *Irvine*  
 Bin Feng, *Pittsburgh*  
 Jenifer Fenton, *East Lansing*  
 Alessandro Fichera, *Chicago*  
 Mitchell P Fink, *Pittsburgh*  
 Leo R Fitzpatrick, *Rancho Cordova*  
 Robert A Forse, *Omaha*  
 Glenn T Furuta, *Aurora*  
 Juan F Gallegos-Orozco, *Scottsdale*  
 Pandu R Gangula, *Nashville*  
 Timothy B Gardner, *Lebanon*  
 Shannon S Glaser, *Temple*  
 Beverley Greenwood-VanMeerveld, *Oklahoma City*  
 John R Grider, *Richmond*  
 Yan-Fang Guan, *Cincinnati*  
 Gregory M Holmes, *Baton Rouge*  
 Richard Hu, *Los Angeles*  
 Hartmut Jaeschke, *Kansas City*  
 Robert T Jensen, *Bethesda*  
 Sreenivasa S Jonnalagadda, *Louis*  
 Michel Kahaleh, *Charlottesville*  
 Andreas M Kaiser, *Los Angeles*  
 Randeep S Kashyap, *Rochester*  
 Richard Kellermayer, *Houston*  
 Chris Kevil, *Shreveport*  
 Sandeep Khurana, *Baltimore*  
 Pawel R Kiela, *Tucson*  
 Tammy L Kindel, *Cincinnati*  
 Gordana Kosutic, *Durham*  
 David Kravetz, *San Diego*

Ashok Kumar, *Detroit*  
 Anthony Kumar, *Los Angeles*  
 John H Kwon, *Chicago*  
 Muriel Larauche, *Los Angeles*  
 Ai-Xuan Le Holterman, *Chicago*  
 I. Michael Leitman, *New York*  
 Felix W Leung, *Sepulveda*  
 Suthat Liangpunsakul, *Indianapolis*  
 Feng-Xin Lu, *Boston*  
 Pauline K Lund, *Chapel Hill*  
 Guang-Xiang Luo, *Lexington*  
 Jay Luther, *Ann Arbor*  
 Ram I Mahato, *Memphis*  
 Akhil Maheshwari, *Birmingham*  
 Kenneth Maiese, *Newark*  
 Adhip PN Majumdar, *Detroit*  
 José E Manautou, *Storrs*  
 Craig J McClain, *Louisville*  
 Dermot PB McGovern, *Los Angeles*  
 Douglas S Merrell, *Bethesda*  
 Murielle Mimeault, *Omaha*  
 Emiko Mizoguchi, *Boston*  
 Huan-Biao Mo, *Denton*  
 Adam J Moeser, *Raleigh*  
 Ramzi M Mohammad, *Detroit*  
 Satdarshan SP Monga, *Pittsburgh*  
 Roger K Moreira, *New York*  
 Sandeep Mukherjee, *Omaha*  
 Karnam S Murthy, *Richmond*  
 Michael J Nowicki, *Jackson*  
 Shuji Ogino, *Boston*  
 Mary F Otterson, *Wisconsin*  
 Chung Owyang, *Ann Arbor*  
 Helieh S Oz, *Lexington*  
 Marco G Patti, *Chicago*  
 Timothy M Pawlik, *Baltimore*  
 Sara Peleg, *Houston*  
 Li-Ya Qiao, *Richmond*  
 Chao Qin, *Oklahoma*  
 Parvaneh Rafiee, *Milwaukee*  
 Sigrid A Rajasekaran, *Wilmington*  
 Vazhaikkurichi Rajendran, *Morgantown*  
 Jean P Raufman, *Baltimore*  
 Ramesh M Ray, *Memphis*  
 Arie Regev, *Indianapolis*  
 Yehuda Ringel, *Chapel Hill*  
 Richard A Rippe, *Rockville*  
 Chantal A Rivera, *Bossier*  
 Andrea Romani, *Cleveland*  
 Praveen K Roy, *Marshfield*  
 Paul A Rufo, *Boston*  
 David B Sachar, *New York*  
 Bimaljit S Sandhu, *Richmond*  
 Sanjaya K Satapathy, *New Hyde Park*

Ron Schey, *Iowa City*  
 Muhammad Y Sheikh, *Fresno*  
 Bo Shen, *Cleveland*  
 Le Shen, *Chicago*  
 Frank A Simmen, *Little Rock*  
 Steven M Singer, *Washington*  
 Shailinder J Singh, *Washington*  
 Adam J Smolka, *Charleston*  
 Ned Snyder, *Houston*  
 Zhen-Yuan Song, *Chicago*  
 Gagan K Sood, *Houston*  
 Rhonda Souza, *Dallas*  
 Stuart J Spechler, *Dallas*  
 Subbaramiah Sridhar, *Augusta*  
 Catia Sternini, *Los Angeles*  
 Veedamali S Subramanian, *Long Beach*  
 Jun Sun, *Rochester*  
 Yvette Taché, *Los Angeles*  
 Xiao-Di Tan, *Chicago*  
 Paul D Terry, *Atlanta*  
 Ma Thomas, *Albuquerque*  
 Jennifer S Tirnauer, *Farmington*  
 Andrea Todisco, *Ann Arbor*  
 George C Tsokos, *Boston*  
 Vic Velanovich, *Detroit*  
 Raj Vuppalachchi, *Indianapolis*  
 Estela Wajcberg, *Cranford*  
 Arnold Wald, *Madison*  
 Li-Xin Wang, *Los Angeles*  
 Horst C Weber, *Boston*  
 Guo-Yao Wu, *Texas*  
 Christian Wunder, *Bethesda*  
 Zuo-Liang Xiao, *Cleveland*  
 Guang-Yin Xu, *Galveston*  
 Guo-Rong Xu, *East Orange*  
 Yoshio Yamaoka, *Houston*  
 Guang-Yu Yang, *Chicago*  
 Jay A Yelon, *Valhalla*  
 Shao-Yong Yu, *Hershey*  
 Yana Zavros, *Cincinnati*  
 Joerg Zehetner, *Los Angeles*  
 Jian X Zhang, *Charlotte*  
 Zhi Zhong, *Charleston*  
 Hui-Ping Zhou, *Richmond*  
 Zhan-Xiang Zhou, *Kannapolis*  
 Qing Zhu, *Bethesda*  
 Yao-Hui Zhu, *Stanford*



**Venezuela**

Fabian Michelangeli, *Caracas*



**FIELD OF VISION**

- 1 Impaired inactivation of digestive proteases: The possible key factor for the high susceptibility of germ-free and antibiotic-treated animals to gut epithelial injury  
*Qin X*

**ORIGINAL ARTICLE**

**Observational Study**

- 3 Polyp detection rate and pathological features in patients undergoing a comprehensive colonoscopy screening  
*Asadzadeh Aghdai H, Nazemalhosseini Mojarad E, Ashtari S, Pourhoseingholi MA, Chaleshi V, Anaraki F, Haghazali M, Zali MR*

## Contents

*World Journal of Gastrointestinal Pathophysiology*  
Volume 8 Number 1 February 15, 2017

### ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Pathophysiology*, Guang-Yu Yang, MD, PhD, Professor, Department of Pathology, Northwestern University, Chicago, IL 60611, United States

### AIM AND SCOPE

*World Journal of Gastrointestinal Pathophysiology* (*World J Gastrointest Pathophysiol*, *WJGP*, online ISSN 2150-5330, DOI: 10.4291), is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

*WJGP* is to report rapidly the most recent results in basic and clinical research on gastrointestinal pathophysiology, including all aspects of normal or abnormal function of the gastrointestinal tract, hepatobiliary system, and pancreas. *WJGP* specifically covers growth and development, digestion, secretion, absorption, metabolism and motility relative to the gastrointestinal organs, as well as immune and inflammatory processes, and neural, endocrine and circulatory control mechanisms that affect these organs. This journal will also report new methods and techniques in gastrointestinal pathophysiological research.

We encourage authors to submit their manuscripts to *WJGP*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

### INDEXING/ABSTRACTING

*World Journal of Gastrointestinal Pathophysiology* is now indexed in PubMed, PubMed Central.

### FLYLEAF

I-IV Editorial Board

### EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*  
Responsible Electronic Editor: *Dan Li*  
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*  
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL  
*World Journal of Gastrointestinal Pathophysiology*

ISSN  
ISSN 2150-5330 (online)

LAUNCH DATE  
April 15, 2010

Frequency  
Quarterly

EDITOR-IN-CHIEF  
**Thomas Y Ma, MD, PhD, Professor, Chief**, Division of Gastroenterology and Hepatology, University of New Mexico, MSC10 5550, 1 UNM, Albuquerque, NM 87131, United States

EDITORIAL BOARD MEMBERS  
All editorial board members resources online at <http://www.wjgnet.com/2150-5330/editorialboard.htm>

EDITORIAL OFFICE  
Xiu-Xia Song, Director  
*World Journal of Gastrointestinal Pathophysiology*  
Baishideng Publishing Group Inc  
8226 Regency Drive, Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>  
<http://www.wjgnet.com>

PUBLISHER  
Baishideng Publishing Group Inc  
8226 Regency Drive, Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>  
<http://www.wjgnet.com>

PUBLICATION DATE  
February 15, 2017

COPYRIGHT  
© 2017 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT  
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS  
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION  
<http://www.wjgnet.com/esps/>

# Impaired inactivation of digestive proteases: The possible key factor for the high susceptibility of germ-free and antibiotic-treated animals to gut epithelial injury

Xiaofa Qin

Xiaofa Qin, GI Biopharma Inc, Westfield, NJ 07090, United States

Author contributions: Qin X contributes solely to this paper.

Conflict-of-interest statement: There is no conflict of interest.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Xiaofa Qin, MD, PhD, GI Biopharma Inc, 918 Willow Grove Road, Westfield, NJ 07090, United States. [xiaofa\\_qin@yahoo.com](mailto:xiaofa_qin@yahoo.com)  
Telephone: +1-908-4637423

Received: October 9, 2016

Peer-review started: October 11, 2016

First decision: December 13, 2016

Revised: December 16, 2016

Accepted: January 2, 2017

Article in press: January 3, 2017

Published online: February 15, 2017

## Abstract

Recent study shows that germ-free and antibiotic-treated animals are highly susceptible to gut epithelial injury. This paper addresses that impaired inactivation of digestive proteases may be the key factor for the increased susceptibility.

**Key words:** Digestive proteases; Germ-free; Antibiotics; Gut microbiota; Gut epithelial injury

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** This paper addresses that impaired inactivation of digestive proteases may be the possible key factor for the high susceptibility of germ-free and antibiotic-treated animals to gut epithelial injury.

Qin X. Impaired inactivation of digestive proteases: The possible key factor for the high susceptibility of germ-free and antibiotic-treated animals to gut epithelial injury. *World J Gastrointest Pathophysiol* 2017; 8(1): 1-2 Available from: URL: <http://www.wjgnet.com/2150-5330/full/v8/i1/1.htm> DOI: <http://dx.doi.org/10.4291/wjgp.v8.i1.1>

## COMMENTARY ON HOT TOPICS

I read with great interest the paper by Hernández-Chirilaque *et al*<sup>[1]</sup> regarding the high susceptibility of germ-free and antibiotic-treated mice to epithelial injury. It is found that germ-free mice treated with dextran sulfate sodium (DSS) showed only minimal inflammation (no colonic thickening, lower myeloperoxidase activity, IL-6, IL-17, TNF-alpha and IFN-gamma secretion by splenocytes and mesenteric cell cultures, *etc.*), but enhanced hemorrhage, epithelial injury and mortality along with weakened intestinal barrier. Animals treated with antibiotics also showed similar but less severe changes with intermediate effects. That paper discussed the interaction among gut bacteria, immune cells, and epithelial cells through inflammatory mediators, cytokines, bioactive molecules, receptors, ligands, tran-



scription factors, and pathways, but failed to provide a coherent mechanistic explanation. Here I suggest that impaired inactivation of digestive proteases may be the key factor for the observed increased susceptibility. Studies have well documented that large amount of digestive proteases can be found in the large intestine of germ-free or antibiotics-treated animals but not in animals raised at conventional condition<sup>[2-7]</sup>. As we know, the gut is mainly protected by a layer of mucus secreted from goblet cells. Mucin, the structural molecule of mucus, contains a central core of peptide and side carbohydrate branches that can constitute up to 85% of the molecule. These carbohydrate side chains greatly retarded degradation of the protective mucus layer and damage of the underlying gut tissue by the digestive proteases being existed in germ-free or antibiotics-treated animals. However, the great destructive potency of these digestive proteases would be released and manifested once the gut barrier is broken as occurred either shortly after treatment with high concentrations of DSS or prolonged treatment with lower dose of DSS, resulting in the severe damage of epithelial cells at gut surface and then blood vessels in the mucosa, leading to excessive blood loss and enhanced mortality, even in the absence of remarkably inflammation. This would be similar to the so-called tryptic hemorrhagic necrosis of the gut as seen in animals under shock, which can be greatly prevented by previous inactivation of digestive proteases or ligation of pancreatic duct<sup>[8]</sup>. In fact, evidences I collected during the last fifteen years make me to believe that impaired inactivation of digestive proteases due to reduction in gut bacteria along with the improved hygiene and inhibition by some dietary chemicals such as saccharin and sucralose may have also played critical causative role in the pathogenesis of inflammatory bowel disease (IBD) in human<sup>[9-11]</sup>. Thus, impaired inactivation of digestive proteases in colitis and

IBD would be worthwhile for further study.

## REFERENCES

- 1 **Hernández-Chirilaque C**, Aranda CJ, Ocón B, Capitán-Cañadas F, Ortega-González M, Carrero JJ, Suárez MD, Zarzuelo A, Sánchez de Medina F, Martínez-Augustin O. Germ-free and Antibiotic-treated Mice are Highly Susceptible to Epithelial Injury in DSS Colitis. *J Crohns Colitis* 2016; **10**: 1324-1335 [PMID: 27117829 DOI: 10.1093/ecco-jcc/jjw096]
- 2 **Norin KE**. Influence of antibiotics on some intestinal microflora associated characteristics. *Anaerobe* 1997; **3**: 145-148 [PMID: 16887579 DOI: 10.1006/anae.1997.0091]
- 3 **Genell S**, Gustafsson BE. Impaired enteric degradation of pancreatic endopeptidases in antibiotic-treated rats. *Scand J Gastroenterol* 1977; **12**: 801-809 [PMID: 594651 DOI: 10.3109/00365527709181723]
- 4 **Genell S**, Gustafsson BE, Ohlsson K. Quantitation of active pancreatic endopeptidases in the intestinal contents of germfree and conventional rats. *Scand J Gastroenterol* 1976; **11**: 757-762 [PMID: 1006149]
- 5 **Gustafsson BE**. The physiological importance of the colonic microflora. *Scand J Gastroenterol Suppl* 1982; **77**: 117-131 [PMID: 6750828]
- 6 **Midtvedt T**, Carlstedt-Duke B, Höverstad T, Midtvedt AC, Norin KE, Saxerholt H. Establishment of a biochemically active intestinal ecosystem in ex-germfree rats. *Appl Environ Microbiol* 1987; **53**: 2866-2871 [PMID: 3124742]
- 7 **Reddy BS**, Pleasants JR, Wostmann BS. Pancreatic enzymes in germfree and conventional rats fed chemically defined, water-soluble diet free from natural substrates. *J Nutr* 1969; **97**: 327-334 [PMID: 5773333]
- 8 **Bounous G**. Metabolic changes in the intestinal mucosa during hemorrhagic shock. *Can J Surg* 1965; **8**: 332-339 [PMID: 5294426]
- 9 **Qin XF**. Impaired inactivation of digestive proteases by deconjugated bilirubin: the possible mechanism for inflammatory bowel disease. *Med Hypotheses* 2002; **59**: 159-163 [PMID: 12208202 DOI: 10.1016/S0306-9877(02)00243-8]
- 10 **Qin X**. Etiology of inflammatory bowel disease: a unified hypothesis. *World J Gastroenterol* 2012; **18**: 1708-1722 [PMID: 22553395 DOI: 10.3748/wjg.v18.i15.1708]
- 11 **Qin X**. Publications by Xiaofa Qin. [accessed 2016 Oct 8]. Available from: URL: <http://www.ncbi.nlm.nih.gov/pubmed/?term=xiaofa+qin>

P- Reviewer: Matowicka-Karna J, Yang CH, Zhu YL S- Editor: Qi Y  
L- Editor: A E- Editor: Li D



Observational Study

# Polyp detection rate and pathological features in patients undergoing a comprehensive colonoscopy screening

Hamid Asadzadeh Aghdaei, Ehsan Nazemalhosseini Mojarad, Sara Ashtari, Mohamad Amin Pourhoseingholi, Vahid Chaleshi, Fakhrosadat Anaraki, Mehrdad Haghazali, Mohammad Reza Zali

Hamid Asadzadeh Aghdaei, Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran 19835-178, Iran

Ehsan Nazemalhosseini Mojarad, Sara Ashtari, Mohamad Amin Pourhoseingholi, Vahid Chaleshi, Mehrdad Haghazali, Mohammad Reza Zali, Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran 19835-178, Iran

Fakhrosadat Anaraki, Colorectal Division of Department of Surgery, Taleghani Hospital, Shahid Beheshti University of Medical Sciences, Tehran 19835-178, Iran

**Author contributions:** Asadzadeh Aghdaei H, Haghazali M and Zali MR performed all colonoscopies; Asadzadeh Aghdaei H drafted the initial manuscript; Nazemalhosseini Mojarad E and Anaraki F were the guarantors and designed the study; Chaleshi V gathered the data; Ashtari S participated in the acquisition, analysis, and interpretation of the data, and wrote and revised the last version of manuscript; Pourhoseingholi MA as PhD of biostatistics revised the article for statistical materials and methods.

**Supported by** Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

**Institutional review board statement:** The study was reviewed and approved by the Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

**Informed consent statement:** All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

**Conflict-of-interest statement:** There are no conflicts of interest to report.

**Data sharing statement:** No additional data are available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Correspondence to:** Sara Ashtari, MSc of Epidemiology in Research Institute of Gastroenterology and Liver Diseases, Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tabnak St, Yaman Ave, Velenjak, Tehran 19835-178, Iran. [sara\\_ashtari@yahoo.com](mailto:sara_ashtari@yahoo.com)  
**Telephone:** +98-21-22432515  
**Fax:** +98-21-22432517

**Received:** September 13, 2016  
**Peer-review started:** September 14, 2016  
**First decision:** October 21, 2016  
**Revised:** November 23, 2016  
**Accepted:** December 7, 2016  
**Article in press:** December 9, 2016  
**Published online:** February 15, 2017

## Abstract

### AIM

To identify the prevalence, and clinical and pathologic characteristic of colonic polyps among Iranian patients undergoing a comprehensive colonoscopy, and determine the polyp detection rate (PDR) and adenoma detection rate (ADR).

## METHODS

In this cross-sectional study, demographics and epidemiologic characteristics of 531 persons who underwent colonoscopies between 2014 and 2015 at Mehrad gastrointestinal clinic were determined. Demographics, indication for colonoscopy, colonoscopy findings, number of polyps, and histopathological characteristics of the polyps were examined for each person.

## RESULTS

Our sample included 295 (55.6%) women and 236 (44.4%) men, with a mean age of  $50.25 \pm 14.89$  years. Overall PDR was 23.5% (125/531). ADR and colorectal cancer detection rate in this study were 12.8% and 1.5%, respectively. Polyps were detected more significantly frequently in men than in women (52.8% *vs* 47.2%,  $P < 0.05$ ). Polyps can be seen in most patients after the age of 50. The average age of patients with cancer was significantly higher than that of patients with polyps (61.3 years *vs* 56.4 years,  $P < 0.05$ ). The majority of the polyps were adenomatous. More than 50% of the polyps were found in the rectosigmoid part of the colon.

## CONCLUSION

The prevalence of polyps and adenomas in this study is less than that reported in the Western populations. In our patients, distal colon is more susceptible to developing polyps and cancer than proximal colon.

**Key words:** Adenoma detection; Polyp detection; Iran; Colonoscopy; Screening

© The Author(s) 2017. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** One of the major reasons for colonoscopy is detection of colon polyps, such as adenomas. Early diagnosis and endoscopic removal of adenomatous polyps is one of the main objectives for screening and prevention of colorectal cancer (CRC). Given that, only few studies are available in the national literature regarding the assessment of colorectal polyps, but none has explicitly noted the rate of polyp detection. Nevertheless, our study provides comprehensive information about clinical and epidemiological features of colorectal polyps. Therefore, the results of this study can provide a good infrastructure for the next preventive program and have clinical implications for CRC screening.

Asadzadeh Aghdaei H, Nazemalhosseini Mojarad E, Ashtari S, Pourhoseingholi MA, Chaleshi V, Anaraki F, Haghazali M, Zali MR. Polyp detection rate and pathological features in patients undergoing a comprehensive colonoscopy screening. *World J Gastrointest Pathophysiol* 2017; 8(1): 3-10 Available from: URL: <http://www.wjgnet.com/2150-5330/full/v8/i1/3.htm> DOI:

## INTRODUCTION

Colorectal cancer (CRC) is the third most common malignant disease in both men and women worldwide, accounting for more than 8% of mortality in the world with approximately 1.4 million new cases a year<sup>[1-4]</sup>.

CRC is also the third most common cancer in men and women in Asia<sup>[5,6]</sup>. In the Asia-Pacific region, the incidence varies between regions, with high incidence in Australia, and Eastern Asia, and low incidence in South-central Asia<sup>[7]</sup>. According to the International Agency for Research on Cancer, the incidence of CRC in many Asian countries is similar to that in many Western countries<sup>[8]</sup>. CRC is also the third most common cancer in Iranians, after excluding skin cancer, and it occurs at younger ages with an increasing trend similar to that in the Asia-Pacific countries<sup>[1,9]</sup>. These increasing rates may result from the young age structure and low rates of colon cancer in older people of these countries<sup>[6,10,11]</sup>.

Almost all CRCs develop from colorectal polyps. Over a period of ten years, most of adenomatous polyps can be converted to colon carcinoma<sup>[12,13]</sup>. Given that the process of conversion of colorectal adenomas into adenocarcinoma is very long and slow<sup>[14]</sup>, early detection and endoscopic removal of these precancerous lesions are very effective in reducing the incidence and mortality rate of CRC<sup>[15-17]</sup>.

CRC is a suitable disease for screening<sup>[18]</sup>. However, due to a lack of comprehensive screening strategy and public acceptance, this program is not implemented in many countries. Nevertheless, access to the CRC screening is an important key to reducing the burden of CRC. Endoscopic screening is comprised of four techniques including sigmoidoscopy, colonoscopy, barium enema, and computed tomographic colonography<sup>[19]</sup>. Colonoscopy is a highly specific and the most effective screening tool for detecting colonic polyps and CRC<sup>[20]</sup>.

Limited data are available in the national literature regarding the assessment of colorectal polyps<sup>[21-25]</sup>. Understanding of the prevalence of colorectal polyps especially adenomas in the general population would help clarify the efficacy of a CRC screening program. Therefore, updating the current knowledge in the scope of colorectal polyps and CRC is essential. Hence, identifying the features of colon polyps (*e.g.*, distribution, location, and histology type) has great implications for developing national screening guidelines for CRC<sup>[26,27]</sup>. In this study, we aimed to determine the baseline polyp and adenoma prevalence in persons who underwent colonoscopies for various indications as well as opportunistic screening for CRC. We also assessed the polyp detection rate (PDR) and adenoma detection rate (ADR), and evaluated the clinical and histological

**Table 1** Patients' characteristics and colonoscopy findings

Variable	All (n = 531)
Sex	
Male, n (%)	236 (44.4)
Female, n (%)	295 (55.6)
Age	
Mean years (SD)	50.3 ± 15.4
Age groups	
≤ 50 yr	250 (47.1)
> 50 yr	281 (52.9)
Family history	
Yes	150 (28.2)
No	381 (71.8)
Indication, n (%)	
Screening	120 (22.6)
Gastrointestinal bleeding	90 (17.0)
Constipation	82 (15.5)
Diarrhea	74 (14.0)
Abdominal pain	70 (13.3)
Inflammatory bowel disease	65 (12.2)
Others	29 (5.4)
Patients with at least 1 polyp, n (%)	
Neoplastic polyps	68 (54.4)
Non-neoplastic polyps	57 (45.6)
Cancer, n (%)	8 (1.5)

characteristics of colorectal polyps in Iranian patients and Iranian volunteers for CRC screening.

## MATERIALS AND METHODS

### Study design

In this cross-sectional study, all data were extracted from a colonoscopy database and pathology reports maintained by Mehrad gastrointestinal clinic in Iran. We included all persons aged 15 to 85 years, who underwent their first time colonoscopy during 2014-2015. Patients who had previously been identified with colon polyp or colorectal malignancies including CRC, colonic resection, active colitis, active diverticulitis and familial adenomatous polyposis were excluded from the study. We collected the data on demographic variables, indications for colonoscopy, and family history of colorectal malignancies. Family history was defined as having a first degree relative with CRC. For all colorectal lesions, data on clinical and pathological features (*i.e.*, number, size, site, and grade of dysplasia) were obtained. The study was approved by the Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences.

### Polyp classification

All polyps identified during colonoscopy were biopsied or removed endoscopically and submitted for histopathology. The overall PDR was defined as the proportion of procedures in which at least one polyp was detected over the total number of colonoscopies. ADR was defined as the number of colonoscopies in which one or more adenomas was detected, divided by the total number of colonoscopies performed by the endoscopist<sup>[28]</sup>.

Pathological features of colorectal lesions were determined using the World Health Organization criteria<sup>[28]</sup>. Hyperplastic and adenomatous polyps were classified as non-neoplastic polyps and neoplastic polyps, respectively. Microscopically, adenomas were categorized architecturally as serrated, tubular, tubular-villous, and villous.

The locations of the polyps were defined as proximal colon including the transverse colon, hepatic flexure, ascending colon, and cecum, and distal colon including the rectum, sigmoid, descending colon, and splenic flexure.

The polyp size was classified as small (< 5 mm), medium (5-9 mm), or large (> 10 mm). Estimation of polyp size was performed by the endoscopist using the diameter of the open biopsy forceps, which is about 8 mm. In the event of multiple polyps, only the size of the largest was considered for the purposes of analysis. Degrees of dysplasia observed in the adenomas were graded as low (mild and moderate) or high grade (severe). Patients with no polyps were regarded as normal.

### Statistical analysis

Categorical variables are expressed as numbers and percentages.  $\chi^2$  or Fisher's exact test, where appropriate, was used for analysis of categorical variables. Continuous variables are expressed as medians, or as means and standard deviation, and 95%CI as appropriate. The Student's *t*-test was used for comparisons of means. All analyses were performed using SPSS version 21.0 (SPSS INC, Chicago, IL, United States). A two-tailed *P* < 0.05 was considered statistically significant.

## RESULTS

### Demographic and historical data

During the period of study (2014-2015), 531 persons met the inclusion criteria. Our sample included 55.6% of women and 44.4% of men. The age range was 15-85 years with a mean of 50.3 ± 15.4 years and a median of 52 years. The patients were divided into two age-groups (≤ 50 and > 50 years). The majority of patients was over 50 years (52.9%). One hundred and fifty (28.2%) patients had a family history of polyps or CRC in our study.

The most common reasons for colonoscopy included screening (asymptomatic adults aged 50 years and older and with a family history of CRC) in 22.6%, and lower gastrointestinal bleeding in 17.0%. Other indications for colonoscopy were constipation (15.4%), diarrhea (13.9%), abdominal pain (13.1%), inflammatory bowel disease (12.2%), and others (5.4%). Other referral indications included bloating, reflux, weight loss, anemia, fatty liver, and irritable bowel syndrome (Table 1).

### Study outcomes and colonoscopy findings

Based on colonoscopy findings, the overall PDR was



**Table 2** Detection rates of different histologic types of polyps and cancer by gender

Histologic type of polyps	Gender		Total
	Male (%)	Female (%)	
Neoplastic			
Tubular	18 (53.0)	16 (47.0)	34 (100)
Tubulo-villous	13 (68.4)	6 (31.6)	19 (100)
Villous	4 (40.0)	6 (60.0)	10 (100)
Serrated	3 (60.0)	2 (40.0)	5 (100)
Adenomatous polyps	38 (55.9)	30 (44.1)	68 (100)
Non-neoplastic			
Hyperplastic	28 (40.6)	29 (45.3)	57 (100)
Total PDR	66 (52.8)	59 (47.2)	125 (100)
Cancer	5 (62.5)	3 (37.5)	8 (100)
Total	69 (51.9)	64 (48.1)	133 (100)

PDR: Polyp detection rate.

**Table 3** Detection rates of different histologic types of polyps and cancer by age

Histologic type of polyps	Age-groups		Total
	≤ 50 yr	> 50 yr	
Neoplastic			
Tubular	6 (17.6)	28 (82.4)	34 (100)
Tubulo-villous	7 (36.8)	12 (63.2)	19 (100)
Villous	3 (30.0)	7 (70.0)	10 (100)
Serrated	3 (60.0)	2 (40.0)	5 (100)
Adenomatous polyps	19 (28.0)	49 (72.0)	68 (100)
Non-neoplastic			
Hyperplastic	19 (33.3)	38 (66.7)	57 (100)
Total PDR	38 (30.4)	87 (69.6)	125 (100)
Cancer	1 (12.5)	7 (87.5)	8 (100)
Total	39 (29.3)	94 (70.7)	133 (100)

PDR: Polyp detection rate.

**Table 4** Detection rates of different histologic types of polyps and cancer by family history

Histologic type of polyps	Family history		Total
	No (%)	Yes (%)	
Neoplastic			
Tubular	26 (76.4)	8 (23.6)	34 (100)
Tubulo-villous	13 (68.4)	6 (31.6)	19 (100)
Villous	5 (50.0)	5 (50.0)	10 (100)
Serrated	2 (40.0)	3 (60.0)	5 (100)
Adenomatous polyps	46 (67.6)	22 (32.4)	68 (100)
Non-neoplastic			
Hyperplastic	46 (80.7)	11 (19.3)	57 (100)
Total PDR	92 (73.6)	33 (26.4)	125 (100)
Cancer	0 (0)	8 (100)	8 (100)
Total	92 (69.2)	41 (30.8)	133 (100)

PDR: Polyp detection rate.

**Table 5** Detection rates of different histologic types of polyps by location

Histologic type of polyps	Location		Total
	Proximal colon (%)	Distal colon (%)	
Neoplastic			
Tubular	10 (26.3)	28 (73.7)	38 (100)
Tubulo-villous	9 (40.9)	13 (59.1)	22 (100)
Villous	3 (30.0)	7 (70.0)	10 (100)
Serrated	3 (37.5)	5 (62.5)	8 (100)
Adenomatous polyps	25 (32.0)	53 (68.0)	78 (100)
Non-neoplastic			
Hyperplastic	19 (31.7)	41 (68.3)	60 (100)
Total PDR	44 (31.9)	94 (68.1)	138 (100)

PDR: Polyp detection rate.

23.5% (125/531). According to histopathology results from 125 patients with at least one polyp, 54.4% of the lesions were neoplastic polyps and 45.6% were hyperplastic polyps (non-neoplastic polyps). The overall ADR in this study was 12.8% (68/531). The percentage of male patients with polyps was significantly higher than that of female patients (52.8% vs 47.2%,  $P < 0.05$ ). CRC was detected in 1.5% (8/531) of the total population (men, 62.5%; women, 37.5%) (Table 2). The mean age of patients with polyp was  $56.4 \pm 13.5$  years. Polyps can be seen in most patients after the age of 50 (69.6%). CRC was more frequently observed in patients aged from 60 to 80 years. The average age of patients with cancer was significantly higher than that of those with polyps ( $61.3 \pm 19.7$  years vs  $56.4 \pm 13.5$  years,  $P < 0.05$ ) (Table 3). In addition, the relationship between PDR and family history of CRC or polyps was assessed and this was found not to be statistically significant ( $P > 0.05$ ), while all patients with CRC in this study had a family history (Table 4).

#### Histopathological characteristics of the polyps

Totally 138 polyps were removed by colonoscopy in this

study. Among the 138 polyps, 56.5% were neoplastic and the others were non-neoplastic. All non-neoplastic polyps were hyperplastic. Among the 78 neoplastic polyps, the common histologic types were tubular adenomas (48.7%), tubule-villous adenomas (28.2%), villous adenomas (12.8%), and serrated adenomas (10.2%).

The prevalence of polyps in distal colon was higher than that in proximal colon (68.1% vs 31.9%,  $P < 0.05$ ). Accordingly, most of cancers were located in the distal colon compared with the proximal colon (75% vs 25%,  $P < 0.05$ ) (Table 5). Overall polyps were frequently detected in the rectum (32.0%), sigmoid (24.6%), transverse colon (16%), and ascending and descending colon (10.1%), and the others located in the cecum (5%), and splenic and hepatic flexure (2.1%) (Table 6).

Data about the size of polyps were available for only 75 polyps; 33.3% were smaller than 5 mm as small size, 40% were between 5–9 mm as medium size and 26.7% were more than 10 mm as large size (Table 7). According to the degree of dysplasia observed in the adenomas, most of patients (52.6%) had mild grade, 24.4% had moderate, and 23% had severe grade of dysplasia (Table 8).

**Table 6** Detection rates of different histologic types of polyps by colonic segments

Polyps	Location								Total
	Ascending colon	Transverse colon	Descending colon	Sigmoid	Rectum	Cecum	Hepatic flexure	Splenic flexure	
Tubular	3 (7.9)	5 (13.2)	4 (10.5)	7 (18.4)	16 (42.1)	1 (2.6)	1 (2.6)	1 (2.6)	38 (100)
Tubulo-villous	4 (18.2)	3 (13.6)	2 (9.1)	6 (27.3)	5 (22.7)	2 (9.1)	0 (0)	0 (0)	22 (100)
Villous	0 (0)	1 (10)	0 (0)	2 (20)	5 (50)	2 (20)	0 (0)	0 (0)	10 (100)
Serrated	0 (0)	3 (37.5)	1 (12.5)	3 (37.5)	1 (12.5)	0 (0)	0 (0)	0 (0)	8 (100)
Hyperplastic	7 (11.7)	10 (16.7)	7 (11.7)	16 (26.7)	17 (28.3)	2 (3.3)	0 (0)	1 (1.7)	60 (100)
Total	14 (10.1)	22 (16.0)	14 (10.1)	34 (24.6)	44 (32.0)	7 (5.1)	1 (0.7)	2 (1.4)	138 (100)

**Table 7** Detection rates of different histologic types of polyps by size

Size	Histologic type					Total
	Hyperplastic	Tubular	Tubul-villous	villous	Serrated	
Small	11 (44.0)	0 (0)	14 (56.0)	0 (0)	0 (0)	25 (100)
Medium	14 (46.7)	0 (0)	12 (40.0)	0 (0)	4 (13.3)	30 (100)
Large	0 (0)	8 (40.0)	3 (15.0)	9 (45.0)	0 (0)	20 (100)
Unknown	35 (55.6)	2 (3.2)	9 (14.3)	13 (20.6)	4 (6.3)	63 (100)
Total	60 (43.5)	10 (7.2)	38 (27.5)	22 (16.0)	8 (5.8)	138 (100)

Data available for only 75 polyps.

**Table 8** Detection rates of different histologic types of polyps by degree of dysplasia

Grade	Histologic type				Total
	Tubular	Tubul-villous	Villous	Serrated	
Mild	21 (51.2)	11 (26.9)	6 (14.6)	3 (7.3)	41 (100)
Moderate	11 (57.9)	3 (15.8)	3 (15.8)	2 (10.5)	19 (100)
Severe	6 (33.3)	8 (44.5)	1 (5.5)	3 (16.7)	18 (100)
Total	38 (48.7)	22 (28.2)	10 (12.9)	8 (10.2)	78 (100)

## DISCUSSION

The PDR and ADR rates obtained in this study are low, when compared to the figures from most Western and some Asian countries. In a large multicenter study from Italy, the median detection rate for polyps was 35%<sup>[29]</sup>. A large colonoscopy series from Spain reported a PDR of 45.8%<sup>[30]</sup>. Similar studies from Mayo Clinic in the United States and France reported PDR of 49% and 35.5%, and ADR of 31% and 17.7%, respectively<sup>[31,32]</sup>. In some Asian countries like Korea, China and Thailand, PDR and ADR were similar to those in Europeans and Americans<sup>[33-36]</sup>. However, our findings are similar to reports from Kuwait, Malaysia and Oman where PDR of 20% and ADR of 10%, 11.5% and 12.1% were reported, respectively<sup>[37-39]</sup>. While in African countries like Nigeria these rates were reported to be lower than our results (PDR, 16.1%; ADR, 6.8%)<sup>[40]</sup>. The mean age of the studied population was relatively young (50.3 years) and it might decrease the ADR because adenomas have been demonstrated to be more frequent in those older than 50 years of age.

The overall estimate for PDR in our study was 23.5%, while ADR was 12.8%. The percentage of male patients with polyps was significantly higher than that of female patients (52.8% vs 47.2%,  $P < 0.05$ ). These data are consistent with the other reports that support

gender differences in the prevalence of colon polyps and cancer<sup>[27,40,41]</sup>.

Older age is the most important predictor for the prevalence of adenomas and cancer. In our study, the PDR and cancer prevalence reached a peak in the 6<sup>th</sup> decade of life. Nonetheless, the average age of patients with cancer was significantly higher than that of patients with polyps (61.3 years vs 56.4 years,  $P < 0.05$ ). Studies from the Middle East and the Western countries also mentioned a significant increase in the risk of CRC, in particular after the age of 50 years<sup>[24,28,42]</sup>.

Based on increasing prevalence of CRC in the sixth decade of life, regular screening should begin at the age of 50. However, people at higher risk of developing CRC should begin screening at a younger age<sup>[43,44]</sup>.

The tubular type was the most common histological feature of adenomas in the present study, in accordance with the results of other reports<sup>[21,27,45]</sup>. Polyps were detected significantly in distal (left sided) colon, comparable with results from Asian and the Western countries<sup>[46,47]</sup>. Nonetheless, because of the significance of adenomatous polyps and hyperplastic polyps present in proximal colon, the location of polyps is not helpful for distinguishing between neoplastic and non-neoplastic polyps. So, complete colonoscopy is recommended in screening guidelines for colon cancer<sup>[43,48]</sup>. In addition, this study showed that only 56.6% of the polyps were found in the rectum and sigmoid region. Our study did not find any association between the age and location of polyps. This is in contrast with previous studies showing that the incidence of right sided polyps increased with increasing age<sup>[44]</sup>.

With regard to the size of polyps, we observed that the hyperplastic polyps and tubular adenoma were always smaller than 10 mm, while tubulo-villous and villous adenoma were always bigger than 10 mm. So,

removing the polyps which are larger than 10 mm is recommended<sup>[36,49]</sup>.

The study faced some limitations. First, this study was not population-based, therefore, the selection bias of the study population must be kept in mind. Second, our sample included mostly symptomatic patients, in which the estimates may be different from screening studies with asymptomatic individuals. Nevertheless, the results of this study can provide a good infrastructure for the next preventive program and have clinical implications for CRC screening.

In conclusion, PDR, ADR and CRC detection rate in this study were 23.5%, 12.8% and 1.5%, respectively. Most of the polyps and CRC were identified in patients aged 50 years or older. The majority of the polyps were adenomatous. More than 50% of the polyps were found in the rectosigmoid part of the colon. Finally, our study did not find any association between the family history and PDR.

## COMMENTS

### Background

Colorectal cancer (CRC) is the third most common malignant disease in both men and women worldwide, accounting for more than 8% of cancer-related death in the world with approximately 1.4 million new cases a year. Almost all CRCs develop from colorectal polyps. CRC largely can be prevented by the detection and removal of adenomatous polyps, and survival is significantly better when CRC is diagnosed while still localized. When CRC is found at an early stage before it has spread, the 5-year relative survival rate is about 90%. But only about 4 out of 10 CRCs are found at this early stage. When cancer has spread outside the colon or rectum, survival rates are lower.

### Research frontiers

Early diagnosis and endoscopic resection of adenomatous polyps is the main approach for screening and prevention of CRC. This study aimed at identifying the prevalence, and clinical and pathologic characteristic of colonic polyps among Iranian patients undergoing a comprehensive colonoscopy and determine the polyp detection rate (PDR) and adenoma detection rate (ADR).

### Innovations and breakthroughs

Only few studies are available in the national literature regarding the assessment of colorectal polyps, but none has explicitly noted the rate of polyp detection. Nevertheless, the study provides comprehensive information about clinical and epidemiological features of colorectal polyps.

### Applications

Older age is the most important predictor for the prevalence of adenomas and cancer. Based on the results of this study, the PDR and cancer prevalence reached a peak in the 6<sup>th</sup> decade of life. Given the increased prevalence of CRC in the sixth decade of life, regular screening beginning at the age of 50 is the key to preventing CRC.

### Terminology

The overall PDR was defined as the proportion of procedures in which at least one polyp was detected over the total number of colonoscopies. ADR was defined as the number of colonoscopies in which one or more adenomas was detected, divided by the total number of colonoscopies performed by the endoscopist.

### Peer-review

Authors report in this paper the detection rates by colonoscopy for cancer and adenoma in an Iranian population.

## REFERENCES

- 1 **Jemal A**, Center MM, DeSantis C, Ward EM. Global patterns of cancer incidence and mortality rates and trends. *Cancer Epidemiol Biomarkers Prev* 2010; **19**: 1893-1907 [PMID: 20647400 DOI: 10.1158/1055-9965.epi-10-0437]
- 2 **Edwards BK**, Ward E, Kohler BA, Ehemann C, Zauber AG, Anderson RN, Jemal A, Schymura MJ, Lansdorp-Vogelaar I, Seeff LC, van Ballegooijen M, Goede SL, Ries LA. Annual report to the nation on the status of cancer, 1975-2006, featuring colorectal cancer trends and impact of interventions (risk factors, screening, and treatment) to reduce future rates. *Cancer* 2010; **116**: 544-573 [PMID: 19998273 DOI: 10.1002/cncr.24760]
- 3 **Ferlay J**, Autier P, Boniol M, Heanue M, Colombet M, Boyle P. Estimates of the cancer incidence and mortality in Europe in 2006. *Ann Oncol* 2007; **18**: 581-592 [PMID: 17287242 DOI: 10.1093/annonc/mdl498]
- 4 **Joseph DA**, King JB, Miller JW, Richardson LC. Prevalence of colorectal cancer screening among adults--Behavioral Risk Factor Surveillance System, United States, 2010. *MMWR Suppl* 2012; **61**: 51-56 [PMID: 22695464]
- 5 **Pourhoseingholi MA**. Increased burden of colorectal cancer in Asia. *World J Gastrointest Oncol* 2012; **4**: 68-70 [PMID: 22532878 DOI: 10.4251/wjgo.v4.i4.68]
- 6 **Yiu HY**, Whittemore AS, Shibata A. Increasing colorectal cancer incidence rates in Japan. *Int J Cancer* 2004; **109**: 777-781 [PMID: 14999789 DOI: 10.1002/ijc.20030]
- 7 **Pourhoseingholi MA**. Epidemiology and burden of colorectal cancer in Asia-Pacific region: what shall we do now? *Transl Gastrointest Cancer* 2014; **3**: 169-173 [DOI: 10.3978/j.issn.2224-4778.2014.08.10]
- 8 **Ferlay J**, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; **136**: E359-E386 [PMID: 25220842 DOI: 10.1002/ijc.29210]
- 9 **Hossein Somi M**, Mirinezhad K, Farhang S, Jazayeri E, Sani A, Seif-Farshadi M, Golzari M, Kashaf S, Sadegy M. Gastrointestinal cancer occurrence in East Azarbaijan: a five year study from North Western Iran. *Asian Pac J Cancer Prev* 2006; **7**: 309-312 [PMID: 16839228]
- 10 **Mousavi SM**, Gouya MM, Ramazani R, Davanlou M, Hajsadeghi N, Seddighi Z. Cancer incidence and mortality in Iran. *Ann Oncol* 2009; **20**: 556-563 [PMID: 19073863 DOI: 10.1093/annonc/mdn642]
- 11 **Ansari R**, Mahdavinia M, Sadjadi A, Nouraie M, Kamangar F, Bishehsari F, Fakheri H, Semnani S, Arshi S, Zahedi MJ, Darvish-Moghadam S, Mansour-Ghanaei F, Mosavi A, Malekzadeh R. Incidence and age distribution of colorectal cancer in Iran: results of a population-based cancer registry. *Cancer Lett* 2006; **240**: 143-147 [PMID: 16288832 DOI: 10.1016/j.canlet.2005.09.004]
- 12 **Noffsinger AE**. Serrated polyps and colorectal cancer: new pathway to malignancy. *Annu Rev Pathol* 2009; **4**: 343-364 [PMID: 19400693 DOI: 10.1146/annurev.pathol.4.110807.092317]
- 13 **Levine JS**, Ahnen DJ. Clinical practice. Adenomatous polyps of the colon. *N Engl J Med* 2006; **355**: 2551-2557 [PMID: 17167138 DOI: 10.1056/NEJMcP063038]
- 14 **Huang CS**, Farraye FA, Yang S, O'Brien MJ. The clinical significance of serrated polyps. *Am J Gastroenterol* 2011; **106**: 229-240; quiz 241 [PMID: 21045813 DOI: 10.1038/ajg.2010.429]
- 15 **García Sánchez J**. [Colonoscopic polypectomy and long-term prevention of colorectal cancer deaths]. *Rev Clin Esp* 2012; **212**: 408 [PMID: 22937540 DOI: 10.1016/j.rce.2012.05.002]
- 16 **Espey DK**, Wu XC, Swan J, Wiggins C, Jim MA, Ward E, Wingo PA, Howe HL, Ries LA, Miller BA, Jemal A, Ahmed F, Cobb N, Kaur JS, Edwards BK. Annual report to the nation on the status of cancer, 1975-2004, featuring cancer in American Indians and Alaska Natives. *Cancer* 2007; **110**: 2119-2152 [PMID: 17939129 DOI: 10.1002/cncr.23044]

- 17 **Atkin WS**, Edwards R, Kralj-Hans I, Wooldrage K, Hart AR, Northover JM, Parkin DM, Wardle J, Duffy SW, Cuzick J. Once-only flexible sigmoidoscopy screening in prevention of colorectal cancer: a multicentre randomised controlled trial. *Lancet* 2010; **375**: 1624-1633 [PMID: 20430429 DOI: 10.1016/s0140-6736(10)60551-x]
- 18 **Inadomi JM**, Vijan S, Janz NK, Fagerlin A, Thomas JP, Lin YV, Muñoz R, Lau C, Somsouk M, El-Nachef N, Hayward RA. Adherence to colorectal cancer screening: a randomized clinical trial of competing strategies. *Arch Intern Med* 2012; **172**: 575-582 [PMID: 22493463 DOI: 10.1001/archinternmed.2012.332]
- 19 **Elmunzer BJ**, Hayward RA, Schoenfeld PS, Saini SD, Deshpande A, Waljee AK. Effect of flexible sigmoidoscopy-based screening on incidence and mortality of colorectal cancer: a systematic review and meta-analysis of randomized controlled trials. *PLoS Med* 2012; **9**: e1001352 [PMID: 23226108 DOI: 10.1371/journal.pmed.1001352]
- 20 **Irvani S**, Nazemalhosseini-Mojarad E, Kashfi SM, Azimzadeh P. Screening of colorectal diseases among individuals without family history in a private hospital, Tehran, Iran from 2011 to 2013. *Transl Gastrointest Cancer* 2014; **3**: 165-168 [DOI: 10.3978/j.issn.2224-4778.2014.08.04]
- 21 **Mirzaie AZ**, Abolhasani M, Moghaddam RM, Kabivar M. The Frequency of gastrointestinal polyps in Iranian population. *Iran J Pathol* 2012; **7**: 183-189
- 22 **Bafandeh Y**, Khosbaten M, Eftekhari Sadat AT, Farhang S. Clinical predictors of colorectal polyps and carcinoma in a low prevalence region: results of a colonoscopy based study. *World J Gastroenterol* 2008; **14**: 1534-1538 [PMID: 18330943 DOI: 10.3748/wjg.14.1534]
- 23 **Bafandeh Y**, Daghestani D, Esmaili H, Aharizad S. Distribution of cancer and adenomatous polyps in the colorectum: study in an Iranian population. *Asian Pac J Cancer Prev* 2006; **7**: 65-68 [PMID: 16629518]
- 24 **Bafandeh Y**, Daghestani D, Esmaili H. Demographic and anatomical survey of colorectal polyps in an Iranian population. *Asian Pac J Cancer Prev* 2005; **6**: 537-540 [PMID: 16436008]
- 25 **Khatibzadeh N**, Ziaee SA, Rahbar N, Molanie S, Arefian L, Fanaie SA. The indirect role of site distribution in high-grade dysplasia in adenomatous colorectal polyps. *J Cancer Res Ther* 2005; **1**: 204-207 [PMID: 17998654 DOI: 10.4103/0973-1482.19587]
- 26 **Eshghi MJ**, Fatemi R, Hashemy A, Aldulaimi D, Khodadoostan M. A retrospective study of patients with colorectal polyps. *Gastroenterol Hepatol Bed Bench* 2011; **4**: 17-22 [PMID: 24834150]
- 27 **Delavari A**, Mardan F, Salimzadeh H, Bishehsari F, Khosravi P, Khanezhad M, Nasser-Moghaddam S, Merat S, Ansari R, Vahedi H, Shahbazzkhani B, Saberifirooz M, Sotoudeh M, Malekzadeh R. Characteristics of colorectal polyps and cancer; a retrospective review of colonoscopy data in Iran. *Middle East J Dig Dis* 2014; **6**: 144-150 [PMID: 25093062]
- 28 **Heitman SJ**, Ronksley PE, Hilsden RJ, Manns BJ, Rostom A, Hemmelgarn BR. Prevalence of adenomas and colorectal cancer in average risk individuals: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2009; **7**: 1272-1278 [PMID: 19523536 DOI: 10.1016/j.cgh.2009.05.032]
- 29 **Ricci E**, Hassan C, Petruzzello L, Bazzoli F, Repici A, Di Giulio E. Inter-centre variability of the adenoma detection rate: a prospective, multicentre study. *Dig Liver Dis* 2013; **45**: 1022-1027 [PMID: 23816699 DOI: 10.1016/j.dld.2013.05.009]
- 30 **Lucendo AJ**, Guagnozzi D, Angueira T, González-Castillo S, Fernández-Fuente M, Friginal-Ruiz AB, Tenias JM. The relationship between proximal and distal colonic adenomas: is screening sigmoidoscopy enough in the presence of a changing epidemiology? *Eur J Gastroenterol Hepatol* 2013; **25**: 973-980 [PMID: 23571611 DOI: 10.1097/MEG.0b013e3283614b57]
- 31 **Boroff ES**, Gurudu SR, Hentz JG, Leighton JA, Ramirez FC. Polyp and adenoma detection rates in the proximal and distal colon. *Am J Gastroenterol* 2013; **108**: 993-999 [PMID: 23567353 DOI: 10.1038/ajg.2013.68]
- 32 **Barret M**, Boustiere C, Canard JM, Arpurt JP, Bernardini D, Bulois P, Chaussade S, Heresbach D, Joly I, Lapuelle J, Laugier R, Lesur G, Pienkowski P, Ponchon T, Pujol B, Richard-Molard B, Robaszkiewicz M, Systchenko R, Abbas F, Schott-Pethelaz AM, Cellier C. Factors associated with adenoma detection rate and diagnosis of polyps and colorectal cancer during colonoscopy in France: results of a prospective, nationwide survey. *PLoS One* 2013; **8**: e68947 [PMID: 23874822 DOI: 10.1371/journal.pone.0068947]
- 33 **Chung SJ**, Kim YS, Yang SY, Song JH, Park MJ, Kim JS, Jung HC, Song IS. Prevalence and risk of colorectal adenoma in asymptomatic Koreans aged 40-49 years undergoing screening colonoscopy. *J Gastroenterol Hepatol* 2010; **25**: 519-525 [PMID: 20370730 DOI: 10.1111/j.1440-1746.2009.06147.x]
- 34 **Choi YS**, Suh JP, Lee DS, Youk EG, Lee IT, Lee SH, Kim DS, Lee DH. Colonoscopy screening for individuals aged 40-49 years with a family history of stomach cancer in Korea. *Int J Colorectal Dis* 2010; **25**: 443-447 [PMID: 20012440 DOI: 10.1007/s00384-009-0855-3]
- 35 **Leung WK**, Tang V, Lui PC. Detection rates of proximal or large serrated polyps in Chinese patients undergoing screening colonoscopy. *J Dig Dis* 2012; **13**: 466-471 [PMID: 22908972 DOI: 10.1111/j.1751-2980.2012.00621.x]
- 36 **Aswakul P**, Prachayakul V, Lohsiriwat V, Bunyaarunnate T, Kachintorn U. Screening colonoscopy from a large single center of Thailand - something needs to be changed? *Asian Pac J Cancer Prev* 2012; **13**: 1361-1364 [PMID: 22799332]
- 37 **Al-Enezi SA**, Alsurayei SA, Ismail AE, Aly NY, Ismail WA, Abou-Bakr AA. Adenomatous colorectal polyps in patients referred for colonoscopy in a regional hospital in Kuwait. *Saudi J Gastroenterol* 2010; **16**: 188-193 [PMID: 20616414 DOI: 10.4103/1319-3767.65194]
- 38 **Rajendra S**, Ho JJ, Arokiasamy J. Risk of colorectal adenomas in a multiethnic Asian patient population: race does not matter. *J Gastroenterol Hepatol* 2005; **20**: 51-55 [PMID: 15610446 DOI: 10.1111/j.1440-1746.2004.03522.x]
- 39 **Ashktorab H**, Brim H, Al-Riyami M, Date A, Al-Mawaly K, Kashoub M, Al-Mjeni R, Smoot DT, Al-Moundhri M, Al-Hashemi S, Ganguly SS, Raeburn S. Sporadic colon cancer: mismatch repair immunohistochemistry and microsatellite instability in Omani subjects. *Dig Dis Sci* 2008; **53**: 2723-2731 [PMID: 18299982 DOI: 10.1007/s10620-007-0189-3]
- 40 **Alatise OI**, Arigbabu AO, Agbakwuro AE, Lawal OO, Sowande OA, Odujoko OO, Adegoke O, Ojo O. Polyp prevalence at colonoscopy among Nigerians: A prospective observational study. *Niger J Clin Pract* 2014; **17**: 756-762 [PMID: 25385915 DOI: 10.4103/1119-3077.144391]
- 41 **Omranipour R**, Doroudian R, Mahmoodzadeh H. Anatomical distribution of colorectal carcinoma in Iran: a retrospective 15-yr study to evaluate rightward shift. *Asian Pac J Cancer Prev* 2012; **13**: 279-282 [PMID: 22502685]
- 42 **Nam JH**, Yang CH. [Clinical characteristics and risk factors of colon polyps in Gyeongju and Pohang area]. *Korean J Gastroenterol* 2008; **52**: 142-149 [PMID: 19077510]
- 43 **Sung JJ**, Ng SC, Chan FK, Chiu HM, Kim HS, Matsuda T, Ng SS, Lau JY, Zheng S, Adler S, Reddy N, Yeoh KG, Tsoi KK, Ching JY, Kuipers EJ, Rabeneck L, Young GP, Steele RJ, Lieberman D, Goh KL. An updated Asia Pacific Consensus Recommendations on colorectal cancer screening. *Gut* 2015; **64**: 121-132 [PMID: 24647008 DOI: 10.1136/gutjnl-2013-306503]
- 44 **Okamoto M**, Shiratori Y, Yamaji Y, Kato J, Ikenoue T, Togo G, Yoshida H, Kawabe T, Omata M. Relationship between age and site of colorectal cancer based on colonoscopy findings. *Gastrointest Endosc* 2002; **55**: 548-551 [PMID: 11923770 DOI: 10.1067/mge.2002.122335]
- 45 **Khodadoostan M**, Fatemi R, Maserat E, Hooshang A, Alizade M, Molaei M, Mashaieky R, Moaddab Y, Poor SY, Hashemy A, Zali MR. Clinical and pathological characteristics of colorectal polyps in Iranian population. *East Afr J Public Health* 2010; **7**: 157-159 [PMID: 21413595]
- 46 **Almadi MA**, Alharbi O, Azzam N, Wadera J, Sadaf N, Aljebreen AM. Prevalence and characteristics of colonic polyps and adenomas in 2654 colonoscopies in Saudi Arabia. *Saudi J Gastroenterol* 2014; **20**: 154-161 [PMID: 24976278 DOI: 10.4103/1319-3767.132986]
- 47 **Kim JH**, Choi YJ, Kwon HJ, Park SJ, Park MI, Moon W, Kim SE. Simple colonoscopy reporting system checking the detection



rate of colon polyps. *World J Gastroenterol* 2015; **21**: 9380-9386 [PMID: 26309364 DOI: 10.3748/wjg.v21.i31.9380]

- 48 **Mehran A**, Jaffe P, Efron J, Vernava A, Liberman A. Screening colonoscopy in the asymptomatic 50- to 59-year-old population. *Surg Endosc* 2003; **17**: 1974-1977 [PMID: 14569451 DOI:

10.1007/s00464-003-8807-4]

- 49 **Hassan C**, Pickhardt PJ, Rex DK. A resect and discard strategy would improve cost-effectiveness of colorectal cancer screening. *Clin Gastroenterol Hepatol* 2010; **8**: 865-869 [PMID: 20621680 DOI: 10.1016/j.cgh.2010.05.018]

**P- Reviewer:** Rey JF, Sali L   **S- Editor:** Ji FF   **L- Editor:** Wang TQ  
**E- Editor:** Li D





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

