

# World Journal of *Dermatology*

*World J Dermatol* 2023 February 6; 11(2): 7-29



**SYSTEMATIC REVIEWS**

- 7 Systematic review of hematidrosis: Time for clinicians to recognize this entity

*Octavius GS, Meliani F, Heriyanto RS, Yanto TA*

**ABOUT COVER**

Editorial Board Member of *World Journal of Dermatology*, Meri K Tulic, PhD, Professor, Research Scientist, INSERM U1065, Mediterranean Centre for Molecular Research, University of Nice Cote d'Azur, Nice 06204, France.  
meri.tulic@unice.fr

**AIMS AND SCOPE**

The primary aim of *World Journal of Dermatology* (WJD, *World J Dermatol*) is to provide scholars and readers from various fields of dermatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of dermatology and covering a wide range of topics including acneiform eruptions, acute generalized exanthematous pustulosis, angiolymphoid hyperplasia with eosinophilia, breast diseases, cutaneous fistula, dermatitis, dermatomyositis, erythema, exanthema, facial dermatoses, foot diseases, hair diseases, hand dermatoses, keratoacanthoma, keratosis, leg dermatoses, lipomatosis, lupus erythematosus, mastocytosis, morgellons disease, nail diseases *etc.*

**INDEXING/ABSTRACTING**

The WJD is now abstracted and indexed in Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

**Production Editor:** Xiang-Di Zhang; **Production Department Director:** Xu Guo; **Editorial Office Director:** Yun-Xiaoqiao Wu.

**NAME OF JOURNAL**

*World Journal of Dermatology*

**ISSN**

ISSN 2218-6190 (online)

**LAUNCH DATE**

June 2, 2012

**FREQUENCY**

Continuous Publication

**EDITORS-IN-CHIEF**

Zekayi Kutlubay

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2218-6190/editorialboard.htm>

**PUBLICATION DATE**

February 6, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Systematic review of hematidrosis: Time for clinicians to recognize this entity

Gilbert Sterling Octavius, Fellisa Meliani, Rivaldo Steven Heriyanto, Theo Audi Yanto

**Specialty type:** Dermatology

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): A

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Shalaby MN, Egypt; Sofou EI, Greece

**Received:** August 22, 2022

**Peer-review started:** August 22, 2022

**First decision:** September 27, 2022

**Revised:** October 12, 2022

**Accepted:** November 29, 2022

**Article in press:** November 29, 2022

**Published online:** February 6, 2023



**Gilbert Sterling Octavius, Fellisa Meliani, Rivaldo Steven Heriyanto, Theo Audi Yanto**, Department of Medicine, Universitas Pelita Harapan, Tangerang 15811, Indonesia

**Corresponding author:** Gilbert Sterling Octavius, MD, MM, Doctor, Researcher, Department of Medicine, Universitas Pelita Harapan, MH Thamrin Boulevard 1100, Klp. Dua, Kec. Klp. Dua, Kota Tangerang, Banten, Tangerang 15811, Indonesia. [sterlinggilbert613@hotmail.com](mailto:sterlinggilbert613@hotmail.com)

### Abstract

#### BACKGROUND

Hematidrosis is a sporadic disease, to a point where its existence is still denied up to date. It is also linked to stigmata, psychological roots, and religious beliefs, which has strengthened clinicians' disbelief in hematidrosis.

#### AIM

To conduct a thorough review to classify the likelihood of hematidrosis cases.

#### METHODS

We searched PubMed, Science Direct, Medline, and Google Scholar, as well as four different preprint databases, including Medrxiv, Research Square, SSRN, and Biorxiv. We included studies from 1996 onwards, with no limitation on language. Hematidrosis was classified as "unlikely", "likely", and "highly likely".

#### RESULTS

There are 74 articles with 106 hematidrosis cases. India ( $n = 40$ ) and China ( $n = 11$ ) report the most cases. Patients are mostly female (76.5%) with a median age of 13 years. The head region is the most common bleeding site ( $n = 168/254$ ). Headaches (26.9%) and abdominal pain (16.4%) are the most common prodromes. Beta-blockers (43%) and anxiolytic (23.2%) are the most commonly prescribed pharmacotherapy. Psychotherapy (37.5%) and counseling (32.5%) are the most utilized non-pharmacotherapy measures. Only 41.1% and 19.8% of all cases reach complete resolution and are highly likely to be hematidrosis, respectively.

#### CONCLUSION

Although hematidrosis is rare and the pathophysiology is still largely unknown, that does not mean hematidrosis does not exist. It is important to note that the most frequent trigger factors are either anxiety, fear, or excessive stress. Clinicians need to exclude other diagnoses and search for stressors to alleviate the bleeding.

**Key Words:** Hematidrosis; Bloody sweat; Stigmata; Blood

**Core Tip:** This is a systematic review on hematidrosis, an entity that is still mostly unknown, even to the experts. While this is not a guide to diagnose hematidrosis, this systematic review will help clinicians understand hematidrosis, the clinical pictures, the current available treatment, as well as the next steps in hematidrosis research.

**Citation:** Octavius GS, Meliani F, Heriyanto RS, Yanto TA. Systematic review of hematidrosis: Time for clinicians to recognize this entity. *World J Dermatol* 2023; 11(2): 7-29

**URL:** <https://www.wjgnet.com/2218-6190/full/v11/i2/7.htm>

**DOI:** <https://dx.doi.org/10.5314/wjd.v11.i2.7>

## INTRODUCTION

The first case of hematidrosis dates back to the 3rd century B.C., where the first two treatises by Aristotle contained sweat mixed with blood[1]. In the Bible, it is mentioned that these symptoms were written during Jesus Christ's sufferings in the garden of Gethsemane (Luke 22:44)[2,3]. Hematidrosis is an eccrine sweat disorder where sweat mixed with blood appear spontaneously without any visible trauma to the skin appendages. It is a diagnosis of exclusion[3]. Under International Classification of Disease 10 (ICD 10) in 2016, hematidrosis is given a diagnosis code of L74.8 under "other eccrine sweat disorders". However, this disease entity is still not widely accepted as a "true pathological disease" for various reasons[2,3].

Initially, the very existence of hematidrosis is questioned up to date. The mythical perception about hematidrosis creates a stigma where the bleeding is more related to pious beliefs than a genuine medical condition[4]. Another contributing factor is that a previous review on hematidrosis was done by Holoubek and Holoubek in 1996, encompassing 76 hematidrosis cases[5]. However, this review has been criticized for reporting the cases without laboratory confirmation[3,5]. Lastly, Favaloro & Lippi[6] mention some factors contributing to the plausible deniability of hematidrosis. Those explanations include the rarity of the disease, the possibility of Munchausen's syndrome (or Munchausen's by proxy), stress and its consequence on self-inflicted injuries, little scientific evidence that hematidrosis is an entity, and most reports are from older literature, in a foreign language, published in non-haematology journals, and there is a lack of publications in high-quality journals[6].

Several authors have attempted to conduct a literature review or systematic review on this topic[2,3,7]. However, the reviews do not present a comprehensive search, do not specify in which journals the articles were published, and more importantly, do not define what constitutes a hematidrosis. Therefore, we conduct a more thorough systematic review about hematidrosis, with a classification of the likelihood of hematidrosis cases in each journal. This review shall guide clinicians in identifying what hematidrosis is, how it presents, the necessary laboratory or radiology tests needed, as well as the treatments.

## MATERIALS AND METHODS

### Eligibility criteria

The Preferred Reporting Items for Systematic Review (PRISMA) 2020 guidance was used[8]. The protocol of this review was registered on the International Prospective Register of Systematic Reviews (PROSPERO) database with a registration number CRD42021289372.

The subjects studied were all patients diagnosed with hematidrosis without any age limitations. The diagnosis of hematidrosis is determined by the respective authors, and we classified the diagnosis as "unlikely", "likely", or "highly likely". A patient was considered to have "highly likely" hematidrosis if: (1) The bleeding episode was witnessed directly by medical professionals, (2) attempts were made to exclude other possible diagnoses, (3) skin biopsies and bleeding analysis were done and found to be normal or inconclusive, and (4) bleeding photos were included in the article; while a patient was determined to have "unlikely" hematidrosis if: (1) The bleeding episode was not witnessed by medical professionals or from family members, (2) minimal or no tests were done to exclude hematidrosis (only routine blood work was done), (3) other disorders possibly explain the bleeding, and (4) no bleeding photos were provided. Lastly, a hematidrosis case was classified as "likely" if it did not satisfy all the highly likely or unlikely criteria.

This review's inclusion criteria are any original articles published after 1996 without any restrictions in language. We restricted the year for literature search to ensure that studies that had been reviewed by Holoubek & Holoubek were not included[5]. We also included grey literature such as conference abstracts, thesis, or dissertations. The exclusion criteria of this study are reviews, opinion-based articles (letter to the editors or commentary), bleeding caused by other disorders, pure hemolacria, and animal studies. Citations from review studies were combed to ensure literature saturation. In order to guarantee that all available studies were included, we conducted citation and hand searches manually and *via* Research Gate.

### Search strategy and study selection

The literature search started on December 6, 2021 and ended on the same day. We searched four different academic databases, including PubMed, Science Direct, Medline, and Google Scholar, and four different preprint databases, including Medrxiv, Research Square, SSRN, and Biorxiv. The keywords used were "hematomidrosis", "hematohidrosis", and "bloody sweat". The Medical Subject Heading (MeSH) terms for each database are listed in Table 1. All records were imported into the Rayyan software, where duplicates were detected automatically and screened manually[9]. This software also allowed authors to collaborate in selecting the relevant studies. Two independent authors conducted the initial search (GSO and RSH), importing all the findings into Rayyan software. Another author (FM) cross-checked the initial searches. These three authors independently screened all available studies. Conflicts were resolved by discussion with the expert (TAY). In the case of studies with overlapping publications (abstracts later published into a full paper), we chose studies that provided more data.

### Data extraction and quality assessment

Data extraction was carried out independently by two authors (FM and RSH), then reviewed by another author (GSO) to ensure accuracy. We extracted relevant information such as study identification (author and year of publication), study characteristics (location and article type), and patients' characteristics (number of patients, age, sex, underlying conditions, characteristics of bleeding, laboratory and radiological examinations, therapies, and outcome).

The Joanna Briggs Institute (JBI) checklist for case reports and case series were used to assess their quality, respectively[10]. Three reviewers (GSO, FM, and RSH) independently assessed the scale, and any discrepancies were sorted with the expert (TAY) until a consensus was attained. If any missing data or further data were needed, corresponding authors were contacted *via* an inquiry email.

### Data synthesis

To incorporate all of the data in this review, pooled descriptive tests were employed. The mean and standard deviation of data reported in median and range (or interquartile range) were calculated[11-13]. The means and standard deviations were then combined into a single value using the Cochrane method[14].

## RESULTS

The initial search yielded 2955 articles, where 188 articles were immediately excluded as duplicates. After title and abstract screening, 2692 articles were excluded. Out of the 75 articles assessed for eligibility, three articles did not have full texts[15-17], three articles were review articles[18], one article mentioned other causes of the bleeding[19], and four articles were purely hemolacria[20-22]. These articles were then excluded, resulting in 68 articles[7,23-28]. We found another six articles through hand-searching and citation searching[29-34]. In total, there are 74 articles included in this review, with a total of 106 patients (Figure 1 and Table 1).

Most of the cases are heavily concentrated in Asia, particularly India ( $n = 40/106$ ) and China ( $n = 11/106$ ). Australia only reports one case of hematomidrosis, the least among other continents (Figure 2). Patients presented with the disease as early as an hour or as late as seven years before consulting to doctors. Anxiety, stress, and fear are the most common trigger for bleeding, while psychiatric disorders are the most common comorbidities in most patients. Notably, 36 cases (34%) do not have any obvious bleeding triggers. Besides bleeding from the skin, patients may also present with hematuria, gastrointestinal (GI) bleed (hematemesis, melena, hematochezia or rectal bleed), or epistaxis. Patients may bleed as frequently as more than 35 times a day or as rare as two times in two years. Most bleeding episodes subside in a few seconds to a few minutes, although in one case, the bleeding stops after 30 min (Table 2).

The majority of the patients are females (76.5%), with a median age of 13 (0.17-72) years old (Table 3). When patients experience prodrome(s) before the bleed, they mostly report headache (26.9%) or abdominal pain (16.4%). Most articles were published in 2019 ( $n = 15/74$ ), followed by 2021 ( $n = 10/74$ ) (Figure 3). When analyzed by the category of the journals, most hematomidrosis cases are published in dermatology journals (25.7%), followed by internal medicine journals (23.0%) and pediatric journals (20.3%).

**Table 1 Medical subject heading terms used in each database**

| Database        | Medical subject heading                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Number of studies found |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------|
| PubMed          | "hematidrosis"[All Fields] OR ("haematohidrosis"[All Fields] OR "hematohidrosis"[All Fields]) OR ("bloody"[All Fields] AND ("tearings"[All Fields] OR "tears"[MeSH Terms] OR "tears"[All Fields] OR "tearing"[All Fields] OR "lacrimal apparatus diseases"[MeSH Terms] OR "lacrimal"[All Fields] AND "apparatus"[All Fields] AND "diseases"[All Fields]) OR "lacrimal apparatus diseases"[All Fields] OR "lacerations"[MeSH Terms] OR "lacerations"[All Fields])) OR ("bloody"[All Fields] AND ("sweat"[MeSH Terms] OR "sweat"[All Fields] OR "sweating"[MeSH Terms] OR "sweating"[All Fields] OR "sweats"[All Fields] OR "sweatings"[All Fields])) OR "hemolacria"[All Fields] | 203                     |
| Medline         | ((hematidrosis) OR haematohidrosis) OR bloody sweat                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | 1057                    |
| Google Scholar  | allintitle: hematidrosis OR haematohidrosis OR bloody sweat                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | 55                      |
| Science Direct  | (Hematidrosis) OR (haematohidrosis) OR (bloody sweat)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 200                     |
| MedRxiv         | (Hematidrosis) OR (haematohidrosis) OR (bloody sweat)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 367                     |
| BioRxiv         | (Hematidrosis) OR (haematohidrosis) OR (bloody sweat)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 839                     |
| SSRN            | (Hematidrosis) OR (haematohidrosis) OR (bloody sweat)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 0                       |
| Research Square | (Hematidrosis) OR (haematohidrosis) OR (bloody sweat)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 134                     |

Out of all the body regions, the head region is the most commonly affected ( $n = 168/254$ ), especially around the ears or earlobes ( $n = 46$ ), forehead ( $n = 24$ ), and nose ( $n = 23$ ). The next most common site is in the upper limbs ( $n = 31/254$ ), with the arms being the most common site of bleeding in this region ( $n = 14$ ). Although most cases do not state the laterality of bleeding ( $n = 81$ ), more cases are bilateral ( $n = 15$ ), as compared to being unilateral bleeding ( $n = 10$ ) (Figure 4).

The bleeding episodes are mostly witnessed by healthcare professionals (65.1%). In the case of family members witnessing the bleeding episode, 50 cases (47.2%) are witnessed directly, and another 50 cases (47.2%) are not explicitly mentioned. The majority of the cases provide bleeding pictures (80.2%). Laboratory tests and radiologic or other tests are done in 94.3% and 38.7% cases, respectively, resulting in a normal result in 86.8% of the cases. An analysis of the bleed is done in 67.0% of cases, generally resulting in the discovery of peripheral blood components. There are many doctors from different specialities involved in the care of hematidrosis patients, with paediatricians (20.5%), psychiatrists (19.3%), dermatologists, and otolaryngologists (both at 16.6%) being the most common specialities consulted.

Skin biopsies, the "best reference standard", is only done in 28.3% of cases, with 63.3% of them being normal or non-specific. Among skin biopsies that have been done, abnormal findings include dilation of blood vessels with extravasation of blood[35-39], possible mild squamous hyperplasia and mild dermal fibrosis[40], papillary dermal edema[41,42] with dermal melanophages[41], bloody exudate from areas that do not contain sweat glands[43,44], a low number of gross cystic disease fluid protein 15-positive eccrine sweat glands[45], loosening areas caused by the separation of collagen fibres[43], and acanthosis of the epidermis with broadening of rete pegs along with hypergranulosis and marked hyperkeratosis [42].

The most common pharmacologic therapy used is beta-blockers (43.0%), with anxiolytic the second (23.2%). Among the non-pharmacologic modality, psychotherapy (37.5%) and counseling (32.5%) are the most common therapies prescribed. Most patients experienced complete resolution (41.1%) in 60 d (2-730 d), as well as an improvement of bleeding symptoms (41.1%) in 28 d (10-720 d).

Among all the 106 cases, the majority of cases belong to the "likely" group with 58 cases (54.7%), followed by "unlikely" with 27 cases (25.5%). Only 21 cases (19.8%) are considered "highly likely" to be hematidrosis.

## DISCUSSION

This systematic review contains 74 articles with 106 hematidrosis cases. We included more articles and cases compared to the other reviews and adjusting to their search timeline[2,3,7]. Hand-searching, citation searching, and including grey literature in our reviews certainly add more cases to our study. Although there has been some scepticism in including grey literature into a review, a proper systematic review shall be thorough during article searching, including finding grey literature[46]. Including the 76 cases from Holoubek and Holoubek's review[5] and our cases, there are currently 182 hematidrosis cases in the world.

Table 2 Characteristics of hematomatosis patients

| Ref.                                 | Patient No. | Time from onset to consultation | Triggers of bleeding                                                                         | Other kinds of bleeding | Comorbidities                                              | Bleeding frequency         | Bleeding duration   | Other Bleeding characteristics                   |
|--------------------------------------|-------------|---------------------------------|----------------------------------------------------------------------------------------------|-------------------------|------------------------------------------------------------|----------------------------|---------------------|--------------------------------------------------|
| Octavius <i>et al</i> [70], 2021     | #1          | One month                       | Not mentioned                                                                                | Not mentioned           | Not mentioned                                              | Three times per day        | Stopped immediately | Not mentioned                                    |
|                                      | #2          | One month                       | Not mentioned                                                                                | Not mentioned           | Not mentioned                                              | Not mentioned              | Not mentioned       | Not mentioned                                    |
| Bhattacharya <i>et al</i> [42], 2013 | #1          | One month                       | Not mentioned                                                                                | Not mentioned           | Not mentioned                                              | > 10 times per day         | Not mentioned       | Not mentioned                                    |
| Manonukul <i>et al</i> [43], 2008    | #1          | 1.5 yr ago                      | Not mentioned                                                                                | Not mentioned           | Not mentioned                                              | Recurrent episodes         | Not mentioned       | Not mentioned                                    |
| Murota <i>et al</i> [45], 2020       | #1          | A few months                    | Excess pressure exerted while playing on the metal bar                                       | Not mentioned           | Not mentioned                                              | Not mentioned              | Not mentioned       | It did not occur during sleeping                 |
| Das <i>et al</i> [89], 2020          | #1          | Two weeks                       | Head trauma                                                                                  | Not mentioned           | Not mentioned                                              | Not mentioned              | Not mentioned       | Not mentioned                                    |
|                                      | #2          | One hour                        | Anxiety symptoms                                                                             |                         | Anxiety                                                    |                            |                     |                                                  |
| Matsuoka <i>et al</i> [36], 2020     | #1          | Two years                       | Not mentioned                                                                                | Not mentioned           | Dissociative disorder and self-harming                     | Few times per week         | Not mentioned       | Bloody sweat-like fluid secretion                |
| Chabchoub <i>et al</i> [49], 2013    | #1          | 6 mau                           | Not mentioned                                                                                | Not mentioned           | Not mentioned                                              | Not mentioned              | Not mentioned       | Not mentioned                                    |
| Shafique <i>et al</i> [38], 2021     | #1          | 3 mo                            | Not mentioned                                                                                | Rectal bleed            | Significant psychosocial stress due to parental separation | 3-35 times per day         | Not mentioned       | Occurred during sleep and wakefulness            |
| Carvalho <i>et al</i> [35], 2008     | #1          | November 2004                   | Physical exertion, increase in room temperature                                              | Not mentioned           | Not mentioned                                              | Not mentioned              | Not mentioned       | Ceased spontaneously                             |
| Rossio <i>et al</i> [32], 2014       | #1          | Not mentioned                   | His family was experiencing an extremely stressful time, and his mother had just passed away | Not mentioned           | Not mentioned                                              | Not mentioned              | 1-2 min             | No predetermined time for recurrence             |
| Gayal <i>et al</i> [90], 2020        | #1          | Six months                      | Hearing voices and anxiety                                                                   | Not mentioned           | Schizophrenia                                              | 1-2 or 10-15 times per day | Not mentioned       | Depending on the intensity of psychotic symptoms |
| Dragan <i>et al</i> [91], 2017       | #1          | Not mentioned                   | Not mentioned                                                                                | Epistaxis, hemoptysis   | Tonsillectomy, benign hypertension                         | Not mentioned              | Not mentioned       | Not mentioned                                    |
| Alsermani <i>et al</i> [23], 2018    | #1          | Five months                     | Bullying in schools                                                                          | Not mentioned           | Celiac disease                                             | Twice weekly               | 1-2 min             | Self-limited, usually in the evening             |
| Jayaraman <i>et al</i> [92], 2017    | #1          | Seven days                      | Scolding from parents due to academic performance                                            | Not mentioned           | Mixed anxiety & depressive disorders, nocturnal enuresis   | 1-2 a day                  | 3-5 min             | Appears when punished or stressed                |
| Jibbawi <i>et al</i> [33], 2021      | #1          | Not mentioned                   | Emotional stress                                                                             | Not mentioned           | Not mentioned                                              | Not mentioned              | Not mentioned       | Appears during stress and self-limited           |

|                                     |    |               |                                                                                 |                                              |                                                                                                  |                                                                             |               |                                                                                                                |
|-------------------------------------|----|---------------|---------------------------------------------------------------------------------|----------------------------------------------|--------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|---------------|----------------------------------------------------------------------------------------------------------------|
| Arakkal <i>et al</i> [24], 2016     | #1 | 5 mau         | Extreme physical exertion                                                       | Not mentioned                                | Not mentioned                                                                                    | Not mentioned                                                               | 2-3 min       | Bleeding when waking up, extreme physical activities, self-limited                                             |
| Corrà <i>et al</i> [93], 2020       | #1 | Five years    | Stress, anxious situation, exercise                                             | Not mentioned                                | The peripheral visual deficit, non-hemorrhagic chronic gastritis, severe intestinal subocclusion | Five times daily                                                            | Few minutes   | Self-limited                                                                                                   |
| Mishra [72], 2009                   | #1 | Not mentioned | Minor trauma                                                                    | Hematuria, gastrointestinal bleed, epistaxis | Platelet function (PF3) dysfunction                                                              | Not mentioned                                                               | Not mentioned | Not mentioned                                                                                                  |
| Mutanabbi <i>et al</i> [29], 2021   | #1 | 4 mau         | Stress                                                                          | Hematemesis, hematuria, epistaxis            | Not mentioned                                                                                    | 1-2 a week                                                                  | 3-5 min       | Ceased spontaneously, precipitated by upcoming stress & family issues                                          |
| Soliman <i>et al</i> [61], 2019     | #1 | Six months    | Not mentioned                                                                   | Not mentioned                                | Hypotension, tachycardia, syncope                                                                | Not mentioned                                                               | 30 seconds    | Never at sleep and self-remitting                                                                              |
| Tshifularo [94], 2013               | #1 | Not mentioned | Severe mental stresses at work, study, and home                                 | Not mentioned                                | Not mentioned                                                                                    | 2-3 a week every 3-4 mo                                                     | 15-20 min     | 2-5 ml blood-stained non-clotting discharge                                                                    |
|                                     | #2 | Two years     | Severe stress                                                                   |                                              |                                                                                                  | Four times in 2 years (2009-2010)                                           | Not mentioned | Blood-stained sweaty discharge, not clotting and watery                                                        |
|                                     | #3 | Not mentioned | Stress                                                                          |                                              |                                                                                                  | Two times in 2 years (2010-2011)                                            | Not mentioned | Painless blood-stained sweat-like discharge                                                                    |
|                                     | #4 | Not mentioned | Severe stress at school and home                                                |                                              |                                                                                                  | 1st episode at July 2013                                                    | Not mentioned | Not mentioned                                                                                                  |
| Meyer <i>et al</i> [95], 2019       | #1 | Two days      | Stumbled and fell on an iron toy train (trauma)                                 | Not mentioned                                | Not mentioned                                                                                    | Almost daily                                                                | N/A           | Self-limiting, episodic. Bleeding occurred from apparently healthy and undamaged skin; no superficial injuries |
| Techasatian <i>et al</i> [81], 2016 | #1 | April 2014    | Not mentioned                                                                   | Not mentioned                                | Not mentioned                                                                                    | > 10 times                                                                  | 10-30 seconds | The time of the bleeding episode was unpredictable, not related to exertion or triggers                        |
| Yeşilova <i>et al</i> [96], 2016    | #1 | 2 d           | Move from cold to a warm environment, fallen out of bed (trauma) one day before | Not mentioned                                | Not mentioned                                                                                    | 20 times in 2 d                                                             | Not mentioned | Not mentioned                                                                                                  |
| Quentric <i>et al</i> [97], 2019    | #1 | Not mentioned | General anxiety related to school                                               | Epistaxis, nipple bleed                      | Not mentioned                                                                                    | 3-4 times a day                                                             | Not mentioned | Spontaneously, mostly at night                                                                                 |
| Mora <i>et al</i> [98], 2013        | #1 | 6 mo          | Stress                                                                          | Not mentioned                                | Not mentioned                                                                                    | Several times a day                                                         | Not mentioned | Spontaneous, self-limited, >30 episodes during hospitalization                                                 |
| Praveen <i>et al</i> [99], 2012     | #1 | 3 mo          | Stress (She had witnessed her elder sister being kidnapped nine months ago)     | Epistaxis                                    | Not mentioned                                                                                    | 2-3 times every week and had two episodes on the same day on four occasions | Not mentioned | Not mentioned                                                                                                  |
| Hansson <i>et al</i> [7], 2019      | #1 | Four weeks    | Not mentioned                                                                   | Not mentioned                                | Not mentioned                                                                                    | 20 episodes in 4 weeks                                                      | Not mentioned | Not mentioned                                                                                                  |

|                                      |    |                |                                                            |                                    |                                                                                              |                                   |                              |                                                                                                               |
|--------------------------------------|----|----------------|------------------------------------------------------------|------------------------------------|----------------------------------------------------------------------------------------------|-----------------------------------|------------------------------|---------------------------------------------------------------------------------------------------------------|
| Wang <i>et al</i> [44], 2014         | #1 | One month      | Not mentioned                                              | Not mentioned                      | Not mentioned                                                                                | 2-3 times a day                   | Not mentioned                | The bleed is described as a fresh, small amount of blood                                                      |
| Jerajani <i>et al</i> [41], 2009     | #1 | Two months     | Continuous mental stress                                   | Not mentioned                      | Depressive disorder                                                                          | Not mentioned                     | Not mentioned                | Not mentioned                                                                                                 |
| Lipsitt <i>et al</i> [100], 2018     | #1 | 9 mo           | Extreme emotional stress, excitement, or physical exertion | Not mentioned                      | Depression, anxiety, PTSD, non-epileptiform seizure                                          | Seven times a day                 | Not mentioned                | Progressive, episodic                                                                                         |
| Bhagwat <i>et al</i> [101], 2009     | #1 | Two years      | Trauma (witnessing a woman beheaded), Feeling anxious      | None                               | Severe depression                                                                            | 1-2 a day                         | 15-20 min                    | Episodic                                                                                                      |
| Sue Tin <i>et al</i> [40], 2015      | #1 | Three weeks    | No obvious trigger                                         | Not mentioned                      | Not mentioned                                                                                | 20 times a day                    | Not mentioned                | Spontaneous, worse during warmer months                                                                       |
| Pari[102], 2018                      | #1 | Five days      | Without any provoking factor                               | Not mentioned                      | Not mentioned                                                                                | 2-3 a day                         | 30 min                       | Not mentioned                                                                                                 |
| Patel <i>et al</i> [79], 2010        | #1 | Sincmaury 2007 | Not mentioned                                              | Not mentioned                      | Not mentioned                                                                                | Once-daily                        | 1 minute                     | Spontaneous, during school hours 9-11 am                                                                      |
| Biswas <i>et al</i> [76], 2013       | #1 | Two years      | Not mentioned                                              | Not mentioned                      | An intelligent quotient (IQ) of 60-70 with a loss of insight                                 | 1-2x daily                        | 10-15 min                    | Episodic, more frequent when waking up in the morning                                                         |
| Jafar <i>et al</i> [69], 2016        | #1 | Two weeks      | Not mentioned                                              | Epistaxis                          | Repeated faint attacks two years ago                                                         | Not mentioned                     | 10-20 min                    | It was spontaneous, unpredictable, and intermittent, with no specific patterns, and it stopped spontaneously. |
| Wang <i>et al</i> [103], 2010        | #1 | 3 yr           | Emotional excitement                                       | Not mentioned                      | Not mentioned                                                                                | > 20 times during hospitalization | Not mentioned                | It always occurred during the middle of the day, never when sleeping at night                                 |
| Thigarajan[83], 2019                 | #1 | Three days     | Stress fr'm father's demand on academics                   | Not mentioned                      | Primary enuresis, head tonsure, and trivial head trauma in a different location to the bleed | Not mentioned                     | Few seconds to a few minutes | Self-limited                                                                                                  |
| Khalid <i>et al</i> [82], 2013       | #1 | Five years     | Emotional stress                                           | None                               | None                                                                                         | Not mentioned                     | Not mentioned                | Not mentioned                                                                                                 |
| Maglie <i>et al</i> [104], 2017      | #1 | Three years    | Not mentioned                                              | Not mentioned                      | Major depressive disorder and panic disorder                                                 | Not mentioned                     | 1-5 min                      | Could occur in sleep or during physical activity & more intense bleeding occurs during emotional stress       |
| Uber <i>et al</i> [71], 2015         | #1 | 6 yr           | Not mentioned                                              | Not mentioned                      | Conversion, dissociative, and generalized anxiety disorder, high peak blood pressure         | Not mentioned                     | Few minutes                  | Spontaneous, sometimes occur during sleep, coincidental with high blood pressure peak                         |
| Salas-Alanis <i>et al</i> [37], 2021 | #1 | Not mentioned  | Acute emotional distress                                   | Not mentioned                      | Not mentioned                                                                                | Numerous                          | 3-25 min                     | From 6 years old, all following emotional distress                                                            |
| Talwar <i>et al</i> [80], 2021       | #1 | One year       | Separation from his aunt                                   | Epistaxis, hematuria, hematochezia | Adjustment disorder                                                                          | Once per week                     | A few minutes                | It can occur during sleep                                                                                     |
| Chowdhury <i>et al</i> [105], 2019   | #1 | Two years      | Migraine                                                   | Epistaxis, previous cesarean scar, | Migraine                                                                                     | Not mentioned                     | Not mentioned                | Associated with headache, her son also had hematomatosis                                                      |

|                                            |    |                      |                                          |                                  |                                                                                   |                   |               |                                                                                                                           |
|--------------------------------------------|----|----------------------|------------------------------------------|----------------------------------|-----------------------------------------------------------------------------------|-------------------|---------------|---------------------------------------------------------------------------------------------------------------------------|
|                                            |    |                      |                                          | menorrhagia                      |                                                                                   |                   |               |                                                                                                                           |
| Das <i>et al</i> [89], 2016                | #1 | 5 mo                 | Not mentioned                            | None                             | Otitis externa                                                                    | Not mentioned     | Not mentioned | More frequent on 06:00 A.M.                                                                                               |
| Hossain <i>et al</i> [77], 2018            | #1 | One and a half years | Not mentioned                            | Not mentioned                    | Not mentioned                                                                     | Not mentioned     | Not mentioned | Not mentioned                                                                                                             |
|                                            | #2 | Last two months      | Not mentioned                            | Not mentioned                    | Repeated episodes of stiffening of limbs with associated self-muttering           | Not mentioned     | Not mentioned | Not mentioned                                                                                                             |
|                                            | #3 | Six months           | Not mentioned                            | Not mentioned                    | Acute anxiety state, repeated episodes of unresponsiveness and conduct disorder   | Not mentioned     | Not mentioned | Not mentioned                                                                                                             |
| Thao <i>et al</i> [39], 2019               | #1 | One month            | Marriage                                 | Not mentioned                    | Continuous mental stress                                                          | Not mentioned     | Not mentioned | Not mentioned                                                                                                             |
| Récher[31], 2021                           | #1 | Not mentioned        | COVID-19 containment induced stress      | Not mentioned                    | Not mentioned                                                                     | Not mentioned     | Not mentioned | A vascular pathway remains after 24 hours of bleeding                                                                     |
| Omar[30], 2018                             | #1 | Not mentioned        | Stress                                   | Not mentioned                    | Not mentioned                                                                     | Occured in bouts  | Not mentioned | Not mentioned                                                                                                             |
| Tirthani <i>et al</i> [86], 2021           | #1 | Three months         | Loss of parents                          | Not mentioned                    | Recurrent panic attacks with tachycardia                                          | Not mentioned     | 2-3 min       | Not mentioned                                                                                                             |
| Ricci <i>et al</i> [106], 2018             | #1 | Three months         | Major stressful event from an earthquake | Not mentioned                    | Not mentioned                                                                     | Not mentioned     | Not mentioned | Not mentioned                                                                                                             |
| Bezner <i>et al</i> [25], 2013             | #1 | Two years            | Not mentioned                            | Hematuria                        | Heavy and irregular menstrual bleed                                               | Up to 20 daily    | Not mentioned | The bleeding episodes reportedly decreased during her menstrual period, which typically lasted for approximately ten days |
| Mahamat Abderraman <i>et al</i> [50], 2019 | #1 | Five years ago       | Not mentioned                            | History of hematuria, epistaxis, | Moderate anxiety-depressive disorder, acute urine retention, seizure, genu flexum | Not mentioned     | 10 min        | Not mentioned                                                                                                             |
| Rharrabti <i>et al</i> [51], 2016          | #1 | Not mentioned        | Family abuse, parental conflicts         | Vulvar bleeding                  | Conversion disorder. primary enuresis                                             | Not mentioned     | Not mentioned | Mostly at day, worsening during parental conflicts                                                                        |
| Morillo <i>et al</i> [52], 2019            | #1 | Seven months         | Not mentioned                            | Not mentioned                    | Bronchial asthma, previous history of abortion                                    | Not mentioned     | Not mentioned | Not mentioned                                                                                                             |
| Kumar <i>et al</i> [78], 2021              | #1 | Not mentioned        | Afraid, stress (during exams), anxiety   | Not mentioned                    | High-grade anxiety                                                                | Not mentioned     | Not mentioned | Spontaneous and recurrent (4-5 times in three days)                                                                       |
| Camargo <i>et al</i> [53], 2020            | #1 | Four years           | Stress and anxiety                       | Hematuria                        | Not mentioned                                                                     | 2-3 times a week  | Not mentioned | Sporadic occurrence                                                                                                       |
| Shen <i>et al</i> [107], 2015              | #1 | Six months           | Not mentioned                            | Not mentioned                    | Tonic seizures                                                                    | 1x every few days | 1-2 min       | Spontaneous, transient, self-limited                                                                                      |
| Shahriari <i>et al</i> [108], 2020         | #1 | Six months           | Going to the toilet or feeling lonely    | Not mentioned                    | Not mentioned                                                                     | Every two weeks   | Not mentioned | Not mentioned                                                                                                             |

|                                                 |    |                              |                                                             |                                      |                                                                                                                  |                                              |                 |               |
|-------------------------------------------------|----|------------------------------|-------------------------------------------------------------|--------------------------------------|------------------------------------------------------------------------------------------------------------------|----------------------------------------------|-----------------|---------------|
|                                                 | #2 | Not mentioned                | Not mentioned                                               | Epistaxis                            | Not mentioned                                                                                                    | Every three weeks                            | Not mentioned   | Not mentioned |
|                                                 | #3 | Not mentioned                | Fighting with her younger sister without any skin damage    | Not mentioned                        | A history of aggression                                                                                          | Not mentioned                                | Not mentioned   | Not mentioned |
| Badry <i>et al</i> [27], 2020                   | #1 | One month                    | Stress                                                      | Hematemesis                          | Depression                                                                                                       | 3-5 per week                                 | A few minutes   | Not mentioned |
| Hoover <i>et al</i> [54], 2019                  | #1 | Almost more than seven years | Illness (malaria) or distress                               | Hematemesis, hematochezia, hematuria | Previous history of treated malaria, human immunodeficiency virus(HIV) and hepatitis B due to blood transfusions | Not mentioned                                | Not mentioned   | Not mentioned |
|                                                 | #2 | Not mentioned                | Irritability and crying                                     | Not mentioned                        | Not mentioned                                                                                                    | Not mentioned                                | Not mentioned   | Not mentioned |
| Latorre Martinez <i>et al</i> [109], 2012       | #1 | One year                     | Stress                                                      | Not mentioned                        | Not mentioned                                                                                                    | Not mentioned                                | A few minutes   | Not mentioned |
| Kleymenova <i>et al</i> [47], 2020 <sup>1</sup> | #1 | Not mentioned                | Weather, stress                                             | Epistaxis                            | Menstruation problems                                                                                            | Not mentioned                                | Not mentioned   | Not mentioned |
| Rani <i>et al</i> [84], 2018                    | #1 | Not mentioned                | Autonomic dysfunction                                       | Hematuria                            | Autonomic dysfunction                                                                                            | Not mentioned                                | Not mentioned   | Not mentioned |
|                                                 | #2 |                              | Autonomic dysfunction                                       | Not mentioned                        | Spontaneous skin breaches resembling incision                                                                    |                                              |                 |               |
|                                                 | #3 |                              | Emotional & Psychological stressors                         | Not mentioned                        | Not mentioned                                                                                                    |                                              |                 |               |
|                                                 | #4 |                              | Emotional & Psychological stressors                         | Not mentioned                        | Not mentioned                                                                                                    |                                              |                 |               |
|                                                 | #5 |                              | Emotional & Psychological stressors                         | Not mentioned                        | Not mentioned                                                                                                    |                                              |                 |               |
|                                                 | #6 |                              | Emotional & Psychological stressors                         | Not mentioned                        | Not mentioned                                                                                                    |                                              |                 |               |
| Wang <i>et al</i> [110], 2021                   | #1 | One year                     | Not mentioned                                               | Not mentioned                        | Not mentioned                                                                                                    | Not mentioned                                | Not mentioned   | Not mentioned |
|                                                 | #2 | Four months                  | Not mentioned                                               | Hematemesis                          | Not mentioned                                                                                                    | Once every few days to dozens of times a day | Several minutes | Not mentioned |
|                                                 | #3 | Five months                  | Carsickness                                                 | Hematuria                            | Canker sores                                                                                                     | Up to 60 times a day                         | Several minutes | Not mentioned |
|                                                 | #4 | Nine months                  | Anxiety or nervousness, hemolacria induced by intense light | Hematemesis, hematuria               | Tonic-clonic seizures                                                                                            | Once every few days to dozens of times a day | Several minutes | Not mentioned |
|                                                 | #5 | Two months                   | Not mentioned                                               | Not mentioned                        | Not mentioned                                                                                                    | Once every few days to dozens of times a day | Several minutes | Not mentioned |
|                                                 | #6 | Four years                   | Military training, minor                                    | Not mentioned                        | Not mentioned                                                                                                    | Once every few days to                       | Several         | Not mentioned |

|                                      |    |                                                     |                                                                                                               |                                                             |                                                    |                                                 |                 |                                                                                |
|--------------------------------------|----|-----------------------------------------------------|---------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------|----------------------------------------------------|-------------------------------------------------|-----------------|--------------------------------------------------------------------------------|
|                                      |    |                                                     | emotional stress, or even during studying                                                                     |                                                             |                                                    | dozens of times a day                           | minutes         |                                                                                |
|                                      | #7 | Six months                                          | Not mentioned                                                                                                 | Not mentioned                                               | Not mentioned                                      | Once every few days to dozens of times a day    | Several minutes | Not mentioned                                                                  |
| Malik <i>et al</i> [34], 2019        | #1 | 6-8 yr                                              | Emotional stress                                                                                              | History of hemolacria                                       | Gestational hypertension (after lost to follow up) | Not mentioned                                   | Not mentioned   | Bleeding started 6-8 years ago                                                 |
|                                      | #2 | Not mentioned                                       | Periods of rage and stress                                                                                    | Epistaxis                                                   | Not mentioned                                      | Not mentioned                                   | Not mentioned   | Unprovoked, self-limited, and not triggered                                    |
| Shahgholi[75], 2018                  | #1 | Several months                                      | Extreme physical or emotional stress                                                                          | Hematemesis                                                 | Not mentioned                                      | Once or twice a day                             | 10-15 min       | Sometimes occurred spontaneously during sleep, subsided as soon as it is wiped |
|                                      | #2 | Not mentioned                                       | Upcoming exam, playing a computer game, watching fiction TV movies or when parents not satisfying his demands | Not mentioned                                               | Not mentioned                                      | Not mentioned                                   | Not mentioned   | Not mentioned                                                                  |
|                                      | #3 | Not mentioned                                       | Not mentioned                                                                                                 | Not mentioned                                               | Not mentioned                                      | Not mentioned                                   | 5 min           | Spontaneous, unpredictable, and intermittent                                   |
| Jagannathan <i>et al</i> [111], 2016 | #1 | Not mentioned                                       | Mental stress                                                                                                 | Epistaxis                                                   | Not mentioned                                      | Not mentioned                                   | Not mentioned   | Not mentioned                                                                  |
|                                      | #2 |                                                     |                                                                                                               | Not mentioned                                               |                                                    |                                                 |                 |                                                                                |
|                                      | #3 |                                                     |                                                                                                               | Not mentioned                                               |                                                    |                                                 |                 |                                                                                |
|                                      | #4 |                                                     |                                                                                                               | Not mentioned                                               |                                                    |                                                 |                 |                                                                                |
|                                      | #5 |                                                     |                                                                                                               | Not mentioned                                               |                                                    |                                                 |                 |                                                                                |
|                                      | #6 |                                                     |                                                                                                               | Not mentioned                                               |                                                    |                                                 |                 |                                                                                |
|                                      | #7 |                                                     |                                                                                                               | Not mentioned                                               |                                                    |                                                 |                 |                                                                                |
|                                      | #8 |                                                     |                                                                                                               | Hematemesis                                                 |                                                    |                                                 |                 |                                                                                |
| Agarwal <i>et al</i> [28], 2017      | #1 | One month                                           | Not mentioned                                                                                                 | Not mentioned                                               | Not mentioned                                      | Not mentioned                                   | 10 seconds      | Self-limited, during the day                                                   |
| Deshpande <i>et al</i> [112], 2014   | #1 | Since February 2013 (around more than one year ago) | Upcoming exams, fight with parents, and parents not satisfying his demands                                    | Hemoptysis, hematuria, epistaxis, hematochezia, hematemesis | Oppositional defiant disorder                      | Not mentioned                                   | Not mentioned   | Not mentioned                                                                  |
| Zhang <i>et al</i> [16], 2004        | #1 | Four months                                         | Not mentioned                                                                                                 | Hematemesis                                                 | A previous diagnosis of Henoch-schonlein purpura   | Several times a day or once every 3-5 d         | Lasts seconds   | The bleeding can occur during sleep                                            |
| Kumar <i>et al</i> [78], 2019        | #1 | One year ago                                        | Suspected fear and anxiety of his lecturer                                                                    | Not mentioned                                               | Not mentioned                                      | Three episodes per day to 20 episodes per month | Not mentioned   | Not mentioned                                                                  |
|                                      | #2 | Three months ago                                    | Fearful dreams                                                                                                | Not mentioned                                               | Not mentioned                                      | > 5 episodes per day                            | Not mentioned   | Episodes are not coinciding with dreams                                        |

|                                                                                                                                   |    |               |               |           |                                                  |               |                             |
|-----------------------------------------------------------------------------------------------------------------------------------|----|---------------|---------------|-----------|--------------------------------------------------|---------------|-----------------------------|
| but occur after the event at night without any triggering factor in the day times as observed by parents at home and the hospital |    |               |               |           |                                                  |               |                             |
| Karpukhina <i>et al</i> [48], 2020                                                                                                | #1 | One month ago | Not mentioned | Epistaxis | High anxiety, neurotic behaviour, and depression | Not mentioned | A few minutes Not mentioned |

<sup>1</sup>There are two cases presented in this literature. However, the second case is purely a hemolacria, so we excluded it.

Favaloro & Lippi's[6] arguments that most cases are outdated and presented in a foreign language are not valid anymore. Although some articles are written in Chinese[16,44] Russian[47,48], French[49-51], Spanish[52], and Portuguese[53], the majority of the articles are written in English language. Since the publications are mostly less than ten years old, we argue that the cases presented here are quite recent. Almost half of the articles were published in the years of 2019-2021. There is one study where the article was published in 2019, but on the website, the article is presented as if it was published in 2021. Therefore, we classify the study as published in 2019[54]. However, we believe that there is still a significant number of cases not published in a "high-quality journal" and thus may affect clinicians' perspectives on the eligibility of these cases[55]. Reflecting on the types of journal specialities, most cases are published in dermatology, pediatric, or internal medicine journals. Only 12.2% of the cases are published in a hematology journal. This point supports Favaloro & Lippi's argument about lacking publication in a high-profile hematology journal[6].

Most of the cases originate from China and India, the top two most populous countries globally. Although it seems convenient to link the connection between the majority of the cases belonging to the top two countries with the most population, the United States of America (USA) only reports three cases. Therefore, genetic, socio-cultural, race or environmental factors might play a role[56]. Lastly, publication bias may play a role in determining which cases get published[57]. Due to the rarity of the cases, many clinicians, even the most senior ones, may not encounter a hematomatidrosis case. Therefore, there is a lack of quality peer-reviewers who understand this case. This problem is confounded because many clinicians still doubt this disease entity[6].

The median age of patients with hematomatidrosis is 13 years, which is consistent with previous review findings of 9-15 years[3]. In this systematic review, 76.5% of the patients are females, while the most common trigger for hematomatidrosis is anxiety, fear, or mental illness. The connection between mental illnesses being more common in females[58-60] has been established and might explain the higher prevalence among these populations. In this study, the bleeding characteristics vary significantly. Some patients experienced bleeding episodes during sleep[38], while others never experienced bleeding during sleeping[45,61]. In some cases, patients usually bleed during the day or evening[23,28]. These patterns may reflect the physiological state of sympathetic nervous systems and hormonal fluctuations and their impacts on blood vessels and the pathophysiology of hematomatidrosis[62-64].

The head is the most common bleeding site, while headaches are the most common prodrome symptom. Spontaneous extracranial hemorrhagic phenomena may be caused by trigemino-autonomic reflex and sterile neurogenic inflammation[65]. Neural activation may also explain the prodromes of dizziness, nausea and vomiting, tingling, pain, photophobia, and phonophobia. Epistaxis is a frequent

**Table 3 Demographics of patients with hematomatidrosis**


| Variables                                               | N (%)        |
|---------------------------------------------------------|--------------|
| Sex ( <i>n</i> = 102)                                   |              |
| Male                                                    | 24 (23.5)    |
| Female                                                  | 78 (76.5)    |
| Age in years ( <i>n</i> = 94) (Median - range)          | 13 (0.17-72) |
| Prodromes <sup>1</sup>                                  |              |
| Headache                                                | 18 (26.9)    |
| Dizziness                                               | 3 (4.5)      |
| Abdominal pain                                          | 11 (16.4)    |
| Nausea                                                  | 4 (6.0)      |
| Vomiting                                                | 4 (6.0)      |
| Tingling                                                | 5 (7.5)      |
| Photophobia                                             | 2 (3.0)      |
| Phonophobia                                             | 2 (3.0)      |
| Pain                                                    | 3 (4.5)      |
| Soreness                                                | 4 (6.0)      |
| Asthenia                                                | 3 (4.5)      |
| Easily irritable                                        | 2 (3.0)      |
| Others                                                  | 6 (8.7)      |
| Presence of hemolacria                                  | 34 (32.1)    |
| Types of journals                                       |              |
| Case report                                             | 4 (5.4)      |
| Dermatology                                             | 19 (25.7)    |
| Otorhinolaryngology                                     | 4 (5.4)      |
| General or internal medicine                            | 17 (23.0)    |
| Hematology and oncology                                 | 9 (12.2)     |
| Pediatric (including pediatric hematology and oncology) | 15 (20.3)    |
| Pharmacy                                                | 1 (1.3)      |
| Preprint (Research Square)                              | 1 (1.3)      |
| Psychiatry                                              | 3 (4.1)      |
| Reproductive health                                     | 1 (1.3)      |
| Healthcare professionals witness the bleeding episodes  |              |
| Yes                                                     | 69 (65.1)    |
| No                                                      | 25 (23.6)    |
| Not explicitly mentioned                                | 12 (11.3)    |
| Family members witness the bleeding episodes            |              |
| Yes                                                     | 50 (47.2)    |
| No                                                      | 6 (5.6)      |
| Not explicitly mentioned                                | 50 (47.2)    |
| Provided bleeding pictures                              | 85 (80.2)    |
| Laboratory tests done                                   | 100 (94.3)   |
| Radiologic and/or other tests done                      | 41 (38.7)    |

|                                                                          |             |
|--------------------------------------------------------------------------|-------------|
| Normal results from all the diagnostic tests done                        | 92 (86.8)   |
| Skin biopsies done                                                       | 30 (28.3)   |
| Normal or non-specific                                                   | 19 (63.3)   |
| Analyzed the blood from the bleeding site                                | 71 (67.0)   |
| Doctors involved in the case <sup>1</sup>                                |             |
| Otolaryngologist                                                         | 25 (16.6)   |
| Ophthalmologist                                                          | 6 (4)       |
| Obstetrics and gynaecologist                                             | 3 (2)       |
| Psychiatrist                                                             | 29 (19.3)   |
| Primary care                                                             | 4 (2.7)     |
| Pediatrics                                                               | 31 (20.5)   |
| Internal medicine                                                        | 19 (12.8)   |
| Dermatologist                                                            | 25 (16.6)   |
| Vascular surgeon                                                         | 1 (0.7)     |
| Neurologist                                                              | 7 (4.8)     |
| Tried to exclude other diagnoses                                         | 90 (84.9)   |
| Pharmacotherapy <sup>2</sup>                                             |             |
| Beta-blockers                                                            | 65 (43.0)   |
| Anxiolytic                                                               | 35 (23.2)   |
| Antidepressants                                                          | 10 (6.6)    |
| Atropine patches                                                         | 2 (1.3)     |
| Drops of adrenaline with gauze                                           | 1 (0.7)     |
| Ascorbic acids                                                           | 6 (4.0)     |
| Antifibrinolytic agents                                                  | 2 (1.3)     |
| Antihistamine receptor blockers                                          | 6 (4.0)     |
| Hemostatics                                                              | 2 (1.3)     |
| Others                                                                   | 22 (14.6)   |
| Psychosocial therapy <sup>2</sup>                                        |             |
| Psychotherapy                                                            | 15 (37.5)   |
| Behavioural therapy                                                      | 6 (15)      |
| Relaxation technique                                                     | 6 (15)      |
| Counseling                                                               | 13 (32.5)   |
| Outcome ( <i>n</i> = 90)                                                 |             |
| Complete resolution                                                      | 37 (41.1)   |
| Time taken for complete resolution in days (median) ( <i>n</i> = 26)     | 60 (2-730)  |
| Improvement of symptoms                                                  | 37 (41.1)   |
| Time taken for improvement of symptoms in days (median) ( <i>n</i> = 19) | 28 (10-720) |
| Relapse                                                                  | 8 (8.9)     |
| Time taken for relapse in days (mean) ( <i>n</i> = 6)                    | 60 (53.4)   |
| Spontaneous resolution                                                   | 4 (4.4)     |
| The patient is still under follow up                                     | 1 (1.1)     |
| Loss to follow up                                                        | 3 (3.3)     |
| Possibilities of hematidrosis                                            |             |

|               |           |
|---------------|-----------|
| Highly likely | 21 (19.8) |
| Likely        | 58 (54.7) |
| Unlikely      | 27 (25.5) |

<sup>1</sup>One patient can be referred to multiple specialities or have multiple prodromes.

<sup>2</sup>One patient can receive more than one modality of treatment and more than one drug.




DOI: 10.5314/wjd.v11.i2.7 Copyright ©The Author(s) 2023.

Figure 1 PRISMA flowchart for selection of included studies.


accompanying bleeding manifestation and may result from activating the trigeminovascular system and subsequent vasodilation of Kiesselbach's plexus[66]. Abdominal pain is the second most common prodrome. While certainly, the cause of abdominal pain is a multitude of etiologies, a case was reported in which bleeding from distant sites preceded by abdominal migraine[67]. Activation of selected nerves and release of neuropeptide and neurotransmitters may facilitate haemorrhage remote from the site of pain[68].

The pathophysiology of bleeding in other areas is still unknown. The blood seeps into the sweat ducts due to vasculopathies in the dermal vasculature and enhanced sympathetic activation due to acute stress and anxiety, resulting in periglandular vascular constriction and subsequent expansion[69,70]. Multiple blood veins arranged in a net-like structure surrounding the sweat glands. These vessels are thought to contract and then widen to the point of rupture when they are under much stress. The blood now travels to the sweat glands, which are pushed to the surface and manifested as blood droplets mixed with sweat[70,71]. These pathways explain why patients who suffer from anxiety, high blood pressure, or elevated sympathetic nervous system tone suffer from hematiidrosis more frequently[71]. However, none of the biopsy results mention any vasculitis and this pathophysiology needs to be confirmed. The pathophysiology mentioned also does not explain the common co-occurrence of hemolacria, GI bleed, and hematuria. While there is a case of PF3 dysfunction[72], other patients do not have any hematological abnormalities after extensive workups. In the past, hematiidrosis was thought to be caused by infections such as *Chromobacterium prodigiosum* and *Micrococcus castellanii*[73]. One author



DOI: 10.5314/wjd.v11.i2.7 Copyright ©The Author(s) 2023.

Figure 2 Worldwide distribution of 106 hematidrosis cases from 74 articles.




DOI: 10.5314/wjd.v11.i2.7 Copyright ©The Author(s) 2023.

Figure 3 The number of cases is based on the year of publication. Note that no studies were found to be published between 2005-2007 and 2011. There is one study published in 2004 and two studies in 2010.

also tested for the presence of these bacteria in their case report, although they did not find any chromogenic bacteria[38]. Figure 5 presents a complete postulated pathophysiological pathway of hematidrosis.

Cutaneous manifestation of Munchausen syndrome or other factitious bleeding related to stigmata usually presents in areas within reach of the arms of the patients (legs, arms, anterior part of the body, and face). These episodes always occur alone and is never witnessed by family members or general practitioners[74]. Therefore, when 34.9% and 52.8% of cases are not witnessed or explicitly mentioned to be witnessed by healthcare professionals and family members, respectively, the diagnosis of hematidrosis becomes weaker. It is important to note that family members as a witness have a lower strength of proof compared to being directly witnessed by healthcare professionals such as doctors, nurses, or residents. Munchausen syndrome cases by proxy where the parents smear their blood to their children to mimic hematidrosis[18].



DOI: 10.5314/wjd.v11.i2.7 Copyright ©The Author(s) 2023.

**Figure 4 Characteristics of bleeding sites.** The head region is the most commonly affected ( $n = 168/254$ ), especially around the ears or earlobes ( $n = 46$ ), forehead ( $n = 24$ ), and nose ( $n = 23$ ). The next most common site is in the upper limbs ( $n = 31/254$ ), with the arms being the most common site of bleeding in this region ( $n = 14$ ). Although most cases do not state the laterality of bleeding ( $n = 81$ ), more cases are bilateral ( $n = 15$ ), as compared to being unilateral bleeding ( $n = 10$ ).

Analysis of the exudate is done to ensure that the reddish or pinkish liquid contains blood components[75]. Some authors add benzidine test[32,41,53,76-82], hemochromogen test[39], chloride test[83], or blood group matching[84] in order to ensure that the blood is likely from the patient. However, as stated above, even though the presence of blood is confirmed *via* observation of red blood cells and other peripheral blood components under the microscope, the blood does not necessarily belong to the patient, even when matched by their blood group. Therefore, we propose that a skin biopsy needs further studies to be the "gold standard" to exclude other skin pathologies before safely confirming that a patient has hematidrosis[43]. However, a biopsy needs to be taken immediately during or after bleeding episodes. When the biopsy is taken any later, the results may be insignificant or even normal[43]. This means that patients need to be admitted just for the biopsy procedures. Clinicians need to weigh the unnecessary risks of hospitalization and added fear and stress against the need for a confirmatory procedure that may add little value to the diagnosis.

The wide involvement of specialists indicates that clinicians are initially perplexed by this condition. Multicollaboration among specialists is needed to exclude other causes of bleeding before diagnosing a patient with hematidrosis. The majority of the cases presented in this review tried to exclude other diagnoses before jumping into diagnosing hematidrosis. The full differential diagnosis of hematidrosis is presented in Table 4.

The most effective treatment seems to be a combination of pharmacologic and non-pharmacologic therapies. Beta-blockers and anxiolytics are the most commonly used pharmacotherapies in concordance with the postulated pathophysiology. However, other therapeutic modalities such as tap-water iontophoresis[45], inosine[44], aluminium chloride hexahydrate[85], and oxybutynin[86] are also used with mixed results. The wide range of therapies indicates that the optimal route, timing, and dosage of therapy is still largely unknown. For example, while atropine transdermal patches may be an effective treatment, certain bleeding locations such as the eyes or vaginal bleeding will exclude this treatment modality[40,76]. When a patient has underlying anxiety or fear, a psychiatrist should be consulted to give appropriate psychosocial therapy[87]. Only 41.1% of patients are fully resolved from bleeding episodes. This emphasizes that the current treatment is still not effective in managing hematidrosis. Clinicians are encouraged to educate and communicate about the aetiology and nature of the disease, emphasizing that this is not a stigma or a "shame" to suffer from hematidrosis. Underlying psychiatric disorders need to be addressed as well.


Shahgholi *et al*[75] also attempt at establishing hematidrosis diagnostic criteria. We agree on health professionals witnessing the bleeding episode to exclude psychiatric causes. However, Shahgholi *et al* [75] mentioned blood analysis as one of their criteria. We opt for skin biopsies as they are the "gold

**Table 4 Differential diagnosis of hematidrosis**

|                                                                                                                                                                       |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Hematologic disorders (Bezner <i>et al</i> [25], 2013; Peretz <i>et al</i> [68], 2016)                                                                                |
| Acquired disorder of hemostatic mechanism                                                                                                                             |
| A hereditary disorder of hemostatic mechanism                                                                                                                         |
| Infection (Jirka, 1968; Peretz <i>et al</i> [68], 2016)                                                                                                               |
| <i>Chromobacterium prodigiosum</i>                                                                                                                                    |
| <i>Micrococcus castellani</i>                                                                                                                                         |
| Disorders of vascular integrity (Bezner <i>et al</i> [25], 2013; Peretz <i>et al</i> [68], 2016)                                                                      |
| Vasculitis                                                                                                                                                            |
| Amyloidosis                                                                                                                                                           |
| Connective tissue disorders                                                                                                                                           |
| Gynecologic disorders (Bezner <i>et al</i> [25], 2013)                                                                                                                |
| Ectopic endometriosis                                                                                                                                                 |
| Pathologic processes causing an acute increase in intracranial pressure                                                                                               |
| Cerebral venous thrombosis                                                                                                                                            |
| Subarachnoid hemorrhage                                                                                                                                               |
| Valsalva maneuver                                                                                                                                                     |
| Drugs (Girolami <i>et al</i> [74], 2014; Peretz <i>et al</i> [68], 2016)                                                                                              |
| Antiplatelets such as non-steroidal anti-inflammatory drugs                                                                                                           |
| Anticoagulants                                                                                                                                                        |
| Glucocorticoids                                                                                                                                                       |
| Antidepressants ( <i>e.g.</i> selective serotonin reuptake inhibitors)                                                                                                |
| Antiepileptics ( <i>e.g.</i> topiramate)                                                                                                                              |
| Antibiotics                                                                                                                                                           |
| Chemotherapeutic agents                                                                                                                                               |
| Intoxication with lysergic acid diethylamide (LSD)                                                                                                                    |
| Herbal supplement ( <i>e.g.</i> Gingko Biloba)                                                                                                                        |
| Trauma (Meyer <i>et al</i> [95], 2019; Peretz <i>et al</i> [68], 2016; Rani <i>et al</i> [84], 2018; Yeşilova <i>et al</i> [96], 2017; Zhang <i>et al</i> [16], 2004) |
| Sleep disorders (parasomnia) (Peretz <i>et al</i> [68], 2016)                                                                                                         |
| Psychiatric disorders (Bezner <i>et al</i> [25], 2013; Karpukhina <i>et al</i> [48], 2020; Peretz <i>et al</i> [68], 2016)                                            |
| Factitious disorder/factitious disorder by proxy                                                                                                                      |
| Psychogenic purpura                                                                                                                                                   |
| "Artistic bleeding"                                                                                                                                                   |
| Religious stigmata or culture-bound phenomenon (Bezner <i>et al</i> [25], 2013; Girolami <i>et al</i> [74], 2014)                                                     |
| Sweat gland disorders ( <i>e.g.</i> chromhidrosis) (Jerajani <i>et al</i> [41], 2009)                                                                                 |

standard" for diagnosis, however more confirmatory studies need to be done before skin biopsies can be considered the "gold standard" for hematidrosis. Blood analysis can be plagued by some issues mentioned above. The last criteria by Shahgholi *et al*[75] are summarised in our criteria as the authors tried to exclude other possible diagnoses. This is imperative as hematidrosis is currently a diagnosis of exclusion. However, to be categorized as "highly likely" as hematidrosis, we also encouraged authors to include the bleeding pictures in the publication. This criterion is usually unable to be met by abstract or poster publications[84]. Our criteria serve only as guidance to consider hematidrosis as a plausible diagnosis and not as a confirmatory guideline.

Our review has some limitations. We could not confirm the true nature of hematidrosis in each case due to limited presentations in some of the articles. Therefore, we judged each case with the best available information. None of the contacted authors replied. Secondly, we could only postulate some



**Figure 5 Postulated pathophysiology of hematidrosis.** There are several hypotheses in hematidrosis pathophysiology. The vasculitis in the dermal vessels, exacerbated by sympathetic activation related to extreme stress and anxiety, leads to periglandular vessel constriction, and subsequent expansion, allowing the blood content to seep into the sweat ducts. Another theory states that multiple blood vessels around the sweat glands are arranged in a net-like form. It is believed that under tremendous stress, the vessels contract. Once anxiety passes, the blood vessels dilate to the point of rupture. The blood, at this point, goes into the sweat glands, which push the blood to the surface and manifests as droplets of blood mixed with sweat. RBCs: Red blood cells.

associations, theories, and hypotheses between each finding without any causal confirmation. A cohort study with a standardized protocol will help determine the nature, progression, and treatment of hematidrosis. The exclusion of three articles without access to the full article is another limitation. However, those three articles are from China screened from their abstracts, they are unlikely to alter our review's findings significantly. Lastly, we do not include pure hemolacria cases in our review because pure hemolacria has its own sets of causes[88]. Therefore, including pure hemolacria in this review will cause heterogeneity.

Despite the limitations, our review is the most up-to-date with the most comprehensive literature search compared to other reviews[7,16,23-25,27-39,40-45,47,48,49-54,61,69-72,75-78,79-84,86,89-112]. Including articles in any language, combined with grey literature, citation searching, and hand searching, ensured that all available articles were included. We present clinical, diagnostic, and other socio-demographic findings that will help clinicians identify hematidrosis.

The future direction in diagnosing hematidrosis is currently evolving. Manonukul *et al*[43] performed electron microscopy and immunoperoxidase studies with normal results on both. Salas-Alanis *et al*[37] performed a genetic analysis and found that 91.5% of the variants are missense variants. The pathogenic variants were found in genes related to the extracellular matrix. These examinations will help us better understand the pathophysiology of hematidrosis.

## CONCLUSION

Hematidrosis is a rare disorder with an increasing number of cases. This disease is more common in Asia and young females with underlying anxiety, fear, stress, or depression. The head region is the most common bleeding site, and some patients may experience prodromal symptoms such as headache and abdominal pain. An important note to take is that hematidrosis might be accompanied by other kinds of

bleeding episodes such as hemolacria, GI bleeding, and epistaxis. The diagnosis is primarily on exclusion after ensuring that all the other diagnostic tests are normal. The most common treatment modality is a combination of pharmacologic (beta-blockers and anxiolytics) and non-pharmacologic (psychotherapy and counselling).

Although hematidrosis is rare and the pathophysiology is still largely unknown, that does not mean hematidrosis does not exist. It can bring severe panic towards parents or family members who care for these patients. Communication about the disease entity is imperative. A statement by Chambers perfectly summarized the rarity of hematidrosis "And as a rule, too, the more common the ailment, the more useful it is for you to hear about it"[113].

## ARTICLE HIGHLIGHTS

### **Research background**

Hematidrosis is a largely unknown entity, even to professional doctors.

### **Research motivation**

In order to bridge the underlying knowledge deficit about hematidrosis, an updated systematic review is conducted.

### **Research objectives**

We aim to conduct a systematic review of hematidrosis and identify the clinical characteristics, laboratory findings, as well as treatments given so far.

### **Research methods**

We conducted a systematic search on eight different databases with no restrictions on the timeline using the English language.

### **Research results**

There are 74 articles with 106 hematidrosis cases with India and China contributing the most cases. Patients are mostly females aged around 13 years. Bleeding occur most in the head region while beta-blockers and anxiolytics are the most frequent treatment given.

### **Research conclusions**

Hematidrosis exists, and it is up to clinical researchers to further dissect this entity so that physicians can give better treatment in the future.

### **Research perspectives**

More controlled skin biopsies and genetic studies with prospective follow-up or case-control studies may be needed to elucidate further and deepen our understanding of the pathophysiology and treatment of this disease.

## FOOTNOTES

**Author contributions:** Octavius GS and Yanto TA did the conception of this research; Data collections and selections are done by Octavius GS, Heriyanto RS, and Meliani F; Octavius GS, Heriyanto RS, and Meliani F drafted the article while Yanto TA did critical revision of the article; Final approval of the version to be published was granted by all authors.

**Conflict-of-interest statement:** All the authors declare that they have no conflict of interest.

**PRISMA 2009 Checklist statement:** The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Indonesia

**ORCID number:** Gilbert Sterling Octavius 0000-0002-6439-6265.

S-Editor: Liu JH

L-Editor: Ma JY

P-Editor: Liu JH

## REFERENCES

- 1 **Stolberg M.** Sweat: learned concepts and popular perceptions, 1500-1800. In: Horstmanshoff M, King H, Zittel C, editors. Blood, sweat and tears: The changing concepts of physiology from antiquity into early modern Europe. Leiden-Boston: Brill; 2012; 503-522
- 2 **Duffin J.** Sweating blood: history and review. *CMAJ* 2017; **189**: E1315-E1317 [PMID: [30978670](#) DOI: [10.1503/cmaj.170756](#)]
- 3 **Kluger N.** Hematomidrosis (bloody sweat): a review of the recent literature (1996-2016). *Acta Dermatovenereol Alp Pannonica Adriat* 2018; **27**: 85-90 [PMID: [29945265](#)]
- 4 **Bonomonte D, Vestita M, Filoni A, Giudice G, Angelini G.** Religious stigmata as malingering artifact: Report of a case and review of the literature. *Medicine (Baltimore)* 2016; **95**: e5354 [PMID: [27930512](#) DOI: [10.1097/MD.0000000000005354](#)]
- 5 **Holoubek JE, Holoubek AB.** Blood, sweat and fear. "A classification of hematomidrosis". *J Med* 1996; **27**: 115-133 [PMID: [8982961](#)]
- 6 **Favaloro EJ, Lippi G.** Commentary: Controversies in Thrombosis and Hemostasis Part 1-Hematomidrosis: "Blood, Sweat and Fears" or A "Pigment of Fertile Imaginations"? *Semin Thromb Hemost* 2018; **44**: 296-297 [PMID: [29220852](#) DOI: [10.1055/s-0037-1608906](#)]
- 7 **Hansson K, Johansson EK, Albåge M, Ballardini N.** Paediatric haematomidrosis: an overview of a rare but clinically distinct condition. *Acta Paediatr* 2019; **108**: 1023-1027 [PMID: [30849192](#) DOI: [10.1111/apa.14773](#)]
- 8 **Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D.** The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Syst Rev* 2021; **10**: 89 [PMID: [33781348](#) DOI: [10.1186/s13643-021-01626-4](#)]
- 9 **Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A.** Rayyan—a web and mobile app for systematic reviews. *Systematic Reviews* 2016; **5**: 210 [DOI: [10.1186/s13643-016-0384-4](#)]
- 10 **Moola S, Munn Z, Tufanaru C, Aromataris E, Sears K, Sfetcu R, Currie M, Lisy K, Qureshi R, Mattis P, Mu P.** Chapter 7: Systematic reviews of etiology and risk. In: Aromataris E, Munn Z (Editors). Joanna Briggs Institute Reviewer's Manual. *JBI Manual for Evidence Synthesis* 2022 [DOI: [10.46658/JBIMES-20-08](#)]
- 11 **Shi J, Luo D, Weng H, Zeng XT, Lin L, Chu H, Tong T.** Optimally estimating the sample standard deviation from the five-number summary. *Res Synth Methods* 2020; **11**: 641-654 [PMID: [32562361](#) DOI: [10.1002/jrsm.1429](#)]
- 12 **Luo D, Wan X, Liu J, Tong T.** Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Stat Methods Med Res* 2018; **27**: 1785-1805 [PMID: [27683581](#) DOI: [10.1177/0962280216669183](#)]
- 13 **Wan X, Wang W, Liu J, Tong T.** Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; **14**: 135 [PMID: [25524443](#) DOI: [10.1186/1471-2288-14-135](#)]
- 14 **Higgins JPT, Li T, Deeks JJ, (editors).** Chapter 6: Choosing effect measures and computing estimates of effect. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions*. 2nd Edition. Chichester (UK): John Wiley & Sons; 2019. p. 143-176 Available from: <https://training.cochrane.org/handbook/current/chapter-06>
- 15 **Chengrong X, Fan G, Yingling C.** [A case of hematomidrosis]. *Chinese Journal of Dermatology* 2010; **43**: 79-81
- 16 **Zhang FK, Zheng YL, Liu JH, Chen HS, Liu SH, Xu MQ, Nie N, Hao YS.** [Clinical and laboratory study of a case of hematomidrosis]. *Zhonghua Xue Ye Xue Za Zhi* 2004; **25**: 147-150 [PMID: [15182582](#)]
- 17 **Wei L, Guoqiang Z, Yixia W, Guang Z, Rusong M.** A case of hematomidrosis in children. *Chinese Journal of Dermatology* 2006; **39**: 113
- 18 **Saha B.** Intriguing hemorrhages: a review of last seven decades (1951-2021). *IJRMS* 2021; **9**: 2520-2534 [DOI: [10.18203/2320-6012.ijrms20213112](#)]
- 19 **Naqvi S, Asadullah Khan R, Rupareliya C, Hanif R, Ali Z, Farooq F.** Bleeding Diathesis or Fabrication: Münchhausen Syndrome. *Cureus* 2017; **9**: e1339 [PMID: [28951819](#) DOI: [10.7759/cureus.1339](#)]
- 20 **Wiese MF.** Bloody tears, and more! *Br J Ophthalmol* 2003; **87**: 1051 [PMID: [12881360](#) DOI: [10.1136/bjo.87.8.1051](#)]
- 21 **Wieser S.** Bloody tears. *Emerg Med J* 2012; **29**: 286 [PMID: [22205781](#) DOI: [10.1136/emered-2011-200955](#)]
- 22 **Barakova L, Reid J.** Haemolacria Associated with Severe Attacks of Migraine with Visual Aura and Hypertension. *Journal of Clinical Studies and Medical Case Reports* 2017; **102**: 104 [DOI: [10.24966/CSMC-8801/100102](#)]
- 23 **Alsermani M, Alzahrani H, El Fakih R.** Hematomidrosis: A Fascinating Phenomenon-Case Study and Overview of the Literature. *Semin Thromb Hemost* 2018; **44**: 293-295 [PMID: [29220853](#) DOI: [10.1055/s-0037-1608905](#)]
- 24 **Arakkal G, Poojari S, Netha G, Kumar B.** Hematomidrosis: A rare case of a female child who sweat blood. *IJPD* 2017; **18**: 327-329 [DOI: [10.4103/2319-7250.193031](#)]
- 25 **Bezner SK, Buchanan GR.** Bleeding from the eyes and through intact skin: physiologic, structural, spiritual, or faked? *Am J Hematol* 2013; **88**: 713-716 [PMID: [23674472](#) DOI: [10.1002/ajh.23485](#)]
- 26 **Varalakshmi B, Doshi VV, Sivalingam D, Nambi S.** The story of a girl with weeping blood: Childhood depression with a rare presentation. *Indian J Psychiatry* 2015; **57**: 88-90 [PMID: [25657464](#) DOI: [10.4103/0019-5545.148533](#)]
- 27 **Badry M, Elbadry M, Ragab A-R, Ahmed M.** Hematomidrosis: Reports and update of clinically mysterious phenomenon. *Indian Journal of Otolaryngology* 2020; **26**: 99-102 [DOI: [10.4103/indianjotol.INDIANJOTOL\\_135\\_19](#)]

- 28 Agarwal S, Rani S. Hematomidrosis: a hematology challenge. *HTIJ* 2017; **4**: 130-131 [DOI: [10.15406/htij.2017.04.00096](https://doi.org/10.15406/htij.2017.04.00096)]
- 29 Mutanabbi M, Ali A, Akhter S, Mosleh T, Morshed J. Haematomidrosis: A Rare Case Report of A 12 year Old Girl with Unexplained & Puzzling Spontaneous Multiple Site Bleeding. *Bangladesh Journal of Child Health* 2021; **45**: 45-47 [DOI: [10.3329/bjch.v45i1.55474](https://doi.org/10.3329/bjch.v45i1.55474)]
- 30 Omar S. Hematomidrosis, the Annoying Rare Condition, Case Report. *Experiments in Rhinology & Otolaryngology* 2018; **1**: 83-84 [DOI: [10.31031/ERO.2018.01.000524](https://doi.org/10.31031/ERO.2018.01.000524)]
- 31 Récher C. Hematomidrosis as a manifestation of COVID-19 containment-induced stress. *EJHaem* 2021; **2**: 25 [PMID: [35846085](https://pubmed.ncbi.nlm.nih.gov/35846085/) DOI: [10.1002/jha2.45](https://doi.org/10.1002/jha2.45)]
- 32 Rossio I, Gonçalves A. Haematomidrosis: The Rare Phenomenon of Sweating Blood. *Eur J Case Rep Intern Med* 2014; **1**-3 [DOI: [10.12890/2014\\_000144](https://doi.org/10.12890/2014_000144)]
- 33 Jibbawi A, Dirani F, Hassam I. An Extremely Rare Case Report of Hematomidrosis in Lebanon ISSN Print : 2709-3522 | ISSN Online : 2709-3530 Frequency : Bi-Monthly Language : English An Extremely Rare Case Report of Hematomidrosis in Lebanon I NTRODUCTION : 2021; 0-3 [DOI: [10.47310/Hjcmph.2021.v02i01.001](https://doi.org/10.47310/Hjcmph.2021.v02i01.001)]
- 34 Malik KZ, Abdullah A, Maka TA. Hematomidrosis-A Rare Knocker At Otolaryngologist's Door. *Pakistan Armed Forces Medical Journal* 2019; **69**: 730-732
- 35 Carvalho AC, Machado-Pinto J, Nogueira GC, Almeida LM, Nunes MB. Hematomidrosis: a case report and review of the literature. *Int J Dermatol* 2008; **47**: 1058-1059 [PMID: [18986356](https://pubmed.ncbi.nlm.nih.gov/18986356/) DOI: [10.1111/j.1365-4632.2008.03746.x](https://doi.org/10.1111/j.1365-4632.2008.03746.x)]
- 36 Matsuoka R, Tanaka M. Hematomidrosis in a Japanese girl: Treatment with propranolol and psychotherapy. *Pediatr Int* 2020; **62**: 1001-1002 [PMID: [32744359](https://pubmed.ncbi.nlm.nih.gov/32744359/) DOI: [10.1111/ped.14223](https://doi.org/10.1111/ped.14223)]
- 37 Salas-Alanis JC, Salas-Garza M, Goldust MM, Fajardo-Ramirez OR. Haematomidrosis and haemolacria in a young adult. *Clin Exp Dermatol* 2021; **46**: 394-396 [PMID: [33284990](https://pubmed.ncbi.nlm.nih.gov/33284990/) DOI: [10.1111/ced.14348](https://doi.org/10.1111/ced.14348)]
- 38 Shafique DA, Hickman AW, Thorne A, Elwood HR, Zlotoff BJ. Pediatric hematomidrosis - A case report and review of the literature and pathogenesis. *Pediatr Dermatol* 2021; **38**: 994-1003 [PMID: [34515356](https://pubmed.ncbi.nlm.nih.gov/34515356/) DOI: [10.1111/pde.14792](https://doi.org/10.1111/pde.14792)]
- 39 Thao NM, Lam THT, Doanh LH, Linh TT, Khang TH. A Vietnamese Case of Hematomidrosis. *Madridge J Case Rep Stud* 2019; **3**: 148-150 [DOI: [10.18689/mjcrs-1000139](https://doi.org/10.18689/mjcrs-1000139)]
- 40 Sue Tin A, Cohn A. Blood sweat and tears - tackling the stigma of stigmata. *J Paediatr Child Health* 2015; **51**: 1134-1136 [PMID: [26096319](https://pubmed.ncbi.nlm.nih.gov/26096319/) DOI: [10.1111/jpc.12936](https://doi.org/10.1111/jpc.12936)]
- 41 Jerajani HR, Jaju B, Phiske MM, Lade N. Hematomidrosis - a rare clinical phenomenon. *Indian J Dermatol* 2009; **54**: 290-292 [PMID: [20161867](https://pubmed.ncbi.nlm.nih.gov/20161867/) DOI: [10.4103/0019-5154.55645](https://doi.org/10.4103/0019-5154.55645)]
- 42 Bhattacharya S, Das MK, Sarkar S, De A. Hematomidrosis. *Indian Pediatr* 2013; **50**: 703-704 [PMID: [23942438](https://pubmed.ncbi.nlm.nih.gov/23942438/) DOI: [10.1007/s13312-013-0178-x](https://doi.org/10.1007/s13312-013-0178-x)]
- 43 Manonukul J, Wisuthsarewong W, Chantorn R, Vongirad A, Omeapinyan P. Hematomidrosis: a pathologic process or stigmata. A case report with comprehensive histopathologic and immunoperoxidase studies. *Am J Dermatopathol* 2008; **30**: 135-139 [PMID: [18360116](https://pubmed.ncbi.nlm.nih.gov/18360116/) DOI: [10.1097/DAD.0b013e318164cf4b](https://doi.org/10.1097/DAD.0b013e318164cf4b)]
- 44 Wang LJ, An CX, Li YM, Cao CQ. [A case report of childhood hematomidrosis]. *Zhongguo Dang Dai Er Ke Za Zhi* 2014; **16**: 214-215 [PMID: [24568922](https://pubmed.ncbi.nlm.nih.gov/24568922/)]
- 45 Murota H, Kotobuki Y, Yamaga K, Yoshioka Y. Female child with hematomidrosis of the palm: Case report and published work review. *J Dermatol* 2020; **47**: 166-168 [PMID: [31793058](https://pubmed.ncbi.nlm.nih.gov/31793058/) DOI: [10.1111/1346-8138.15179](https://doi.org/10.1111/1346-8138.15179)]
- 46 Saleh AA, Ratajeski MA, Bertolet M. Grey Literature Searching for Health Sciences Systematic Reviews: A Prospective Study of Time Spent and Resources Utilized. *Evid Based Libr Inf Pract* 2014; **9**: 28-50 [PMID: [25914722](https://pubmed.ncbi.nlm.nih.gov/25914722/) DOI: [10.18438/b8dw3k](https://doi.org/10.18438/b8dw3k)]
- 47 Kleymentova M, Khashchenko E, Kiseleva I, Uvarova E. Hemolacria and hematomidrosis: literature review and description of rare clinical cases in adolescent girls. Reproductivnoe zdorov'e detey i podrostkov [Pediatric and Adolescent Reproductive Health]. 2020; **16**: 102-16. [DOI: [10.33029/1816-2134-2020-16-4-102-116](https://doi.org/10.33029/1816-2134-2020-16-4-102-116)]
- 48 Karpukhina VN, Rudneva NS, Drachyov SN. Hematomidrosis or artistic dermatitis? *Medical Alphabet* 2020; **6**: 34-35 [DOI: [10.33667/2078-5631-2020-6-34-35](https://doi.org/10.33667/2078-5631-2020-6-34-35)]
- 49 Chabchoub RBA, Safi F, Trabelsi L, Maalej B, Gargouri L, Turki F, Amouri M, Halima NB, Turki, H, Mahfoud A. L'hématidrose: a propos d'un cas pédiatrique. *Archives de Pédiatrie*. 2013; **20**: 559-566 [DOI: [10.1016/j.arcped.2013.02.068](https://doi.org/10.1016/j.arcped.2013.02.068)]
- 50 Mahamat Abderraman G, Djidita Hagne Y, Mahamat Hissein A, Boudalia A, Sauvage J, Ibrahim H, Glenn R, Brahim N, Achta AF, Tara F, Charfadine S, Aboubacar A, Haddoum F. Haematomidrosis responding favorably to propranolol. *Press Medicale* 2019; **48**: 324-326
- 51 Rharrabti S, Khattala K, Belahsen M, Aalouane R. [Hematomidrosis and hemolacria associated with conversion disorder. About a pediatric case]. *Presse Med* 2016; **45**: 712-714 [PMID: [27371357](https://pubmed.ncbi.nlm.nih.gov/27371357/) DOI: [10.1016/j.lpm.2016.04.007](https://doi.org/10.1016/j.lpm.2016.04.007)]
- 52 Morillo Z, Ureña Correa W, Guzmán D, Ortiz Y. A propósito de un caso: "la niña que suda sangre: un raro caso de hematomidrosis". *Ciencia y Salud* 2019; **3**: 65-69 [DOI: [10.22206/cysa.2019.v3i2.pp65-69](https://doi.org/10.22206/cysa.2019.v3i2.pp65-69)]
- 53 Camargo KdCFd, Rezende LV, Galisteu BL, Muniz DN, Gatica JL, Ribeiro TS, Kinoshita JN. Hematomidrose: Doença Rara Relatada em Paciente Pediátrico. *Press Medicale* 2020; **1**-10
- 54 Hoover A, Fustino N, Sparks AO, Rokes C. Sweating Blood: A Case Series of 2 Siblings With Hematomidrosis. *J Pediatr Hematol Oncol* 2021; **43**: 70-72 [PMID: [31743319](https://pubmed.ncbi.nlm.nih.gov/31743319/) DOI: [10.1097/MPH.0000000000001661](https://doi.org/10.1097/MPH.0000000000001661)]
- 55 Kratz JE, Strasser C. Researcher perspectives on publication and peer review of data. *PLoS One* 2015; **10**: e0117619 [PMID: [25706992](https://pubmed.ncbi.nlm.nih.gov/25706992/) DOI: [10.1371/journal.pone.0117619](https://doi.org/10.1371/journal.pone.0117619)]
- 56 Bamshad M. Genetic influences on health: does race matter? *JAMA* 2005; **294**: 937-946 [PMID: [16118384](https://pubmed.ncbi.nlm.nih.gov/16118384/) DOI: [10.1001/jama.294.8.937](https://doi.org/10.1001/jama.294.8.937)]
- 57 Joobar R, Schmitz N, Annable L, Boksa P. Publication bias: what are the challenges and can they be overcome? *J Psychiatry Neurosci* 2012; **37**: 149-152 [PMID: [22515987](https://pubmed.ncbi.nlm.nih.gov/22515987/) DOI: [10.1503/jpn.120065](https://doi.org/10.1503/jpn.120065)]
- 58 Kuehner C. Why is depression more common among women than among men? *Lancet Psychiatry* 2017; **4**: 146-158 [PMID: [27856392](https://pubmed.ncbi.nlm.nih.gov/27856392/) DOI: [10.1016/S2215-0366\(16\)30263-2](https://doi.org/10.1016/S2215-0366(16)30263-2)]

- 59 **Li SH**, Graham BM. Why are women so vulnerable to anxiety, trauma-related and stress-related disorders? *Lancet Psychiatry* 2017; **4**: 73-82 [PMID: 27856395 DOI: 10.1016/S2215-0366(16)30358-3]
- 60 **Otten D**, Tibubos AN, Schomerus G, Brähler E, Binder H, Kruse J, Ladwig KH, Wild PS, Grabe HJ, Beutel ME. Similarities and Differences of Mental Health in Women and Men: A Systematic Review of Findings in Three Large German Cohorts. *Front Public Health* 2021; **9**: 553071 [PMID: 33614574 DOI: 10.3389/fpubh.2021.553071]
- 61 **Soliman M**, Mowafy K, Soliman R. Synchronous genital and facial hematomidrosis in adult female: first case report from Egypt. *EJMCR* 2019; **3**: 28-32 [DOI: 10.24911/ejmcr/173-1538850634]
- 62 **Dodt C**, Breckling U, Derad I, Fehm HL, Born J. Plasma epinephrine and norepinephrine concentrations of healthy humans associated with nighttime sleep and morning arousal. *Hypertension* 1997; **30**: 71-76 [PMID: 9231823 DOI: 10.1161/01.hyp.30.1.71]
- 63 **Muller JE**, Tofler GH, Verrier RL. Sympathetic activity as the cause of the morning increase in cardiac events. A likely culprit, but the evidence remains circumstantial. *Circulation* 1995; **91**: 2508-2509 [PMID: 7743610 DOI: 10.1161/01.cir.91.10.2508]
- 64 **Maus TL**, McLaren JW, Shepard JW Jr, Brubaker RF. The effects of sleep on circulating catecholamines and aqueous flow in human subjects. *Exp Eye Res* 1996; **62**: 351-358 [PMID: 8795453 DOI: 10.1006/exer.1996.0040]
- 65 **Drummond PD**. Mechanisms of autonomic disturbance in the face during and between attacks of cluster headache. *Cephalalgia* 2006; **26**: 633-641 [PMID: 16686902 DOI: 10.1111/j.1468-2982.2006.01106.x]
- 66 **Jarjour IT**, Jarjour LK. Migraine and recurrent epistaxis in children. *Pediatr Neurol* 2005; **33**: 94-97 [PMID: 16087052 DOI: 10.1016/j.pediatrneurol.2005.02.006]
- 67 **Kakisaka Y**, Uematsu M, Wang ZI, Haginoya K. Abdominal migraine reviewed from both central and peripheral aspects. *World J Exp Med* 2012; **2**: 75-77 [PMID: 24520537 DOI: 10.5493/wjem.v2.i4.7510.5493/wjem.v2.i4.75]
- 68 **Peretz AM**, Woldeamanuel YW, Rapoport AM, Cowan RP. Spontaneous extracranial hemorrhagic phenomena in primary headache disorders: A systematic review of published cases. *Cephalalgia* 2016; **36**: 1257-1267 [PMID: 26611681 DOI: 10.1177/0333102415618951]
- 69 **Jafar A**, Ahmad A. Child Who Presented with Facial Hematomidrosis Compared with Published Cases. *Case Rep Dermatol Med* 2016; **2016**: 5095781 [PMID: 27051537 DOI: 10.1155/2016/5095781]
- 70 **Octavius GS**, Koleta T, Garniasih D, Yanto TA. Hematomidrosis and Hemolacria: Report of Two Cases From Indonesia. *Iranian Journal of Blood and Cancer* 2021; **13**: 98-101
- 71 **Uber M**, Robl R, Abagge KT, Carvalho VO, Ehlke PP, Antoniuk SA, Werner B. Hematomidrosis: insights in the pathophysiology. *Int J Dermatol* 2015; **54**: e542-e543 [PMID: 26227471 DOI: 10.1111/ijd.12932]
- 72 **Mishra KL**. Bloody tears and hematomidrosis in a patient of PF3 dysfunction: a case report. *Cases J* 2009; **2**: 9029 [PMID: 20181216 DOI: 10.1186/1757-1626-0002-0000009029]
- 73 **Jirka M**. Hemingway's sweating waiter. *The Lancet* 1968; **291**: 1429 [DOI: 10.1016/S0140-6736(68)92387-8]
- 74 **Girolami A**, Bertozzi I, Tasinato V, Sambado L, Treleani M. Bleeding manifestations apparently unrelated to coagulation or other organic disorders: A tentative classification and diagnostic clues. *Hematology* 2014; **19**: 293-298 [PMID: 24164712 DOI: 10.1179/1607845413Y.0000000133]
- 75 **Shahgholi E**. A case series of hematomidrosis: A puzzling medical phenomenon. *Turk J Pediatr* 2018; **60**: 757-761 [PMID: 31365218 DOI: 10.24953/turkped.2018.06.022]
- 76 **Biswas S**, Surana T, De A, Nag F. A curious case of sweating blood. *Indian J Dermatol* 2013; **58**: 478-480 [PMID: 24249903 DOI: 10.4103/0019-5154.119964]
- 77 **Hossain A**, Nath S, Ghosh S, Hannan MHS, Pathak, Arnab, Mukherjee S, Dhar T. Hematomidrosis: Case Reports. *Indian Journal of Psychiatry: Medknow Publications & Media Pvt Ltd*; 2018; S124-S68. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5830862/>
- 78 **Kumar S**, Bhoi K, Yelme G. Hematomidrosis: A Rare and Mysterious Case. *Asian Journal of Case Reports in Medicine and Health* 2021; **5**: 10-13
- 79 **Patel RM**, Mahajan S. Hematomidrosis: A rare clinical entity. *Indian Dermatol Online J* 2010; **1**: 30-32 [PMID: 23130190 DOI: 10.4103/2229-5178.73256]
- 80 **Talwar M**, Chidambaram AC, Mekala S, Parameswaran N, Delhikumar CG. Haematomidrosis in a 12-year-old boy: blood, sweat and tears. *Paediatr Int Child Health* 2021; **41**: 300-302 [PMID: 34278983 DOI: 10.1080/20469047.2021.1951555]
- 81 **Techasatian L**, Warasawapati S, Jetsrisuparb C, Jetsrisuparb A. Hematomidrosis: a report with histological and biochemical documents. *Int J Dermatol* 2016; **55**: 916-918 [PMID: 26753524 DOI: 10.1111/ijd.13214]
- 82 **Khalid SR**, Maqbool S, Raza N, Mukhtar T, Ikram A, Qureshi S. Ghost spell or hematomidrosis. *J Coll Physicians Surg Pak* 2013; **23**: 293-294 [PMID: 23552544]
- 83 **Thigarajan KRT**. The vanishing ghostly bleeds - hematomidrosis. *University Journal of Medicine and Medical Specialities* 2019; **5**: 1-2
- 84 **Rani S**, Sandhya V, Varshini B, Vishnuvardhan, Agarwal S. Hematomidrosis-Case Series From A Tertiary Care Center. *Pediatric Hematology Oncology Journal* 2018 [DOI: 10.1016/j.phoj.2018.11.017]
- 85 **Patel NA**. Pediatric COVID-19: Systematic review of the literature. *Am J Otolaryngol* 2020; **41**: 102573 [PMID: 32531620 DOI: 10.1016/j.amjoto.2020.102573]
- 86 **Tirthani K**, Sardana K, Mathachan SR. Hematomidrosis of the mid-face and hands treated with oral oxybutynin. *Pediatr Dermatol* 2021; **38**: 962-963 [PMID: 34041778 DOI: 10.1111/pde.14621]
- 87 **Shamoon ZA**, Lappan S, Blow AJ. Managing Anxiety: A Therapist Common Factor. *Contemporary Family Therapy* 2017; **39**: 43-53 [DOI: 10.1007/s10591-016-9399-1]
- 88 **Ho VH**, Wilson MW, Linder JS, Fleming JC, Haik BG. Bloody tears of unknown cause: case series and review of the literature. *Ophthalmic Plast Reconstr Surg* 2004; **20**: 442-447 [PMID: 15599244 DOI: 10.1097/01.iop.0000143713.01616.cf]
- 89 **Das D**, Kumari P, Poddar A, Laha T. Bleeding to Life: A Case Series of Hematomidrosis and Hemolacria. *Indian J Pediatr*

- 2020; **87**: 84 [PMID: [31529380](#) DOI: [10.1007/s12098-019-03075-3](#)]
- 90 **Gayal TD**, Devi MG. A rare case of hematomidrosis in a patient with paranoid schizophrenia. *IJRMS* 2020; **8**: 3085-3087 [DOI: [10.18203/2320-6012.ijrms20203468](#)]
  - 91 **Dragan J**, Parrish RC. A 13-Year Old Female With Hemolacria and Hemoptysis. *IJCMPR* 2017; **3**: 2584-2586 [DOI: [10.24327/23956429.ijcmpr20170288](#)]
  - 92 **Jayaraman AR**, Kannan P, Jayanthini V. An Interesting Case Report of Hematomidrosis. *Indian J Psychol Med* 2017; **39**: 83-85 [PMID: [28250564](#) DOI: [10.4103/0253-7176.198953](#)]
  - 93 **Corrà A**, Quintarelli L, Caproni M. Bleeding from the oral cavity: a new case of hematomidrosis. *Int J Dermatol* 2020; **59**: e421-e422 [PMID: [32578211](#) DOI: [10.1111/ijd.15021](#)]
  - 94 **Tshifularo M**. Blood otorrhea: blood stained sweaty ear discharges: hematomidrosis; four case series (2001-2013). *Am J Otolaryngol* 2014; **35**: 271-273 [PMID: [24315735](#) DOI: [10.1016/j.amjoto.2013.09.006](#)]
  - 95 **Meyer J**, Spacil K, Stehr M, Hinrichs W, Haller S, Schäfer FM. Hematomidrosis After Head Injury - A Case Report. *Klin Padiatr* 2019; **231**: 326-327 [PMID: [31724140](#) DOI: [10.1055/a-1005-7328](#)]
  - 96 **Yeşilova Y**, Turan E, Aksoy M. Hematomidrosis on the forehead following trauma: a case report. *Int J Dermatol* 2017; **56**: 212-214 [PMID: [26945943](#) DOI: [10.1111/ijd.13274](#)]
  - 97 **Quentric PV**, Sauvêtre G. Unusual bleeding. *Eur J Intern Med* 2019; **68**: e1-e2 [PMID: [31358354](#) DOI: [10.1016/j.ejim.2019.07.013](#)]
  - 98 **Mora E**, Lucas J. Hematomidrosis: blood sweat. *Blood* 2013; **121**: 1493 [PMID: [23570065](#) DOI: [10.1182/blood-2012-09-450031](#)]
  - 99 **Praveen BK**, Vincent J. Hematomidrosis and hemolacria: a case report. *Indian J Pediatr* 2012; **79**: 109-111 [PMID: [21617906](#) DOI: [10.1007/s12098-011-0449-2](#)]
  - 100 **American Society of Pediatric Hematology/Oncology (ASPHO)**, David L. Lawrence Convention Center, Pittsburgh, PA, May 2-5, 2018. *Pediatr Blood Cancer* 2018; **65** Suppl 1: e27057 [PMID: [29603868](#) DOI: [10.1002/pbc.27057](#)]
  - 101 **Bhagwat PV**, Tophakhane RS, Rathod RM, Shashikumar BM, Naidu V. Hematomidrosis. *Indian J Dermatol Venereol Leprol* 2009; **75**: 317-318 [PMID: [19439898](#) DOI: [10.4103/0378-6323.51267](#)]
  - 102 **Pari T**. Hematomidrosis - A Rare Case. *Indian Dermatol Online J* 2019; **10**: 334-335 [PMID: [31149588](#) DOI: [10.4103/idoj.IDOJ\\_252\\_18](#)]
  - 103 **Wang Z**, Yu S, Su J, Cao L, Zhao X, Bai X, Zhan S, Wu T, Jin L, Zhou P, Ruan C. A case of hematomidrosis successfully treated with propranolol. *Am J Clin Dermatol* 2010; **11**: 440-443 [PMID: [20666570](#) DOI: [10.2165/11531690-000000000-00000](#)]
  - 104 **Maglie R**, Caproni M. A case of blood sweating: hematomidrosis syndrome. *CMAJ* 2017; **189**: E1314 [PMID: [29061857](#) DOI: [10.1503/cmaj.161298](#)]
  - 105 **Chowdhury A**, Islam M, Rahman H, Jannat M, Rahman M. Haematomidrosis with Headache-A Rare Phenomenon of Sweating Blood: A Case Report. *Bangladesh Med Res Counc Bull* 2019; **45**: 205-207 [DOI: [10.3329/bmrcb.v45i3.44654](#)]
  - 106 **Ricci F**, Oranges T, Novembre E, Della Bona ML, la Marca G, de Martino M, Filippi L. Haematomidrosis treated with propranolol: a case report. *Arch Dis Child* 2019; **104**: 171 [PMID: [29434019](#) DOI: [10.1136/archdischild-2017-314170](#)]
  - 107 **Shen H**, Wang Z, Wu T, Wang J, Ren C, Chen H, Yu Z, Don W. Haematomidrosis associated with epilepsy in a girl successfully treated with oxcarbazepine: case report. *J Int Med Res* 2015; **43**: 263-269 [PMID: [25673645](#) DOI: [10.1177/0300060514562488](#)]
  - 108 **Shahriari M**, Bazrafshan A, Karimi M, Shakibazad N. Case series of bloody sweating; a scary event for families. *Acta Haematologica Polonica* 2020; **51**: 258-260 [DOI: [10.2478/ahp-2020-0044](#)]
  - 109 **Latorre Martinez N**, Betlloch Mas I, Monteagudo Paz AF, Lucas Boronat J. Recurrent bleeding in an 18-year-old girl. *Arch Dermatol* 2012; **148**: 960-961 [PMID: [22911204](#) DOI: [10.1001/archdermatol.2012.1007](#)]
  - 110 **Wang Z**, Yin J, Yu Z, Han Y, Cao L, Kong D, Gu C, Bai X, Zhang X. Hematomidrosis: Pathophysiology and Therapeutic Strategy. Research Square 2021, Preprint, Available from: <https://doi.org/10.21203/rs.3.rs-256984/v1>
  - 111 **Jagannathan VV**, Rajasekar M, Mathivanan S, Mehta MP. Spontaneous ear nose throat bleed: hematomidrosis an unknown entity series of eight cases. *Int J of Otolaryngology and Head & Neck Surgery* 2016; **2**: 164-167
  - 112 **Deshpande M**, Indla V, Kumar V, Reddy IR. Child who presented with hematomidrosis (sweating blood) with oppositional defiant disorder. *Indian J Psychiatry* 2014; **56**: 289-291 [PMID: [25316941](#) DOI: [10.4103/0019-5545.140649](#)]
  - 113 **Chambers TK**. Clinical Lecture On A Case Of "Bloody Sweat." Given at St. Mary's Hospital, Feb. 9th, 1861. *Lancet* 1861; **77**: 207-209 [DOI: [10.1016/S0140-6736\(02\)41899-5](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

