

# World Journal of *Hematology*

*World J Hematol* 2014 February 6; 3(1): 1-17





# World Journal of Hematology

A peer-reviewed, online, open-access journal of hematology

## Editorial Board

2012-2016

The *World Journal of Hematology* Editorial Board consists of 123 members, representing a team of worldwide experts in hematology. They are from 29 countries, including Argentina (2), Austria (1), Belgium (1), Brazil (1), Canada (1), China (4), Croatian (1), Denmark (1), France (6), Germany (4), Greece (4), India (1), Iran (1), Ireland (1), Israel (2), Italy (11), Japan (9), Luxembourg (1), Mexico (1), Netherlands (7), Norway (2), Romania (1), Singapore (1), South Korea (2), Spain (6), Thailand (1), Turkey (5), United Kingdom (9), and United States (36).

### EDITORS-IN-CHIEF

Xiaoyan Jiang, *Vancouver*  
Thomas J Kipps, *San Diego*

### GUEST EDITORIAL BOARD MEMBERS

Hwei-Fang Tien, *Taipei*

### MEMBERS OF THE EDITORIAL BOARD



#### Argentina

Ricardo Forastiero, *Buenos Aires*  
Mirta A Schattner, *Buenos Aires*



#### Austria

Richard H Moriggl, *Vienna*



#### Belgium

Xavier Sagaert, *Leuven*



#### Brazil

Constantino José Fernandes Jr, *São Paulo*



#### China

Anskar YH Leung, *Hong Kong*  
Raymond HS Liang, *Hong Kong*  
Xiao-Yu Tian, *Hong Kong*



#### Croatian

Mariastefania Antica, *Zagreb*



#### Denmark

Erik Lerkevang Grove, *Aarhus*



#### France

Emmanuel Andres, *Strasbourg*  
Claude Bagnis, *Marseille*  
Bernard Binetruy, *Marseille*  
Cyril Fauriat, *Marseille*  
Florence Nguyen Khac, *Paris*  
Xavier Georges Thomas, *Pierre Benite*



#### Germany

Arnold Ganser, *Hannover*  
Dirk Matthias Hermann, *Essen*  
Rory R Koenen, *Munich*  
Zhixiong Li, *Hannover*



#### Greece

Anastasios G Kriebardis, *Athens*  
Marie-Christine Kyrtsolis, *Athens*  
Gerassimos A Pangalis, *Athens*  
Issidora S Papassideri, *Athens*



#### India

Gurudutta U Gangenahalli, *Delhi*



#### Iran

Shahram Teimourian, *Tehran*



#### Ireland

Eva Szegezdi, *Galway*



#### Israel

Jacob George, *Rehovot*  
Avichai Shimoni, *Tel-Hashomer*



#### Italy

Luca Arcaini, *Pavia*  
Vincenzo Casolaro, *Baronissi*  
Alessia Colosimo, *Teramo*  
Raimondo De Cristofaro, *Rome*  
Claudio Fozza, *Sassari*  
Edoardo G Giannini, *Genova*  
Giampiero La Rocca, *Palermo*  
Pier Paolo Piccaluga, *Bologna*  
Alessandro Poggi, *Genoa*  
Sergio M Siragusa, *Palermo*  
Elena Zocchi, *Genova*



#### Japan

Xian Wu Cheng, *Nagoya*  
Seiji Fukuda, *Izumo*  
Satoshi Hagiwara, *Yufu*  
Shinsaku Imashuku, *Takasago*  
Masanobu Kitagawa, *Tokyo*  
Tetsuya Nosaka, *Tsu*

Masao Seto, *Nagoya*  
Toshiki Shimizu, *Moriguchi*  
Masafumi Takahashi, *Shimotsuke*



#### **Luxembourg**

Jacques Zimmer, *Luxembourg*



#### **Mexico**

Agustin Avilés, *Mexico City*



#### **Netherlands**

Miranda Buitenhuis, *Rotterdam*  
Roland P Kuiper, *Nijmegen*  
Jan Jacques Michiels, *Erasmus*  
Gerry AF Nicolaes, *Maastricht*  
Pieter Sonneveld, *Rotterdam*  
Arnold Spek, *Amsterdam*  
Ruurd Torensma, *Nijmegen*



#### **Norway**

Brynjar Foss, *Stavanger*  
Mikhail Sovershaev, *Tromsø*



#### **Romania**

Adriana Georgescu, *Bucharest*



#### **Singapore**

Jerry Chan, *Singapore*



#### **South Korea**

Jung Weon Lee, *Seoul*  
Myung-Geun Shin, *Gwangju*



#### **Spain**

Matilde Canelles, *Granada*  
Slaven Erceg, *Seville*  
Pedro Cosme Redondo Liberal, *Cáceres*  
Julian Pardo, *Zaragoza*  
Josep-Maria Ribera, *Badalona*  
Juan M Zapata, *Madrid*



#### **Thailand**

Chirayu Udomsakdi Auewarakul, *Bangkok*



#### **Turkey**

Mutay Aslan, *Antalya*  
Murat Biteker, *Istanbul*  
Taner Demirer, *Ankara*  
Selami Kocak Toprak, *Ankara*  
Ertan Yetkin, *Mersin*



#### **United Kingdom**

Dominique Bonnet, *London*  
Olga Tura Ceide, *Edinburgh*  
Helen A Ireland, *London*  
Charles Henderson Lawrie, *Oxford*  
Drew Provan, *London*  
Dipak Purshottam Ramji, *Cardiff*  
Sabrina Tosi, *Uxbridge*

Shao-An Xue, *London*  
Jian guo Zhuang, *Liverpool*



#### **United States**

Ivo Abraham, *Tucson*  
Julia E Brittain, *Chapel Hill*  
Chung-Che Chang, *Houston*  
Edward Alan Copelan, *Cleveland*  
Zeev Estrov, *Houston*  
Steve Fiering, *Lebanon*  
Suzanne T Ildstad, *Louisville*  
Elias Jabbour, *Houston*  
Ming Jiang, *Nashville*  
Katsuhiko Kita, *Galveston*  
Robert G Lerner, *Valhalla*  
Shaoguang Li, *Worcester*  
Dazhi Liu, *Sacramento*  
Ming-Lin Liu, *Philadelphia*  
Surya Nauli, *Toledo*  
Steffan Nawrocki, *San Antonio*  
Xuyang Peng, *Nashville*  
Manuel L Penichet, *Los Angeles*  
Luis Francisco Porrata, *Rochester*  
Rehan Qayyum, *Baltimore*  
L Vijaya Mohan Rao, *San Diego*  
Guangwen Ren, *New Brunswick*  
Xiaoping Ren, *Cincinnati*  
Jatin J Shah, *Houston*  
Angus M Sinclair, *Thousand Oaks*  
Ali A Sovari, *Chicago*  
Christopher A Tormey, *New Haven*  
Olga Volpert, *Chicago*  
Zack Zhengyu Wang, *Scarborough*  
EllenLori Weisberg, *Boston*  
Wen-Shu Wu, *Scarborough*  
Yi Wu, *Newark*  
Feng-Chun Yang, *Indiana*  
Karina Yazdanbakhsh, *New York*  
Xin-Fu Zhou, *Adelaide*

**REVIEW**

- 1 A concise, practical guide to diagnostic assessment for mast cell activation disease

*Afrin LB, Molderings GJ*

## Contents

*World Journal of Hematology*  
Volume 3 Number 1 February 6, 2014

**APPENDIX** I-V Instructions to authors

**ABOUT COVER** Editorial Board Member of *World Journal of Hematology*, Claudio Fozza, MD, Research Fellow, Blood Diseases, Institute of Haematology, University of Sassari, Viale San Pietro 12, 07100 Sassari, Italy

**AIM AND SCOPE** *World Journal of Hematology* (*World J Hematol*, *WJH*, online ISSN 2218-6204, DOI: 10.5315) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

*WJH* covers topics concerning experimental, clinical, oncological and transplant hematology, transfusion science, hemostasis and thrombosis, evidence-based medicine, epidemiology and nursing. Priority publication will be given to articles concerning diagnosis and treatment of hematological diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJH*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

**INDEXING/ABSTRACTING** *World Journal of Hematology* is now indexed in Digital Object Identifier.

**FLYLEAF** I-II Editorial Board

## EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xin-Xin Che*  
Responsible Electronic Editor: *Huan-Liang Wu*  
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ling-Ling Wen*

**NAME OF JOURNAL**  
*World Journal of Hematology*

**ISSN**  
ISSN 2218-6204 (online)

**LAUNCH DATE**  
June 6, 2012

**FREQUENCY**  
Quarterly

**EDITOR-IN-CHIEF**  
**Xiaoyan Jiang, MD, PhD, Associate Professor,** Medical Genetics, University of British Columbia, Terry Fox Laboratory, British Columbia Cancer Agency, 675 West 10th Ave, Vancouver, BC, V5Z 1L3, Canada

**Thomas J Kipps, MD, PhD, Professor of Medicine,** University of California, San Diego, Moores Cancer Center, 3855 Health Sciences Drive, MC 0820, La Jolla, CA 92093-0820, United States

### EDITORIAL OFFICE

Jin-Lei Wang, Director  
Xiu-Xia Song, Vice Director  
*World Journal of Hematology*  
Room 903, Building D, Ocean International Center,  
No. 62 Dongsihuan Zhonglu, Chaoyang District,  
Beijing 100025, China  
Telephone: +86-10-85381891  
Fax: +86-10-85381893  
E-mail: [wjhematol@wjgnet.com](mailto:wjhematol@wjgnet.com)  
<http://www.wjgnet.com>

### PUBLISHER

Baishideng Publishing Group Co., Limited  
Flat C, 23/F, Lucky Plaza,  
315-321 Lockhart Road, Wan Chai,  
Hong Kong, China  
Fax: +852-6555-7188  
Telephone: +852-3177-9906  
E-mail: [bpgooffice@wjgnet.com](mailto:bpgooffice@wjgnet.com)  
<http://www.wjgnet.com>

### PUBLICATION DATE

February 6, 2014

### COPYRIGHT

© 2014 Baishideng Publishing Group Co., Limited. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

### SPECIAL STATEMENT

All articles published in this journal represent the viewpoints of the authors except where indicated otherwise.

### INSTRUCTIONS TO AUTHORS

Full instructions are available online at [http://www.wjgnet.com/2218-6204/g\\_info\\_20100722173604.htm](http://www.wjgnet.com/2218-6204/g_info_20100722173604.htm).

### ONLINE SUBMISSION

<http://www.wjgnet.com/esp/>

## A concise, practical guide to diagnostic assessment for mast cell activation disease

Lawrence B Afrin, Gerhard J Molderings

Lawrence B Afrin, Division of Hematology/Oncology, Medical University of South Carolina, Charleston, SC 29425-6350, United States

Gerhard J Molderings, Institute of Human Genetics, University Hospital of Bonn, D-53127 Bonn, Germany

**Author contributions:** Afrin LB wrote the initial draft; Afrin LB and Molderings GJ contributed equally to the final revision processes; and Molderings GJ subsequent edits.

**Correspondence to:** Lawrence B Afrin, MD, BSB103, MSC635, Division of Hematology/Oncology, Medical University of South Carolina, 173 Ashley Avenue, Charleston, SC 29425-6350, United States. [afrinl@musc.edu](mailto:afrinl@musc.edu)

Telephone: +1-843-7924271 Fax: +1-843-7920644

Received: September 5, 2013 Revised: October 6, 2013

Accepted: November 15, 2013

Published online: February 6, 2014

mastocytosis vs MCAS. Appropriate tissue examinations are needed to diagnose mastocytosis, while elevated levels of relatively specific mast cell mediators are sought to support diagnosis of MCAS. Whether assessing for mastocytosis or MCAS, testing is fraught with potential pitfalls which can easily yield false negatives leading to erroneous rejection of diagnostic consideration of MCAD in spite of a clinical history highly consistent with MCAD. Efforts at accurate diagnosis of MCAD are worthwhile, as many patients then respond well to appropriately directed therapeutic efforts.

© 2014 Baishideng Publishing Group Co., Limited. All rights reserved.

**Key words:** Mast cell activation disease; Mastocytosis; Mast cell activation syndrome; Mast cell mediators; Tryptase; KIT mutations

### Abstract

As recognition of mast cell (MC) involvement in a range of chronic inflammatory disorders has increased, diagnosticians' suspicions of MC activation disease (MCAD) in their chronically mysteriously inflamed patients have similarly increased. It is now understood that the various forms of systemic mastocytosis - diseases of inappropriate activation and proliferation of MCs seemingly driven by a small set of rare, usually constitutively activating mutations in assorted MC regulatory elements - comprise merely the tip of the MCAD iceberg, whereas the far larger and far more clinically heterogeneous (and thus more difficult to recognize) bulk of the iceberg consists of assorted forms of MC activation syndrome (MCAS) which manifest little to no abnormal MC proliferation and may originate from a far more heterogeneous set of MC mutations. It is reasonable to suspect MCAD when symptoms and signs of MC activation are present and no other diagnosis better accounting for the full range of findings is present. Initial laboratory assessment should include not only routine blood counts and serum chemistries but also a serum total tryptase level, which helps direct further evaluation for

**Core tip:** Mast cell activation disease (MCAD) is characterized by accumulation of genetically altered mast cells and/or abnormal release of these cells' mediators, affecting functions in potentially every organ system, often without causing abnormalities in routine laboratory or radiologic testing. Recent data suggest a high prevalence of MCAD. Thus, MCAD should be considered routinely in the differential diagnosis of patients with chronic multisystem polymorbidity of a generally inflammatory theme or patients in whom established diagnoses do not well account for the patient's presentation of symptoms consistent with mast cell mediator release. Mediator testing can be challenging but typically is manageable. Diagnostic efforts are worthwhile, as diagnosis often leads to effective therapy.

Afrin LB, Molderings GJ. A concise, practical guide to diagnostic assessment for mast cell activation disease. *World J Hematol* 2014; 3(1): 1-17 Available from: URL: <http://www.wjgnet.com/2218-6204/full/v3/i1/1.htm> DOI: <http://dx.doi.org/10.5315/wjh.v3.i1.1>

## INTRODUCTION

Of hematopoietic origin, mast cells (MCs) are found in all human tissues, especially at the environmental interfaces and perivascular/perineural sites<sup>[1]</sup>. They serve largely as sentinels of environmental change and bodily insults and respond by releasing large and variable assortments of molecular mediators which directly and indirectly influence behavior in other local and distant cells and tissues to respond to changes and insults so as to maintain, or restore, homeostasis.

MCs and the related rare cutaneous disease urticaria pigmentosa (UP) were first discovered in the latter half of the 19<sup>th</sup> century<sup>[2,3]</sup>, and the existence of seemingly even rarer systemic MC disease first became apparent in the middle of the 20<sup>th</sup> century<sup>[4,5]</sup>. For several decades it was thought that virtually all MC diseases were neoplastic, with symptoms resulting principally from accompanying inappropriate mediator release. Nearly a quarter century ago, the recurring somatic D816V mutation was discovered<sup>[6]</sup> in the dominant MC regulatory element, transmembrane tyrosine kinase receptor KIT<sup>[7]</sup>, in a high proportion of patients with systemic mastocytosis (SM)<sup>[8,9]</sup>. This constitutively activating mutation has since been found to drive many features of SM including aberrant aggregation and spindled morphology of MCs in SM, tryptase and histamine overexpression, and aberrant MC surface co-expression of CD25<sup>[10-12]</sup>. Several other KIT mutations, also quite rare, have since been found in other cases of SM, mostly in KIT's kinase domain 1 at or near codon 816 or in the juxtamembrane region of KIT<sup>[11]</sup>. Recently, evidence has emerged that the KIT<sup>D816V</sup> mutation induces the above-mentioned immunohistochemical and morphological changes in affected MCs but seems not to be solely responsible for the clinical symptoms of the MC disease<sup>[13,14]</sup>. This finding fits well with previous reports in healthy people bearing the KIT<sup>D816V</sup> mutation in their peripheral blood leukocytes<sup>[15]</sup>.

Around the same time as the discovery of KIT<sup>D816V</sup>, though, the notion was first advanced that some portion of MC disease might be due to inappropriate mediator release with little to no accompanying MC proliferation<sup>[16]</sup>. This theory appeared validated when the first recognized cases of what is now called MC activation syndrome (MCAS) were published recently<sup>[17,18]</sup>. Although not yet independently confirmed, there soon followed provocative, repeated findings from one group of a very wide array of (presumably mostly constitutively activating) mutations scattered across all domains of KIT in small cohorts of MCAS patients<sup>[19,20]</sup>. Many of these patients appeared to bear multiple mutations in MC KIT, with no apparent recurring patterns. Similar mutational complexity has been found, too, across the spectrum of chronic myeloproliferative neoplasms (MPNs) within which the MC disorders reside<sup>[21]</sup>, and in advanced mastocytosis itself<sup>[22]</sup>.

In recognition of the fact that all MC disease is, first and foremost, disease of inappropriate MC activation, Akin *et al.*<sup>[23]</sup> have proposed a new umbrella term of MC



**Figure 1 Spectrum of mast cell disease.** SM-AHNMD: Systemic mastocytosis with associated clonal hematologic non-mast-cell-lineage disorder.

activation disease (MCAD) to describe the full spectrum of MC disease (Figure 1). MCAS is estimated to be more prevalent<sup>[13,24]</sup>, but also more difficult to recognize, than other diseases traditionally ascribed to MC dysfunction. It also has been proposed that the various systemic MCAD variants and clinical phenotypes represent not distinct disease entities but rather varying presentations of a common generic root process of mast cell dysfunction<sup>[13]</sup>. The various forms of mastocytosis (principally cutaneous and systemic) may be the tip of a proverbial MCAD iceberg, fairly readily recognizable (in spite of their rarity) because of their defined unique immunohistochemical and relatively uniform clinical presentations, while the bulk of the iceberg - hidden below the waterline of easy clinical recognizability - may be a far larger, and far more heterogeneous, collection of variants of MCAS, some specifically named (*e.g.*, idiopathic anaphylaxis<sup>[18]</sup>, cryopyrin-associated periodic syndrome<sup>[25]</sup>) but most not<sup>[23,26]</sup>. It seems logical that marked mutational heterogeneity would drive the marked heterogeneity of aberrant mediator expression and clinical presentation which are observed in MCAD and which can easily confound the diagnostician. For example, although MCAD can readily impact any or all systems in the body, cases have been described of MCAS causing hematologic presentations as diverse as pure red cell aplasia<sup>[27]</sup>, erythrocytosis<sup>[28]</sup>, and agranulocytosis<sup>[29]</sup>.

Limited familial studies performed to date interestingly show high familial loading of MCAD despite a typical absence of germline mutations<sup>[24]</sup>; instead, most of the few families carefully studied show distinct sets of somatic mutations in the affected members with correspondingly varying clinical presentations. Clinical correlation suggests acquisition of the initial disease-causing mutations occurs relatively early in life, but additional mutations may develop in subclones over time. Efforts to reconcile the apparent dichotomy of high familial loading of MCAD with absence of apparent germline genetic mutations in affected families have led to thoughts that certain epigenetic alterations (GJM, unpublished data) may confer a state of genetic fragility which, upon interaction with varying biochemical milieus induced by assorted stressors (*e.g.*, infection), in turn

Form of disease		Features
Cutaneous mastocytosis		Usually found in childhood; prevalence in the general population of approximately 0.01% <sup>[116,117]</sup> ; various forms such as urticaria pigmentosa or telangiectasia macularis eruptiva perstans; diagnosed by skin biopsy
SM		Usually emerges in middle age; prevalence in the general population of approximately 0.0003% <sup>[13]</sup>
	Indolent SM	Survival equal to that of the general population <sup>[118]</sup>
	Smoldering SM	Proposed category
	SM with associated clonal hematologic non-MC-lineage disorder	<i>e.g.</i> , SM with lymphoma or leukemia; outcomes are best if the SM and the AHNMD are treated concurrently <sup>[38]</sup>
	Aggressive SM	SM with end-organ failure such as hepatic fibrosis, portal hypertension, malabsorption, or cytopenias
	Mast cell leukemia	Rare; the most lethal and most therapeutically challenging of all mast cell diseases
Mast cell sarcoma		Rare
Extracutaneous mastocytoma		Rare

**Figure 2 Diagnostic classification of mastocytosis<sup>[33]</sup>.** SM: Systemic mastocytosis; AHNMD: Associated clonal hematologic non-mast-cell-lineage disorder.

induces MCAD-causing mutations in MC regulatory elements.

Diagnosticians willing to pursue the possibility of MCAD in their patients whose chronic multisystem polymorbidity (often, but not always, of a generally inflammatory theme) has defied extensive prior diagnostic efforts (typically focused on subspecialty-oriented symptom subsets rather than broad multisystem considerations) currently face many challenges including controversies in precise diagnostic criteria (particularly for MCAS), uncertainty regarding the utility of various diagnostic tests, and difficulties properly managing specimens for such testing. Whether assessing for mastocytosis or MCAS, testing is fraught with potential pitfalls which can easily yield erroneous conclusions that MCAD is not present and thereby potentially add yet much more time and cost to the very long and expensive path most patients with MCAD (particularly MCAS) require to establish diagnosis, let alone effective therapy. We describe here in detail our thoughts regarding current proposals for diagnostic criteria for MCAD and our approach to diagnostic testing for MCAD, *i.e.*, both mastocytosis and MCAS.

## DIAGNOSTIC CRITERIA FOR MCAD

Classification systems and diagnostic criteria for mastocytosis were first proposed in the 1980s<sup>[30,31]</sup>. Since 2001 the World Health Organization (WHO) consensus criteria have guided classification and diagnosis globally<sup>[32]</sup>. The latest revision of these criteria, published in 2008<sup>[33]</sup>, divide mastocytosis broadly into cutaneous and systemic forms as well as the even rarer solid MC tumors of MC sarcoma and extracutaneous mastocytoma (Figure 2).

The WHO 2008 consensus diagnostic criteria for SM are shown in Figure 2. Given that it is now understood that MC KIT codon 816 mutations (a minor criterion) drive MC aggregation (a major criterion) and certain other minor criteria including MC spindling, tryptase

overexpression, and CD25 co-expression, reorganization of the WHO diagnostic criteria for SM may be in order.

Recognition of the need for an alternative, MCAS-like diagnosis came about in part because of the discovery of patients whose clinical presentation was highly consistent with SM but who did not satisfy the major SM diagnostic criterion and only satisfied one or two of the minor criteria. In particular, many patients significantly affected by aberrant MC activation previously have been denied diagnosis of, and treatment for, MCAD because of a serum tryptase level < 20 ng/mL. Presently there are two principal proposals for diagnostic criteria for MCAS (Figure 3). Potential problems with the Valent *et al.*<sup>[34]</sup> criteria<sup>[28]</sup> include (1) non-recognition of many of the symptoms that can result from MC activation; (2) lack of published validation that the described tryptase increase reliably distinguishes ordinary baseline fluctuation of tryptase from fluctuation induced by aberrant MC activation; (3) practical difficulties in providing/obtaining a specimen for serum total tryptase within 4 h of onset of an exacerbation of symptoms; and (4) practical difficulties in many patients in finding - in such a heterogeneous disease - MC-targeting agents that can effect at least partial response. Thus, for example, patients whose aberrant MC activation causes substantial muscle/joint/bone aching, constipation and abdominal pain, paresthesias, adenitis, and cognitive dysfunction but not the symptoms listed in these criteria would not qualify for the diagnosis. Similarly, a rise in serum tryptase (whose normal range typically is approximately 0.5 to 11 ng/mL) from 2.0 ng/mL to 4.2 ng/mL would be considered “20% + 2 ng/mL” evidence of aberrant MC activation, but a rise to 4.1 ng/mL would not. Furthermore, patients often are sufficiently disabled during a flare of symptoms that they cannot easily get to a medical center, and those who do travel to an urgent care facility or emergency department often encounter providers resistant to pursuing tests not needed for immediate care of the presenting symptoms.

WHO 2008 diagnostic criteria for systemic mastocytosis<sup>[33]</sup>

Major criterion:

Multifocal, dense aggregates of MCs (15 or more) in sections of bone marrow or other extracutaneous tissues and confirmed by tryptase immunohistochemistry or other special stains

Minor criteria:

- 1 Atypical or spindled appearance of at least 25% of the MCs in the diagnostic biopsy
- 2 Expression of CD2 and/or CD25 by MCs in marrow, blood, or extracutaneous organs
- 3 KIT codon 816 mutation in marrow, blood or extracutaneous organs
- 4 Persistent elevation of serum total tryptase > 20 ng/mL

Diagnosis of SM made by either (1) major criterion + any one or more minor criteria; or (2) any three minor criteria

Proposed diagnostic criteria for MCAS: Valent *et al*<sup>[34]</sup> criteria

- 1 Chronic/recurrent symptoms (flushing, pruritus, urticaria, angioedema, nasal congestion or pruritus, wheezing, throat swelling, headache, hypotension, and/or diarrhea) consistent with aberrant MC mediator release
- 2 Absence of any other known disorder that can better account for these symptoms
- 3 Increase in serum total tryptase of 20% above baseline plus 2 ng/mL during or within 4 h after a symptomatic period
- 4 Response of symptoms to histamine H<sub>1</sub> and/or H<sub>2</sub> receptor antagonists or other "MC-targeting" agents such as cromolyn

Proposed diagnostic criteria for MCAS: Molderings *et al*<sup>[26]</sup> criteria

Major criteria:

- 1 Multifocal MC aggregates as per WHO major criterion for SM
- 2 Clinical history consistent with chronic/recurrent aberrant MC mediator release (symptoms per Table 4 in<sup>[28]</sup>)

Minor criteria:

- 1 Abnormal MC morphology as per WHO SM minor criterion 1
- 2 CD2 and/or CD25 expression as per WHO SM minor criterion 2
- 3 Detection of known constitutively activating mutations in MCs in blood, marrow, or extracutaneous organs
- 4 Elevation in serum tryptase or chromogranin A, plasma heparin or histamine, urinary N-methylhistamine, and/or other MC-specific mediators such as (but not limited to) relevant leukotrienes (B<sub>4</sub>, C<sub>4</sub>, D<sub>4</sub>, E<sub>4</sub>) or PGD<sub>2</sub> or its metabolite 11-β-PGF<sub>2α</sub>

Diagnosis of MCAS made by either (1) both major criteria, or (2) the second major criterion plus any one of the minor criteria, or (3) any three minor criteria

**Figure 3 Diagnostic criteria for systemic mastocytosis and mast cell activation syndrome.** WHO: World Health Organization; MCs: Mast cells; MCAS: Mast cell activation syndrome; SM: Systemic mastocytosis PG: Prostaglandin.

Such resistance often persists even when the patient presents a prescription from a MC disease specialist specifically requesting MC mediator testing at times of such flares. Finally, in concert with the observed marked heterogeneity of the clinical presentation of MCAS (perhaps due to underlying marked mutational heterogeneity), some MCAS patients benefit little from the first few or several MC-targeting therapies tried, risking premature rejection of the diagnosis.

A potential problem with the Molderings *et al*<sup>[26]</sup> criteria for diagnosis of MCAS is the lack of mention of excluding other diagnoses (including mastocytosis) better accounting for the full range of findings in the patient, but such an exclusion would seem to be implicit. In practice, this scheme most often leads to diagnosis of MCAS by pairing of the second major criterion with the last of the minor criteria, and it appears to permit applying the diagnosis of MCAS - and therefore also pursuing therapy for MCAS - in a wider population of otherwise mysteriously chronically multi systemically ill patients, possibly increasing risk of misdiagnosis.

There have been no studies of the diagnostic accuracy or efficiency of these two diagnostic schemes. Also, at present there are no "gold standards" distinguishing

levels of MC mediators seen in normal MC activation/reaction from levels seen in aberrant MC activation. We believe such distinctions will be difficult to develop, therefore also making it difficult to perform comparative studies of accuracy of different diagnostic schemes. Routine use of MC whole exome/genome sequencing and mutational analysis in patients with clinical histories suspicious for MCAD may become the most efficient route to definitively diagnosing MCAD, but currently the specific polymerase chain reaction (PCR) assay for KIT<sup>D816V</sup> (found often in SM but seldom in MCAS) is the only mutational analysis routinely available in most clinics. At present, though, it is worth noting that in view of the demonstrated great mutational complexity across the MPN spectrum, it may be premature to declare any given MCAS patient's disease as "non-clonal" based on negative clonality evaluations less complete than whole KIT sequencing. The term "uncertain clonality" may be more accurate.

Given the very limited mutational and clonality testing for MCAD presently available in most clinics, laboratory evaluation for MCAS will continue, at least for the near future, to depend far more on demonstration of elevated levels of MC-specific mediators.

## DIAGNOSIS OF MCAD

### **Establishing suspicion**

Accurate diagnosis of any condition begins with inclusion of the correct diagnosis in the considered differential diagnosis. Despite its rarity, development of clinical suspicion for mastocytosis is somewhat easier than for MCAS given the often flagrant nature of the clinical presentation, either with classic appearance of UP or telangiectasia macularis eruptiva perstans (TMEP) in cutaneous mastocytosis or with classic appearance of recurrent unprovoked flushing and/or anaphylaxis in systemic mastocytosis.

Initial suspicion of MCAS is more challenging due to its heterogeneity and, often, lack of flagrant acute presentation. Diagnosticians need to be cognizant that MCAD can affect every system, usually affects multiple systems, and usually manifests symptoms in a subacute or chronic waxing/waning or episodic fashion, though episodes also can arise acutely, *i.e.*, the so-called “flares” or “attacks” or “spells” that many patients have of one symptom set or another. Many symptoms often are categorized as inflammatory in nature (*e.g.*, pain, diarrhea), though non-inflammatory symptoms (or at least symptoms not traditionally thought to be inflammatory in nature) are prevalent, too (*e.g.*, fatigue, paresthesias).

Another clue that MCAD may be present is simply the oddity or unexpectedness of certain clinical events or findings in the patient. Given the very large array of mediators normally produced and released by the MC<sup>[35]</sup> and therefore potentially abnormally expressed in MCAD, with each mediator causing a unique, and usually wide, range of direct and indirect, local and distant effects, the potential for the disease to manifest “odd”, “weird”, “strange”, “inexplicable” and “bizarre” clinical presentations is substantial. Such descriptors often are found throughout MCAD patients’ charts for years prior to diagnosis. Especially when seen in the context of pre-existing chronic multisystem inflammatory illness, the appearance in the patient of “unusual” new clinical phenomena should provoke consideration of MCAD. (Of course, the diagnostician is abetted in such recognition by remaining abreast of the steadily enlarging scope of the chronic inflammatory diseases. For example, diabetes mellitus type 2, obesity, and atherosclerotic vascular disease have been recognized as chronic inflammatory diseases only relatively recently.)

By the time MCAD is diagnosed, most patients have seen many providers, have undergone extensive testing (often frustratingly yielding normal or non-specifically/minimally abnormal results), and have been assigned many diagnoses (often with less than a full measure of confidence) which explain assorted subsets of findings but do not well account for the full range of findings, including chronicity often dating back decades. Treatments for these preliminary diagnoses sometimes help somewhat but often do not help at all, further perplexing the diagnostician. It is reasonable to suspect MCAD when at least several symptoms and signs of MC activation are

present (Table 1)<sup>[26,28,36,37]</sup> and no other diagnosis better accounting for the full range of findings is present. In particular, presence of a definitively diagnosed condition (*e.g.*, lymphoma) which does not well account for all symptoms and findings (*e.g.*, presyncope, erythrocytosis, *etc.*), or poor response of a definitively diagnosed condition to standard treatment for that condition, should raise suspicion for the presence of either a significant independent diagnosis or an alternative underlying ailment better accounting for the full range of symptoms and findings. In such cases, definitive diagnosis of a comorbid (and potentially even underlying) MCAD permits dual-directed therapy<sup>[38]</sup> which may lead to improved outcomes. Standardized detection of a MC mediator release syndrome can be achieved by a validated questionnaire (Figure 4). Although routine complete blood counts (CBCs) and metabolic panels often are confoundingly normal in MCAD patients, it is also the case that abnormalities in these and other common blood tests in these patients are commonly seen and are typically modest and stable. Frequent, though not necessarily constant, relative or absolute monocytosis, eosinophilia, basophilia, and/or reactive lymphocytosis, typically to just modest degrees, can be seen along with similar patterns of abnormality in routine chemistries such as aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (AP) (LBA and GJM, unpublished data), hyperbilirubinemia, and diet-independent hypercholesterolemia and hypertriglyceridemia<sup>[39]</sup>. Viewed in encounter-specific isolation, such abnormalities often raise no concern and garner no further attention, but trend analysis made easy by electronic medical record systems can quickly highlight the persistence of these abnormalities and permit expansion of the considered differential diagnosis to include entities such as MCAD which can account for these additional findings.

### **Mastocytosis or MCAS? Initial laboratory assessment**

It is now understood that the serum total tryptase level much more reflects the total body MC load than the total body MC activation state<sup>[40-42]</sup>. As such, serum tryptase is expected to be elevated in mastocytosis but usually is elevated little to none in MCAS. “Spillover” into blood circulation of tryptase and other MC mediators initially released in the tissues can be influenced by many factors. Hence, a normal value for a mediator in blood or other bodily fluid does not rule out its unregulated release in tissue. In fact, about 20% of SM patients present with serum tryptase < 20 ng/mL<sup>[37,43,44]</sup>, but such patients rarely harbor the more advanced forms of SM [aggressive SM (ASM) and MC leukemia (MCL)] which require therapeutic approaches different from those used for the more common indolent SM and MCAS. Marrow aspiration/biopsy, an uncomfortable procedure for some patients, is warranted in cases of suspected mastocytosis but seldom yields diagnostic findings in cases of MCAS not already manifesting significant hematologic abnormalities (LBA and GJM, unpublished data). Therefore, a serum tryptase persistently elevated to > 20 ng/mL in two or more spec-

**Table 1** Symptoms and findings in mast cell activation disease

System	Potential manifestations of MCAD (most are chronic, low-grade; some are persistent, but many are either episodic or fluctuant)
Constitutional	Fatigue, malaise, asthenia, "chronic fatigue syndrome", subjective and/or objective hyperthermia and/or hypothermia, "sense of feeling cold much of the time", sweats/diaphoresis (not always nocturnal), flushing, plethora or pallor, increased or decreased appetite, early satiety, weight gain or loss, pruritus, chemical and/or physical environmental sensitivities (often odd)
Dermatologic/integument	Rashes and lesions of many sorts (classic urticaria pigmentosa, "freckles", telangiectatic/angiomatic lesions, xerosis, warts, tags, folliculitis, ulcers, dyshidrotic eczema, diffusely migratory but sometimes focally persistent patchy macular erythema), pruritus (often diffusely migratory, sometimes aquagenic), flushing, angioedema, striae, dermatographism, hair thinning and alopecia, onychodystrophy (brittle nails, longitudinal ridges), poor healing
Ophthalmologic	Irritated eyes, increased or decreased lacrimation, suffusion, conjunctivitis, episodic difficulty focusing, lid tremor/tic (blepharospasm), solar sensitivity, infectious or sterile inflammation
Otologic/osmic	Infectious or sterile otitis externa and/or media, hearing loss or hyperacusis, tinnitus, otosclerosis, dysosmia, coryza, congestion
Oral/oropharyngeal	Pain or irritation (sometimes "burning"), leukoplakia, fibrosis, lichen planus, ulcers, sores, angioedema, dental decay, dysgeusia, throat tickle/discomfort/irritation/pain, post-nasal drip
Lymphatic	Adenopathy, usually sub-pathologic and often waxing/waning in size, sometimes asymptomatic but not uncommonly tender, sometimes focal, sometimes migratory, pathology usually shows reactive lymphocytosis or sometimes an atypical non-specific lymphoproliferative disorder; left upper quadrant discomfort (likely from release of mediators from splenic mast cells with or without detectable splenomegaly)
Pulmonary	Rhinitis, sinusitis, pharyngitis, laryngitis, bronchitis, pneumonitis (often confused with infectious pneumonia), cough, dyspnea (often low-grade, inconstant, "I just can't catch a deep breath" despite normal pulmonary function tests), wheezing, obstructive sleep apnea, pulmonary hypertension
Cardiovascular	Presyncope (lightheadedness, weakness, dizziness, vertigo) and/or syncope (patients may have been diagnosed with postural orthostatic tachycardia syndrome or neurocardiogenic syncope), hypertension and/or hypotension, palpitations, dysrhythmias, chest discomfort or pain (usually non-anginal in character), coronary and peripheral arterial atherosclerosis/spasm/infarction, idiopathic acute or chronic heart failure (e.g., takotsubo), aneurysms, hemorrhoids, varicosities, aberrant angiogenesis (hemangiomas, arteriovenous malformations, telangiectasias), migratory edema (often non-dependent and with normal cardiac and renal function)
Gastrointestinal	Aerophagia, angioedema in any segment of the luminal tract, dysphagia (often proximal, possibly due to pharyngeal angioedema), bloating/gas, pain/inflammation (often migratory) in one or more segments of the luminal tract (from esophagitis to proctitis) and/or one or more solid organs (e.g., hepatitis, pancreatitis), queasiness, nausea, vomiting (sometimes "cyclical"), diarrhea and/or constipation (often alternating), malabsorption (more often selective micronutrient malabsorption than general protein-calorie malabsorption), ascites either from portal hypertension and/or peritoneal serositis; gastroesophageal reflux disease (often "treatment-refractory") and inflammatory/irritable bowel syndrome are common pre-existing diagnoses
Genitourinary	Inflammation (often migratory) in one or more segments of the luminal tracts (ureteritis, cystitis, urethritis, vaginitis, vestibulitis) and/or one or more solid organs (e.g., nephritis, prostatitis), chronic kidney disease, endometriosis, chronic low back pain or flank pain or abdominal pain, hydronephrosis (likely from ureteral angioedema), infertility, erectile dysfunction, decreased libido; in the appropriate setting of multisystem morbidity, miscarriages should prompt consideration of antiphospholipid antibody syndrome potentially due to MCAD
Musculoskeletal	Clinical myositis, often diffusely migratory (fibromyalgia is a common pre-existing diagnosis), subclinical myositis (i.e., asymptomatic elevated creatine kinase not otherwise explained), arthritis (typically migratory), joint laxity/hypermobility (patients may have been diagnosed with Ehlers-Danlos Syndrome Type III), osteoporosis/osteopenia, osteosclerosis, sometimes mixed osteoporosis/osteopenia/osteosclerosis; MCAD-driven musculoskeletal pain not uncommonly is poorly responsive to non-steroidal anti-inflammatory drugs and narcotics
Neurologic	Headache (esp. migraine), presyncope and/or syncope, peripheral (usually distal) sensory and/or motor neuropathies including paresthesias, tics, tremors (typically resting), chronic inflammatory demyelinating polyneuropathy, seizure disorders (can be "treatment-refractory"), pseudoseizures, dysautonomia
Psychiatric	Mood disturbances (e.g., anger, depression), bipolar affective disorder, attention deficit-hyperactivity disorder, post-traumatic stress disorder, anxiety and panic, psychoses, memory difficulties, word-finding difficulties, other cognitive dysfunction, wide variety of sleep disruptions
Endocrinologic/metabolic	Abnormal electrolytes (including magnesium) and liver function tests, delayed puberty, dysmenorrhea, endometriosis, osteosclerosis and/or osteoporosis, hypothyroidism, hyperthyroidism, dyslipidemia, hyperferritinemia, selective vitamin and/or other micronutrient deficiencies, weight change, possibly diabetes mellitus
Hematologic/coagulopathic	Polycythemia or anemia (may be macrocytic, normocytic, or microcytic), leukocytosis or leukopenia, chronic (usually mild) monocytosis or eosinophilia or basophilia, thrombocytosis or thrombocytopenia, arterial and/or venous thromboembolic disease, "easy" bruising/bleeding; in mast cell activation syndrome the marrow usually does not show increased (or even flow-cytometrically aberrant) mast cells; marrow histology often read as normal or as unspecified myelodysplastic/myeloproliferative syndrome; standard cytogenetic studies are almost always normal or show culture failure
Immunologic	Type I, II, III and IV hypersensitivity reactions, increased risk for malignancy, autoimmunity, impaired healing, increased susceptibility to infection, elevated or decreased levels of one or more isotypes of immunoglobulin; modest monoclonal gammopathy of undetermined significance not uncommon

MCAD: Mast cell activation disease.

imens facilitates an initial decision whether to evaluate further for SM with biopsies as appropriate including at least marrow, and possibly also skin, upper and lower gastrointestinal (GI) tract mucosa, and potentially other tissues, especially those at the environmental interfaces. An

initial serum tryptase < 20 ng/mL makes SM (especially ASM and MCL) much less likely, and since the prognosis of, and therapeutic approach toward, indolent SM and MCAS are presently indistinguishable, there appear to be no adverse consequences if the clinician initially misdiag-

*Clinical signs*

- The patient complains about recurring or continuing burning and/or crampy abdominal pain of unknown cause and/or recurring or continuing diarrhea of unknown cause and/or frequently intense meteorism/gassiness (independent of the composition of diet) and/or about episodically occurring nausea. ☐ 1
- The symptoms respond to treatment with H1-antihistamines. ☐ 1
- The progression of the symptoms occurred in episodes with symptom-free periods becoming shorter. ☐ 1
- The patient complains about episodically occurring burning and/or choking chest pain attacks, which are often experienced as life-threatening. Electrocardiographic findings are without pathological signs. ☐ 1
- The patient complains about occasional or continuing pain in the urinary bladder and/or pelvis accompanied by painful desire to void and/or blood in the urine. There is no bacteriuria. ☐ 1
- The patient complains about occasional or continuing paresthesia (burning, pins and needles, numbness) and/or pain which does not respond to treatment with analgesics. ☐ 1
- Gastroscopy and biopsies from the stomach and duodenum
- are without pathological findings. ☐ 0
  - or
  - show minor signs of inflammation. ☐ 1
  - or
  - show *Helicobacter pylori*- and NSAID-negative erosion and/or ulcer. ☐ 3
  - or
  - show clusters of mast cells and/or a considerable number of spindle-shaped mast cells and/or CD25-positive mast cells. ☐ 10
- Colonoscopy and intestinal biopsies
- are without pathological findings. ☐ 0
  - or
  - show minor signs of inflammation. ☐ 1
  - or
  - show melanosis coli (abuse of anthra-cenediones ruled out). ☐ 1
  - or
  - show clusters of mast cells and/or a considerable number of spindle-shaped mast cells and/or CD25-positive mast cells. ☐ 10
- The patient reports the following signs of episodically occurring symptoms of autonomic dysfunction:
- tachycardia or palpitation/dysrhythmia ☐ 1
  - flush (redness, feeling of heat) ☐ 2
  - hot flash, sweat ☐ 2
  - paroxysmal hypo/hypertension with dizziness to the point of syncope ☐ 2
- Although there are no pathological findings in routine laboratory parameters and imaging methods, the patient present with
- a pronounced asthenia. ☐ 1
  - fatigue. ☐ 1
  - loss in weight. ☐ 1
- During the symptomatic periods of the disorder the patient is afflicted with anal pruritus and/or anal eczema. ☐ 1
- Intestinal adhesions are present without prior history of abdominal surgery ☐ 1

*Triggering factors*

- Deprivation of sleep ☐ 1
- Fasting for 24 h ☐ 1
- Histamine containing food (*e.g.*, red wine, cheese, tuna) ☐ 1

*Laboratory parameters*

- The patient shows signs of a bleeding diathesis (*e.g.*, abnormal secondary bleeding or bruises after minimal trauma and/or lesions). ☐ 1
- During the symptomatic periods of the disorder the patient showed, at least once, hyperbilirubinemia (up to 2.5 mg/dL), and/or an increase of transaminases (up to twice their upper limits of normal) and/or diet-independent hypercholesterolemia (up to 300 mg/dL). ☐ 1
- There are low titers of autoantibodies without clinical signs in the organs or tissues against which the autoantibodies are directed. ☐ 1
- The serum total tryptase was normal. ☐ 0
- or
  - was elevated > 11 and < 20 ng/mL. ☐ 3
  - or
  - was elevated more than 20 ng/mL. ☐ 10
- The level of heparin in blood was normal. ☐ 0

	or	
	was elevated > 0.05 anti-Factor Xa units/mL.	<input type="checkbox"/> 3
The level of N-methylhistamine in a 12-h urine collection was normal.		<input type="checkbox"/> 0
	or	
	was marginally elevated.	<input type="checkbox"/> 1
	or	
	was elevated up to tenfold of the reference value.	<input type="checkbox"/> 5
	or	
	was elevated by more than tenfold of the reference value.	<input type="checkbox"/> 10
<i>Imaging methods</i>		
The patient has splenomegaly and/or hepatomegaly.		<input type="checkbox"/> 1
The patient has bone pain with signs of osteoporosis and/or osteopenia and/or osteosclerosis.		<input type="checkbox"/> 1
<i>Medical history</i>		
The patient shows involvement of the skin in terms of		
	brown-Reddish maculopapulous rash/eruption.	<input type="checkbox"/> 2
	angioedema of the lips, lids of the eye, infraorbital.	<input type="checkbox"/> 2
	pruritus (itching) without rash/eruption and/or disease-related folliculitis	<input type="checkbox"/> 1
	a clear increase in the number of telangiectasias.	<input type="checkbox"/> 1
The patient reports sudden attacks of migraine-like headache.		<input type="checkbox"/> 1
The patient reports memory loss (ability to remember names or words) and/or concentration difficulty and/or sleep disturbances.		<input type="checkbox"/> 1
The patient reports tinnitus attacks and/or ocular discomfort (dry eyes, red eyes, stinging eyes) and/or rhinorrhea/chronic nasal congestion and/or stomatitis (score if two or more of these symptoms are present).		<input type="checkbox"/> 1
The patient reports non-allergic respiratory ailments such as asthma, compulsion to clear the throat, tickling/ticklish feeling in the respiratory tract and/or shortness of breath during routine tasks.		<input type="checkbox"/> 1
In the past, common viral infections of the upper respiratory tract were frequently complicated by bacterial superinfection.		<input type="checkbox"/> 1
The patient can state precisely the date of the first clinical manifestation of the mast cell mediator release syndrome because it appears to him to be associated with an infectious disease.		<input type="checkbox"/> 1

**Figure 4** Validated questionnaire to recognize symptoms as part of a mast cell activation disease in a standardized manner (modified from<sup>[38,119]</sup>). NSAID: Non-steroidal anti-inflammatory drugs.

noses the uncommon low-tryptase SM as MCAS.

Tryptase has been well established as a highly specific MC mediator<sup>[45]</sup>. Its biology is complex; many isoforms with different behaviors and functions have been elucidated<sup>[46]</sup>. It is heat-labile<sup>[47,48]</sup> and has a relatively short half-life *in vivo* (from 6-8 min in healthy subjects to 1.5-2.3 h in patients with hypersensitivity reactions), longer (approximately 4 d) in separated serum<sup>[34,49]</sup>. The WHO 2008 diagnostic criteria for mastocytosis call only for the measurement of total serum tryptase, setting a threshold of 20 ng/mL as the minimum level consistent with a diagnosis of SM<sup>[33]</sup>. It is unclear whether measurement of specific isoforms of tryptase would be beneficial in diagnosing MCAD in any form, and thus at present only the measurement of total serum tryptase can be recommended in the evaluation of a patient suspected of having MCAD. Although the serum total tryptase usually is not elevated in MCAS<sup>[20,26]</sup>, an elevated level (though almost always < 20 ng/mL) can be found in a minority of MCAS patients, and any elevation at all can help buttress a diagnostician's suspicions of the involvement of MC activation in the patient's illness.

If the serum tryptase persistently exceeds 20 ng/mL, or if other clinical features of the presentation are characteristic for SM (*e.g.*, initial onset of symptoms

in middle age, MCs observed in the peripheral blood smear, *etc.*), then further evaluation for SM is warranted. Marrow aspiration/biopsy is a standard part of such evaluation, and given both the patchy distribution of the disease in involved tissues and the observation that unilateral marrow biopsies are non-diagnostic in one-sixth of patients ultimately diagnosed with SM<sup>[50]</sup>, bilateral aspirations/biopsies are preferred. Core biopsies should undergo routine staining as well as MC-targeted immuno/histochemical staining (*e.g.*, CD117, CD25, CD2, tryptase, Giemsa, toluidine blue, Alcian blue, *etc.*). There is no particular subset of these stains which has been established as a standard initial assessment. It has been proposed that CD30, too, be routinely examined when assessing for MCAD<sup>[38]</sup>. Also, CD68 - more classically associated with macrophages - can be displayed by MCs and thus may be useful in diagnosing MCAD<sup>[51]</sup>. All of the above having been said, MCs express surface CD117 (the extracellular portion of *c-kit*) roughly an order of magnitude more brightly than any other CD117-expressing cell, so often only CD117 staining is needed to estimate MC density and to characterize aggregation. Each aspirate should be sent for standard cytogenetic analysis as well as multi-color flow cytometric assessment for co-expression of cell-surface doublets CD117/

CD25 and CD117/CD2; occasionally even the triplet of CD117/CD25/CD2 is seen<sup>[52]</sup>. These flow cytometric signatures are widely considered to be pathognomonic for monoclonal MC disease. Each aspirate should be subjected to the full extent of MC mutational analysis available at the time, though at present this typically is limited to PCR analysis for the KIT<sup>D816V</sup> mutation. If leukocytosis, erythrocytosis, or thrombocytosis is present, assessment is also warranted for the Janus kinase 2 (JAK2) mutations often found in the MPNs (and perhaps the MPL<sup>W515L/K</sup> mutation, too, in cases of leukocytosis or thrombocytosis without erythrocytosis which might signal the presence of essential thrombocythemia or myelofibrosis). Similarly, if prominent eosinophilia is observed, tyrosine kinases platelet-derived growth factor receptor alpha (PDGFR $\alpha$ ), PDGFR $\beta$ , and fibroblast growth factor receptor 1 should be assessed for mutations as recently reviewed elsewhere<sup>[53]</sup>.

The presence of skin lesions characteristic of UP or TMEP warrants biopsy of one or more such lesions, to be immediately processed with routine staining and the above-noted immunohistochemical staining plus assessment for clonality with flow cytometry. As solid tissue flow cytometry requires fresh tissue but PCR mutation analysis does not, it may be reasonable to defer PCR of skin lesion biopsies until other testing finds at least some evidence suggestive of MC disease.

Spontaneously appearing/disappearing, diffusely migratory macular erythematous patchy rashes are common in MCAS, as are scatterings of small ulcerative lesions, too. In the authors' experience, though, biopsies of such sites rarely reveal increased or otherwise aberrant MC populations, suggesting such lesions may be merely the end result of aberrant release of mediators from distantly located MCs.

If clinical suspicion for SM persists (whether due to serum tryptase persistently > 20 ng/mL or other factors) but two or more marrow biopsies (and possibly also biopsies of suspect skin lesions) are negative, upper and lower endoscopic examination of the gastrointestinal tract is reasonable. Biopsies should be performed of any macroscopically apparent lesions, but blind biopsies of macroscopically normal mucosa should be pursued, too, at multiple points along the tract. All such biopsies should be processed as described above for skin biopsies. Several studies have reported that mast cell counts in GI or genitourinary (GU) tract mucosal biopsies normally number fewer than 20 per high power field<sup>[54-58]</sup>. However, cut-offs between "normal" and "abnormal" numbers of non-aggregated mast cells are not addressed in either of the two currently proposed diagnostic schemes for MCAS. Furthermore, it is important to remember that the diagnosis of any form of MCAD rests in meeting a set of criteria, so an isolated finding of "increased" tissue mast cells does not establish the diagnosis regardless of the cut-off used to define "increased", and especially when such cells are not tightly clustered. In the context of other data meeting diagnostic criteria, though, a finding of "increased" tissue mast cells lends additional cre-


dence to the diagnosis.

MCs are highly pleomorphic. On routine hematoxylin and eosin (H and E) staining, MCs can be indistinguishable from lymphocytes, plasma cells, macrophages, histiocytes, or spindle cells<sup>[59,60]</sup>. Thus, in MC disease, not all "lymphocytes" seen with H and E staining are necessarily lymphocytes, and immuno/histochemical staining is required in cases of suspected MCAD. Not infrequently, because of their chronic idiopathic (and often treatment-refractory) gastrointestinal symptoms, patients being evaluated for MCAD have previously undergone GI tract endoscopies with biopsies. Thus, simple reassessment of archived tissue with MC-targeted immuno/histochemical staining may be sufficient to demonstrate that the "benign mucosa" or "mild chronic inflammation" previously seen with H and E staining actually harbors increased MCs. As PCR testing can be performed on preserved tissue, positive immuno/histochemical findings on archived tissue should lead, if sufficient tissue remains, to mutation analysis as possible.

Clinical judgment is required as to whether respiratory or GU tract biopsies are warranted in particularly diagnostically challenging cases, but if pursued, such tissues should be processed as for skin and GI tract biopsies.

Newly or persistently pathologically enlarged (> 1 cm) lymph nodes require biopsy to rule out malignant lymphoproliferative and granulomatous disorders (regardless of whether such disorders can be proven to be independent of, or consequent to, MCAD). However, diffusely migratory, spontaneously waxing/waning adenopathy and/or adenitis is common in MCAD and far more often is a reactive rather than malignant process, though the reactive patterns seen on biopsy can be bizarre (*e.g.*, sinus histiocytosis) and can have the pathologist on the verge of declaring a malignancy. In patients diagnosed with MCAD (or being evaluated for suspected MCAD) who have a longstanding history of spontaneous enlargement of nodes to pathologic size soon followed by spontaneous regression of such adenopathy, the new appearance of a modestly pathologically enlarged lymph node can be watched closely without reflexively proceeding to biopsy, but persistence or progression of adenopathy beyond the range, or outside of the pattern, the patient has usually experienced may warrant biopsy. Node biopsies should be excisional whenever possible and should be submitted not only for the above-noted analyses for MC disease but also for routine analyses for lymphoproliferative disorders and perhaps even - in appropriate clinical settings - specific plasma cell dyscrasias or granulomatous diseases such as amyloidosis, Castleman's disease, or sarcoidosis.

In addition to checking a serum tryptase, other initial laboratory assessment for MCAD (whether suspected to be mastocytosis or MCAS) should include a variety of common tests as listed in Figure 5. As MCAD commonly causes osteolysis (and, less commonly, osteosclerosis - and occasionally even both processes at different sites in the same patient), it is appropriate to establish a baseline for bone densitometry after the diagnosis of



**Figure 5 Diagnostic approach to mast cell activation disease.** MCs: Mast cells; MCAS: Mast cell activation syndrome; GI: Gastrointestinal; PG: Prostaglandin; PCR: Polymerase chain reaction; NSAID: Non-steroidal anti-inflammatory drugs.

MCAD has been established, as the finding of osteopenia or osteoporosis has therapeutic implications. MCAD can drive erythropenia or erythrocytosis (either of which may be macrocytic, normocytic, or microcytic, engendering appropriate differential diagnostic considerations of secondary processes), leukopenia or leukocytosis, and/or thrombocytopenia or thrombocytosis, but there are no findings in the CBC that are specific for MCAD (other than significant numbers of circulating mast cells signaling mast cell leukemia). Also, as previously noted, certain subtle abnormalities are not uncommonly seen in the leukocyte differential, but often the differential is as normal as the CBC. Indeed, it is a confounding point for hematologists that this disease of fundamentally hematologic classification often presents with not a single abnormality in the CBC or differential. However, many MCAD patients - with either mastocytosis or MCAS - do manifest subtle to gross abnormalities in one or more elements of the CBC, providing the diagnostician important opportunities to consider alternative diagnoses. Of note, though, gross abnormalities in the CBC in advanced mastocytosis may be due more to marrow replacement by MCs, while gross abnormalities in the

CBC in less advanced mastocytosis and MCAS are likely due more to MC mediator-driven interference with normal hematopoiesis.

Many MCAD patients also manifest abnormalities in the common serum chemistries, again typically modest in degree but providing the diagnostician opportunities to consider alternative diagnoses regardless of whether they may have risen independently of, or consequent to, MCAD.

Both hypergammaglobulinemia (usually a polyclonal inflammatory phenomenon but occasionally a monoclonal gammopathy of undetermined significance or more advanced clonal plasma cell dyscrasia) and hypogammaglobulinemia can be seen in MCAD<sup>[61]</sup>; patients sometimes have been previously diagnosed with common variable immunodeficiency. Markedly IgG-deficient patients (< 400 mg/dL) who have suffered major infections or substantial healing complications may be candidates for prophylactic immune globulin therapy.

Histories of “easy” bruising and/or bleeding are frequent amongst MCAD patients. Multiple mechanisms are possible (and not necessarily mutually exclusive) including coagulation factor deficiencies, antiphospholipid

antibodies, and/or MC release into the local tissues of (antithrombin-activating) heparin. However, fibrinolysis (as stimulated by heparin release<sup>[62]</sup> and other fibrinolytic mediators such as tissue plasminogen activator<sup>[63]</sup>) appears likely to be the dominant cause, explaining why antifibrinolytics such as tranexamic acid can be helpful in controlling MCAD-driven bleeding. The prothrombin time (PT) and activated partial thromboplastin time (aPTT) remain reasonable screening tests in patients with suspected MCAD who report histories of easy bruising/bleeding, but the diagnostician should be aware that local MC release of mediators with sufficient direct or indirect anticoagulant effect to cause local bleeding usually do not provoke systemic anticoagulant effect sufficient to cause rises in PT or aPTT to abnormal levels. Thus, measurement of fibrinolytic mediators<sup>[63]</sup>, as proximate to the onset of bleeding as possible, may be required to prove fibrinolysis if felt clinically important. Otherwise inexplicable prolongations in PT or aPTT should prompt appropriate further evaluation for clotting factor deficiencies. Also, assessment for antiphospholipid antibodies may not be unreasonable in suspected MCAD patients with otherwise inexplicable abnormalities (high or low) in either PT or aPTT and histories of not only easy bruising/bleeding but also thromboembolism or miscarriage.

If not already performed, a careful evaluation for known hypercoagulable disorders should be performed in the patient with suspected MCAD and a history of idiopathic thromboembolism. Such evaluation should include, among other considerations, determination of the plasma level of Factor VIII (produced by various sources including the MC<sup>[64]</sup> and associated with hypercoagulability<sup>[65]</sup>) and, with a history of portal/splanchnic vein thrombosis, assessment for the JAK2<sup>V617F</sup> mutation which might suggest the occult presence of an MPN<sup>[66]</sup>. Testing for inborn hypercoagulable disorders may not be warranted in cases of initial thromboembolism at ages above 40.

Most patients presenting for evaluation of MCAD have been chronically unwell and have undergone extensive diagnostic evaluation, but the diagnostician considering MCAD nevertheless must carefully review past evaluations and ensure other diagnostic considerations potentially fitting the patient's unique course have been excluded (*e.g.*, hypothyroidism, celiac sprue, Epstein-Barr virus disease, carcinoid, amyloidosis, porphyria, sarcoidosis, *etc.*). Even if the root cause of the patient's multi-system unwellness is MCAD, identification of specific secondary phenomena (*e.g.*, autoimmunity, infection, malignancy, *etc.*) may permit better outcomes via dual-directed therapy.

#### **Laboratory assessment of other MC-specific mediators**

If the serum tryptase is < 20 ng/mL and there are no other clinical hints that SM is more likely than MCAS, then an initial laboratory survey of other MC mediators should be pursued in patients clinically suspected of having MCAS. In the authors' opinion, such an initial media-

tor survey for evidence of MCAS should include serum chromogranin A; chilled plasma for prostaglandin D<sub>2</sub> (PGD<sub>2</sub>) (and/or 11-β-PGF<sub>2α</sub>); chilled plasma histamine; chilled plasma heparin (in patients not on exogenous heparin products); and chilled random and 24-h urine collections for PGD<sub>2</sub> (and/or 11-β-PGF<sub>2α</sub>) and *N*-methylhistamine<sup>[26]</sup>. In some situations (see below), urinary levels of leukotrienes B<sub>4</sub>, C<sub>4</sub>, D<sub>4</sub> and E<sub>4</sub> may also be worth pursuing<sup>[26,67]</sup>.

Although specimen collection as soon as possible following an acute flare of symptoms is ideal, there is no need to wait for such an event when initially assessing the patient with longstanding baseline symptoms consistent with aberrant MC mediator release. However, if the initial laboratory assessment in a patient with a history suspicious for MCAD is negative, repeat testing is usually warranted but preferably should be deferred until the presentation of an acute flare. If possible, hourly determinations of serum tryptase, plasma PGD<sub>2</sub> and histamine, and spot urinary PGD<sub>2</sub> and *N*-methylhistamine should be pursued at baseline and over the next 2-3 h as a flare evolves.

Chromogranin A (CgA) is another known MC product<sup>[35,68]</sup> which appears quite heat-stable<sup>[69]</sup> and was found elevated in the serum in 17% (21 out of 122) of a cohort of MCAD patients investigated (LBA, unpublished data). Elevated serum levels of CgA can be helpful toward diagnosing MCAS, but other potential causes of such elevations (heart failure, renal insufficiency, proton pump inhibitor (PPI) use, and neuroendocrine cancer) should be excluded. Changes in serum CgA level in response to initiation or cessation of PPI therapy are fairly reliable and rapid (within 5 d<sup>[70]</sup>) but highly variable in magnitude<sup>[71,72]</sup>, likely due to multiple mechanisms including pharmacogenomic polymorphisms (largely in cytochrome p450 isoenzyme 2C19<sup>[73]</sup>) and increase in density of CgA-secreting enterochromaffin cells (especially in gastric tissue) in response to gastric acid reduction<sup>[74]</sup>. Cessation of PPI therapy at least 5 days prior to submission of a serum sample for CgA assessment should permit an accurate gauge of the baseline CgA level.

Although it was the first MC mediator to be discovered<sup>[75,76]</sup> and is a highly sensitive and specific indicator of MC activity<sup>[16,63]</sup>, the plasma heparin level is difficult to measure accurately in the clinical setting. The metabolism of heparin is complex<sup>[77]</sup>, but it is clear that it has a short half-life (generally less than one hour) and decomposes rapidly even at chilled temperatures<sup>[63]</sup>. Seidel *et al.*<sup>[63]</sup> reported an upper limit of normal plasma heparin of 0.02 anti-Factor Xa units/mL, but the utility of assessing the plasma heparin level for evidence of MCAS may be constrained by the fact that most of the commercially available assays have a lower limit of detectability of 0.05 or 0.10 anti-Factor Xa units/mL since such assays are used far more to assess anticoagulant adequacy of therapeutic heparinization than to assess endogenous heparin levels. However, about 50% of MCAD patients will manifest a plasma heparin level exceeding 0.05 anti-Factor Xa units/mL on commercially available assays

(105 out of 202 MCAD patients in a current study; unpublished data), and in such patients whose detectable plasma heparin levels cannot be attributed to therapeutic heparin or to the only other known endogenous source of heparin in humans (basophils), MC activation almost certainly is the source.

Like heparin, histamine is slightly less specific for MC disease than tryptase in that histamine is also released by basophils. Unlike heparin, histamine is not particularly heat-labile<sup>[78]</sup> but nevertheless has a short half-life *in vivo*<sup>[79,80]</sup> (estimates have ranged from 1 min to 1 h and may vary based on increased histaminase activity during anaphylaxis<sup>[81]</sup>). Histamine appears to be stable in separated plasma at room temperature for at least 48 h<sup>[82]</sup>. However, because histamine assays often must be performed at distant reference laboratories (requiring specimen storage and transport exceeding 48 h), plasma specimens for histamine levels should be kept continuously chilled at all stages of handling. Histamine's major metabolite, N-methylhistamine, has a longer half-life *in vivo*<sup>[80]</sup> and may be a preferable measure over histamine, especially when a sample cannot be collected during a flare of symptoms or when other constraints preclude measurement of both molecules.

Although also produced in macrophages<sup>[83-86]</sup>, Langerhans cells<sup>[87]</sup>, liver endothelium<sup>[86]</sup>, platelets<sup>[88]</sup>, Th2 helper T cells<sup>[89]</sup>, stimulated osteoblasts<sup>[90]</sup>, and possibly adipocytes<sup>[91]</sup>, PGD<sub>2</sub> appears to be far dominantly produced in MCs<sup>[92,93]</sup>, yielding attractive specificity for clinical detection of MCAD. In patients with MC activation producing increases in urinary N-methylhistamine, the fold increase in urinary PGD<sub>2</sub> is substantially greater than seen for the histamine metabolite<sup>[94]</sup>. However, PGD<sub>2</sub> appears to have an even shorter half-life than histamine (on the order of 1-30 min in various studies)<sup>[95-97]</sup>. In fact, PGD<sub>2</sub> is metabolized so rapidly that its measured levels may substantially underestimate its total production<sup>[98]</sup>. Some PGD<sub>2</sub> metabolites, though, are more stable than the parent compound (*e.g.*, 9 $\alpha$ ,11 $\beta$ -PGF<sub>2 $\alpha$</sub> )<sup>[92]</sup>, leading some to preferentially test such metabolites over PGD<sub>2</sub>. Of note, though, far more PGF<sub>2 $\alpha$</sub>  comes from reduction of PGH<sub>2</sub>, and even some from PGE<sub>2</sub>, which are produced by a range of other types of cells<sup>[99]</sup>, making the value of PGF<sub>2 $\alpha$</sub>  as a marker for MC activity less clear). Assays of levels of histamine and prostaglandin metabolites are further complicated by the need of many clinical laboratories to ship samples to a reference laboratory for measurement. Care must be taken by the patient and laboratory staff to maintain the samples (particularly for heparin and PGD<sub>2</sub>) in chilled condition throughout collection, storage, and transport until final processing. It is unclear whether urinary PGD<sub>2</sub> is exclusively a product of the kidneys<sup>[100]</sup> or dominantly a filtering of plasma PGD<sub>2</sub><sup>[92]</sup>. If the former is true, a low urinary PGD<sub>2</sub> could be due to chronic kidney disease (CKD), though certainly patients with CKD due to activated MCs can manifest elevated urinary PGD<sub>2</sub> levels. Therefore, CKD is not a reason to forego testing of the urinary PGD<sub>2</sub> level.

Ideally, a 24-h urine collection is preferred over a spot/random collection given that the evanescence of mediator flares and the short half-lives of many mediators make spot urine collections and plasma assays less likely to catch elevated levels of mediators. However, from an efficiency perspective, testing both 24-h and random specimens at the same time may not be unreasonable in patients whose mysterious chronic illness has already consumed extensive resources.

Non-steroidal anti-inflammatory drugs (NSAIDs), which inhibit cyclooxygenase and thus limit prostaglandin production, can result in low PGD<sub>2</sub> levels in the blood and urine. Similar to the precaution with PPI therapy when assessing for serum CgA elevation, the patient's history of use of NSAIDs, in both prescription and over-the-counter products, should be reviewed when planning the laboratory assessment for prostaglandin levels. Like most PPIs, some NSAIDs have long half-lives permitting daily dosing, and thus patients should be cautioned to abstain, if possible, from all NSAIDs for at least 5 d prior to specimen collection for assessment of prostaglandins.

Given the heterogeneity of MCAD and the above-noted challenges in detecting elevated levels of MC mediators, it should not be surprising that some patients present not only with histories which are classic for MCAS but also with initial screenings for the most sensitive and specific MC mediators which are normal. When repeated efforts to identify aberrant MC activation using the above-described screening approach all fail, consideration can be given to screening for aberrant expression of less specific MC mediators such as Factor VIII<sup>[64,101]</sup>, plasma free norepinephrine<sup>[102]</sup>, tumor necrosis factor alpha<sup>[103-111]</sup> and interleukin-6<sup>[103]</sup>. Plasma free norepinephrine is often ordered as part of a plasma free catecholamine profile; in many MCAS patients a pattern of elevated norepinephrine, low-normal or low epinephrine, and sometimes elevated dopamine is seen (LBA and GJM, unpublished data).

Although commercial testing (whether through local or reference laboratories) is not yet widely available, levels of MC mediators leukotriene B<sub>4</sub> and cysteinyl leukotrienes C<sub>4</sub>, D<sub>4</sub> and E<sub>4</sub> have been reported to be elevated in patients with SM as compared to healthy controls<sup>[67]</sup>. Plasma and urinary leukotrienes also increase with acute attacks of asthma<sup>[112]</sup>, a disease increasingly suspected to involve aberrantly active MCs<sup>[113]</sup>. However, leukotriene expression has not been specifically studied in the MCAS population. Furthermore, leukotrienes are synthesized in a variety of myeloid cells (and epithelial cells as well)<sup>[114,115]</sup>, so the specificity of elevated leukotriene metabolite levels for MCAD is unclear and may depend substantially on the presence of a clinical history compatible with MCAD. Whether elevated leukotriene levels are related more to MC load or to MC activity remains to be determined. However, if the clinical history is best explained by MC activation but abnormal levels of the previously discussed biochemical markers cannot be identified, it would not be unreasonable to also exam-

ine levels of plasma and urinary leukotriene metabolites if commercial assays of such are accessible.

## SPECIMEN HANDLING ISSUES

The short half-lives and thermolability of many MC mediators require continuous specimen chilling throughout collection, storage, and transport. Particularly with regard to 24-h urine collections for MC mediator testing, patients should be carefully educated to pre-chill the collection container overnight before beginning the collection and then to keep the container continuously chilled while following an otherwise standard 24-h urine collection protocol; the container should be removed from the refrigerator or ice chest only when imminently needed and should be returned to chilling as soon as possible. Patients should also be cautioned to maintain the container in a chilled environment throughout transport. We recommend the container be placed in a bag filled with ice and sealed, with the bag then placed in an ice chest filled with ice and sealed for transport to the accessioning lab, whereupon the bag can be removed from the chest and provided to the technician with a reminder of the criticality of keeping the specimen chilled.

Laboratory staff often are unfamiliar with MC mediator testing, previously a rarely undertaken endeavor. Consideration should be given by the diagnostician to sharing with laboratory personnel the ultimate clinical goal of such testing and the importance of maintaining thermal integrity of these specimens at all times, including at the time of initial accessioning as well as when packing specimens for transport to reference laboratories that may be thousands of miles distant, transits that may involve long periods sitting in unventilated cargo containers on hot tarmacs. Use of well-insulated containers, and liberal placement of cold packs in the insulated container, should be *de rigueur* when packing such specimens for long-distance transport.

If PGD<sub>2</sub> (or 11-β-PGF<sub>2α</sub>) levels below the lower limit of normal are determined and the patient denies any recent use of NSAIDs, or if “normal” MC mediator levels are repeatedly seen despite specimens being accessioned at particularly symptomatic times, it may be useful to ask the patient about his observations, while at the lab, as to whether the laboratory staff maintained the specimen in a chilled environment. Aside from NSAID use, loss of thermal integrity is the most common reason for low PGD<sub>2</sub> levels in the authors’ experience.

## CONCLUSION

With growing appreciation of the prevalence of MCAD (particularly MCAS), there is a growing need to pursue diagnostic evaluation for MCAD outside of the few centers specializing in this area. An understanding of the spectrum of MCAD and the similarities and differences amongst the various forms of the disease is helpful in guiding the approach to testing. Especially if MCAS is

confirmed to usually be a clonal disease, the rise of personal genomics may come to obviate many of the present challenges in diagnosing MCAD, but with appropriate collaboration amongst the diagnostician, patient, other relevant clinicians, and laboratory staff, today’s challenges can be surmounted. Especially given that the majority of the MCAD iceberg seems likely to be MCAS and that significantly helpful therapy can be found for most MCAD patients (including most MCAS patients), efforts to smooth the path toward diagnosis of MCAD are worthwhile.

## REFERENCES

- 1 Kalesnikoff J, Galli SJ. New developments in mast cell biology. *Nat Immunol* 2008; **9**: 1215-1223 [PMID: 18936782 DOI: 10.1038/ni.f.216]
- 2 von Recklinghausen F. Ueber Eiter- und Bindegewebskörperchen. *Virchows Arch* 1863; **28**: 157-197 [DOI: 10.1007/BF01930779]
- 3 Unna P. Beiträge zur Anatomie und Pathogenese der Urticaria simplex und pigmentosa. *Monatschrift der praktischen Dermatologie* 1887; **6**: 9-18
- 4 Ellis JM. Urticaria pigmentosa; a report of a case with autopsy. *Arch Pathol (Chic)* 1949; **48**: 426-435 [PMID: 18149230]
- 5 Efrati P, Klajman A, Spitz H. Mast cell leukemia? Malignant mastocytosis with leukemia-like manifestations. *Blood* 1957; **12**: 869-882 [PMID: 13471655]
- 6 Furitsu T, Tsujimura T, Tono T, Ikeda H, Kitayama H, Koshimizu U, Sugahara H, Butterfield JH, Ashman LK, Kanayama Y. Identification of mutations in the coding sequence of the proto-oncogene c-kit in a human mast cell leukemia cell line causing ligand-independent activation of c-kit product. *J Clin Invest* 1993; **92**: 1736-1744 [PMID: 7691885 DOI: 10.1172/JCI116761]
- 7 Akin C, Metcalfe DD. The biology of Kit in disease and the application of pharmacogenetics. *J Allergy Clin Immunol* 2004; **114**: 13-19; quiz 20 [PMID: 15241338 DOI: 10.1016/j.jaci.2004.04.046]
- 8 Nagata H, Worobec AS, Oh CK, Chowdhury BA, Tannenbaum S, Suzuki Y, Metcalfe DD. Identification of a point mutation in the catalytic domain of the protooncogene c-kit in peripheral blood mononuclear cells of patients who have mastocytosis with an associated hematologic disorder. *Proc Natl Acad Sci USA* 1995; **92**: 10560-10564 [PMID: 7479840]
- 9 Garcia-Montero AC, Jara-Acevedo M, Teodosio C, Sanchez ML, Nunez R, Prados A, Aldanondo I, Sanchez L, Dominguez M, Botana LM, Sanchez-Jimenez F, Sotlar K, Almeida J, Escribano L, Orfao A. KIT mutation in mast cells and other bone marrow hematopoietic cell lineages in systemic mast cell disorders: a prospective study of the Spanish Network on Mastocytosis (REMA) in a series of 113 patients. *Blood* 2006; **108**: 2366-2372 [PMID: 16741248 DOI: 10.1182/blood-2006-04-015545]
- 10 D'Ambrosio C, Akin C, Wu Y, Magnusson MK, Metcalfe DD. Gene expression analysis in mastocytosis reveals a highly consistent profile with candidate molecular markers. *J Allergy Clin Immunol* 2003; **112**: 1162-1170 [PMID: 14657877 DOI: 10.1016/j.jaci.2003.07.008]
- 11 Orfao A, Garcia-Montero AC, Sanchez L, Escribano L. Recent advances in the understanding of mastocytosis: the role of KIT mutations. *Br J Haematol* 2007; **138**: 12-30 [PMID: 17555444 DOI: 10.1111/j.1365-2141.2007.06619.x]
- 12 Mayerhofer M, Gleixner KV, Hoelbl A, Florian S, Hoermann G, Aichberger KJ, Bilban M, Esterbauer H, Krauth MT, Sperr WR, Longley JB, Kralovics R, Moriggl R, Zappulla J, Liblau RS, Schwarzwinger I, Sexl V, Sillaber C, Valent P. Unique effects of KIT D816V in BaF3 cells: induction of cluster forma-

- tion, histamine synthesis, and early mast cell differentiation antigens. *J Immunol* 2008; **180**: 5466-5476 [PMID: 18390729]
- 13 **Haenisch B**, Nöthen MM, Molderings GJ. Systemic mast cell activation disease: the role of molecular genetic alterations in pathogenesis, heritability and diagnostics. *Immunology* 2012; **137**: 197-205 [PMID: 22957768 DOI: 10.1111/j.1365-2567.2012.03627.x]
  - 14 **Broesby-Olsen S**, Kristensen T, Vestergaard H, Brixen K, Møller MB, Bindslev-Jensen C. KIT D816V mutation burden does not correlate to clinical manifestations of indolent systemic mastocytosis. *J Allergy Clin Immunol* 2013; In press [DOI: 10.1016/j.jaci.2013.02.019]
  - 15 **Lawley W**, Hird H, Mallinder P, McKenna S, Hargadon B, Murray A, Bradding P. Detection of an activating c-kit mutation by real-time PCR in patients with anaphylaxis. *Mutat Res* 2005; **572**: 1-13 [PMID: 15790486 DOI: 10.1016/j.mrfmmm.2004.08.015]
  - 16 **Roberts LJ**, Oates JA. Biochemical diagnosis of systemic mast cell disorders. *J Invest Dermatol* 1991; **96**: 19S-24S; discussion 24S-25S; 60S-65S [PMID: 16799604 DOI: 10.1111/1523-1747.ep12468945]
  - 17 **Sonneck K**, Florian S, Müllauer L, Wimazal F, Födinger M, Sperr WR, Valent P. Diagnostic and subdiagnostic accumulation of mast cells in the bone marrow of patients with anaphylaxis: Monoclonal mast cell activation syndrome. *Int Arch Allergy Immunol* 2007; **142**: 158-164 [PMID: 17057414 DOI: 10.1159/000096442]
  - 18 **Akin C**, Scott LM, Kocabas CN, Kushnir-Sukhov N, Brittain E, Noel P, Metcalfe DD. Demonstration of an aberrant mast-cell population with clonal markers in a subset of patients with "idiopathic" anaphylaxis. *Blood* 2007; **110**: 2331-2333 [PMID: 17638853 DOI: 10.1182/blood-2006-06-028100]
  - 19 **Molderings GJ**, Kolck UW, Scheurlen C, Brüss M, Homann J, Von Kügelgen I. Multiple novel alterations in Kit tyrosine kinase in patients with gastrointestinally pronounced systemic mast cell activation disorder. *Scand J Gastroenterol* 2007; **42**: 1045-1053 [PMID: 17710669 DOI: 10.1080/00365520701245744]
  - 20 **Molderings GJ**, Meis K, Kolck UW, Homann J, Frieling T. Comparative analysis of mutation of tyrosine kinase kit in mast cells from patients with systemic mast cell activation syndrome and healthy subjects. *Immunogenetics* 2010; **62**: 721-727 [PMID: 20838788 DOI: 10.1007/s00251-010-0474-8]
  - 21 **Kralovics R**. Genetic complexity of myeloproliferative neoplasms. *Leukemia* 2008; **22**: 1841-1848 [PMID: 18754034 DOI: 10.1038/leu.2008.233]
  - 22 **Schwaab J**, Schnittger S, Sotlar K, Walz C, Fabarius A, Pfirmann M, Kohlmann A, Grossmann V, Meggendorfer M, Horny HP, Valent P, Jawhar M, Teichmann M, Metzgeroth G, Erben P, Ernst T, Hochhaus A, Haferlach T, Hofmann WK, Cross NC, Reiter A. Comprehensive mutational profiling in advanced systemic mastocytosis. *Blood* 2013; **122**: 2460-2466 [PMID: 23958953 DOI: 10.1182/blood-2013-04-496448]
  - 23 **Akin C**, Valent P, Metcalfe DD. Mast cell activation syndrome: Proposed diagnostic criteria. *J Allergy Clin Immunol* 2010; **126**: 1099-1104.e4 [PMID: 21035176 DOI: 10.1016/j.jaci.2010.08.035]
  - 24 **Molderings GJ**, Haenisch B, Bogdanow M, Fimmers R, Nöthen MM. Familial occurrence of systemic mast cell activation disease. *PLoS One* 2013; **8**: e76241 [PMID: 24098785 DOI: 10.1371/journal.pone.0076241]
  - 25 **Nakamura Y**, Kambe N, Saito M, Nishikomori R, Kim YG, Murakami M, Núñez G, Matsue H. Mast cells mediate neutrophil recruitment and vascular leakage through the NLRP3 inflammasome in histamine-independent urticaria. *J Exp Med* 2009; **206**: 1037-1046 [PMID: 19364881 DOI: 10.1084/jem.20082179]
  - 26 **Molderings GJ**, Brettner S, Homann J, Afrin LB. Mast cell activation disease: a concise practical guide for diagnostic workup and therapeutic options. *J Hematol Oncol* 2011; **4**: 10 [PMID: 21418662 DOI: 10.1186/1756-8722-4-10]
  - 27 **Afrin LB**. Mast cell activation disorder masquerading as pure red cell aplasia. *Int J Hematol* 2010; **91**: 907-908 [PMID: 20526893 DOI: 10.1007/s12185-010-0605-x]
  - 28 **Afrin LB**. Polycythemia from mast cell activation syndrome: lessons learned. *Am J Med Sci* 2011; **342**: 44-49 [PMID: 21642812 DOI: 10.1097/MAJ.0b013e31821d41dd]
  - 29 **Afrin LB**. Mast cell activation syndrome masquerading as agranulocytosis. *Mil Med* 2012; **177**: 113-117 [PMID: 22338992]
  - 30 **Travis WD**, Li CY, Bergstralh EJ, Yam LT, Sweet RG. Systemic mast cell disease. Analysis of 58 cases and literature review. *Medicine* (Baltimore) 1988; **67**: 345-368 [PMID: 3054417]
  - 31 **Travis WD**, Li CY, Yam LT, Bergstralh EJ, Sweet RG. Significance of systemic mast cell disease with associated hematologic disorders. *Cancer* 1988; **62**: 965-972 [PMID: 3409177]
  - 32 **Valent P**, Horny HP, Escribano L, Longley BJ, Li CY, Schwartz LB, Marone G, Nuñez R, Akin C, Sotlar K, Sperr WR, Wolff K, Brunning RD, Parwaresch RM, Austen KF, Lennert K, Metcalfe DD, Vardiman JW, Bennett JM. Diagnostic criteria and classification of mastocytosis: a consensus proposal. *Leuk Res* 2001; **25**: 603-625 [PMID: 11377686 DOI: 10.1016/S0145-2126(01)00038-8]
  - 33 **Horny HP**, Metcalfe DD, Bennett J, Bain BJ, Akin C, Escribano L, Valent P. Mastocytosis. In: Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, Thiele J, Vardiman JW. WHO Classification of Tumors of Hematopoietic and Lymphoid Tissues (4th edition.). Lyon, France: International Agency for Research and Cancer, 2008: 54-63
  - 34 **Valent P**, Akin C, Arock M, Brockow K, Butterfield JH, Carter MC, Castells M, Escribano L, Hartmann K, Lieberman P, Nedoszytko B, Orfao A, Schwartz LB, Sotlar K, Sperr WR, Triggiani M, Valenta R, Horny HP, Metcalfe DD. Definitions, criteria and global classification of mast cell disorders with special reference to mast cell activation syndromes: a consensus proposal. *Int Arch Allergy Immunol* 2012; **157**: 215-225 [PMID: 22041891 DOI: 10.1159/000328760]
  - 35 **Ibelgaufits H**. Mast Cells. In: COPE: Cytokines and Cells Online Pathfinder Encyclopaedia, 2013, Sept. 3. Available from: URL: [http://www.copewithcytokines.de/cope.cgi?key=mast cells](http://www.copewithcytokines.de/cope.cgi?key=mast%20cells)
  - 36 **Afrin L**. Presentation, Diagnosis, and Management of Mast Cell Activation Syndrome. In: Murray D. Mast Cells: Phenotypic Features, Biological Functions, and Role in Immunity. Happaugue, NY: Nova Science Publishers, 2013: 155-231
  - 37 **Hermine O**, Lortholary O, Leventhal PS, Catteau A, Soppelsa F, Baude C, Cohen-Akenine A, Palmérini F, Hanssens K, Yang Y, Sobol H, Fraytag S, Ghez D, Suarez F, Barete S, Casassus P, Sans B, Arock M, Kinet JP, Dubreuil P, Moussy A. Case-control cohort study of patients' perceptions of disability in mastocytosis. *PLoS One* 2008; **3**: e2266 [PMID: 18509466 DOI: 10.1371/journal.pone.0002266]
  - 38 **Valent P**, Sperr WR, Akin C. How I treat patients with advanced systemic mastocytosis. *Blood* 2010; **116**: 5812-5817 [PMID: 20855864 DOI: 10.1182/blood-2010-08-292144]
  - 39 **Alfter K**, von Kügelgen I, Haenisch B, Frieling T, Hülsdonk A, Haars U, Rolfs A, Noe G, Kolck UW, Homann J, Molderings GJ. New aspects of liver abnormalities as part of the systemic mast cell activation syndrome. *Liver Int* 2009; **29**: 181-186 [PMID: 18662284 DOI: 10.1111/j.1478-3231.2008.01839.x]
  - 40 **Schwartz LB**, Sakai K, Bradford TR, Ren S, Zweiman B, Worobec AS, Metcalfe DD. The alpha form of human tryptase is the predominant type present in blood at baseline in normal subjects and is elevated in those with systemic mastocytosis. *J Clin Invest* 1995; **96**: 2702-2710 [PMID: 8675637 DOI: 10.1172/JCI18337]
  - 41 **Borer-Reinhold M**, Haerberli G, Bitzenhofer M, Jandus P, Hausmann O, Fricker M, Helbling A, Müller U. An increase in serum tryptase even below 11.4 ng/mL may indicate a mast cell-mediated hypersensitivity reaction: a prospective

- study in Hymenoptera venom allergic patients. *Clin Exp Allergy* 2011; **41**: 1777-1783 [PMID: 22092437 DOI: 10.1111/j.1365-2222.2011.03848.x]
- 42 **Sperr WR**, Jordan JH, Fiegl M, Escribano L, Bellas C, Dirnhofer S, Semper H, Simonitsch-Klupp I, Horny HP, Valent P. Serum tryptase levels in patients with mastocytosis: correlation with mast cell burden and implication for defining the category of disease. *Int Arch Allergy Immunol* 2002; **128**: 136-141 [PMID: 12065914 DOI: 10.1159/000059404]
  - 43 **Alvarez-Twose I**, González de Olano D, Sánchez-Muñoz L, Matito A, Esteban-López MI, Vega A, Mateo MB, Alonso Díaz de Durana MD, de la Hoz B, Del Pozo Gil MD, Caballero T, Rosado A, Sánchez-Matas I, Teodósio C, Jara-Acevedo M, Mollejo M, García-Montero A, Orfao A, Escribano L. Clinical, biological, and molecular characteristics of clonal mast cell disorders presenting with systemic mast cell activation symptoms. *J Allergy Clin Immunol* 2010; **125**: 1269-1278. e2 [PMID: 20434205 DOI: 10.1016/j.jaci.2010.02.019]
  - 44 **van Doormaal JJ**, van der Veer E, van Voorst Vader PC, Kluin PM, Mulder AB, van der Heide S, Arends S, Kluin-Nelemans JC, Oude Elberink JN, de Monchy JG. Tryptase and histamine metabolites as diagnostic indicators of indolent systemic mastocytosis without skin lesions. *Allergy* 2012; **67**: 683-690 [PMID: 22435702 DOI: 10.1111/j.1398-9995.2012.02809.x]
  - 45 **Schwartz LB**, Metcalfe DD, Miller JS, Earl H, Sullivan T. Tryptase levels as an indicator of mast-cell activation in systemic anaphylaxis and mastocytosis. *N Engl J Med* 1987; **316**: 1622-1626 [PMID: 3295549 DOI: 10.1056/NEJM198706253162603]
  - 46 **Hallgren J**, Pejler G. Biology of mast cell tryptase. An inflammatory mediator. *FEBS J* 2006; **273**: 1871-1895 [PMID: 16640553 DOI: 10.1111/j.1742-4658.2006.05211.x]
  - 47 **Schwartz LB**, Bradford TR. Regulation of tryptase from human lung mast cells by heparin. Stabilization of the active tetramer. *J Biol Chem* 1986; **261**: 7372-7379 [PMID: 3519608]
  - 48 **Fajardo I**, Pejler G. Human mast cell beta-tryptase is a gelatinase. *J Immunol* 2003; **171**: 1493-1499 [PMID: 12874242]
  - 49 **Goldstein SM**, Wintroub BU. Mast cell proteases. In: Kaliner MA, Metcalfe DD. *The Mast Cell in Health and Disease*. New York: Marcel Dekker, 1993: 343-380
  - 50 **Butterfield JH**, Li CY. Bone marrow biopsies for the diagnosis of systemic mastocytosis: is one biopsy sufficient? *Am J Clin Pathol* 2004; **121**: 264-267 [PMID: 14983941 DOI: 10.1309/2EWQKN00PG02JKY0]
  - 51 **Horny HP**, Valent P. Diagnosis of mastocytosis: general histopathological aspects, morphological criteria, and immunohistochemical findings. *Leuk Res* 2001; **25**: 543-551 [PMID: 11377679 DOI: 10.1016/S0145-2126(01)00021-2]
  - 52 **Escribano L**, Orfao A, Díaz-Agustín B, Villarrubia J, Cerveró C, López A, Marcos MA, Bellas C, Fernández-Cañadas S, Cuevas M, Sánchez A, Velasco JL, Navarro JL, Miguel JF. Indolent systemic mast cell disease in adults: immunophenotypic characterization of bone marrow mast cells and its diagnostic implications. *Blood* 1998; **91**: 2731-2736 [PMID: 9531582]
  - 53 **Havelange V**, Demoulin JB. Review of current classification, molecular alterations, and tyrosine kinase inhibitor therapies in myeloproliferative disorders with hypereosinophilia. *J Blood Med* 2013; **4**: 111-121 [PMID: 23976869 DOI: 10.2147/JBM.S33142]
  - 54 **Yeom JS**, Choi MB, Seo JH, Park JS, Lim JY, Park CH, Woo HO, Youn HS, Ko GH, Baik SC, Lee WK, Cho MJ, Rhee KH. Relationship between headache and mucosal mast cells in pediatric *Helicobacter pylori*-negative functional dyspepsia. *Cephalalgia* 2013; **33**: 323-329 [PMID: 23291287 DOI: 10.1177/0333102412472070]
  - 55 **Zare-Mirzaie A**, Lotfi M, Sadeghipour A, Haghi-Ashtiani MT. Analysis of colonic mucosa mast cell count in patients with chronic diarrhea. *Saudi J Gastroenterol* 2012; **18**: 322-326 [PMID: 23006460 DOI: 10.4103/1319-3767.101128]
  - 56 **Martínez C**, Lobo B, Pigrau M, Ramos L, González-Castro AM, Alonso C, Guilarte M, Guilá M, de Torres I, Azpiroz F, Santos J, Vicario M. Diarrhoea-predominant irritable bowel syndrome: an organic disorder with structural abnormalities in the jejunal epithelial barrier. *Gut* 2013; **62**: 1160-1168 [PMID: 22637702 DOI: 10.1136/gutjnl-2012-302093]
  - 57 **Vivinus-Nébot M**, Dainese R, Anty R, Saint-Paul MC, Nano JL, Gonthier N, Marjoux S, Frin-Mathy G, Bernard G, Hébuterne X, Tran A, Theodorou V, Piche T. Combination of allergic factors can worsen diarrheic irritable bowel syndrome: role of barrier defects and mast cells. *Am J Gastroenterol* 2012; **107**: 75-81 [PMID: 21931380 DOI: 10.1038/ajg.2011.315]
  - 58 **Bassotti G**, Villanacci V, Nascimbeni R, Cadei M, Manenti S, Sabatino G, Maurer CA, Cathomas G, Salerni B. Colonic mast cells in controls and slow transit constipation patients. *Aliment Pharmacol Ther* 2011; **34**: 92-99 [PMID: 21539589 DOI: 10.1111/j.1365-2036.2011.04684.x]
  - 59 **Brunning RD**, McKenna RW, Rosai J, Parkin JL, Risdall R. Systemic mastocytosis. Extracutaneous manifestations. *Am J Surg Pathol* 1983; **7**: 425-438 [PMID: 6614308]
  - 60 **Swieter M**, Lee TD, Stead RH, Fujimaki H, Befus D. Mast cell pleomorphism: properties of intestinal mast cells. *Adv Exp Med Biol* 1987; **216A**: 613-623 [PMID: 2446472]
  - 61 **Worobec AS**, Semere T, Nagata H, Metcalfe DD. Clinical correlates of the presence of the Asp816Val c-kit mutation in the peripheral blood mononuclear cells of patients with mastocytosis. *Cancer* 1998; **83**: 2120-2129 [PMID: 9827716]
  - 62 **Upchurch GR**, Valeri CR, Khuri SF, Rohrer MJ, Welch GN, MacGregor H, Ragno G, Francis S, Rodino LJ, Michelson AD, Loscalzo J. Effect of heparin on fibrinolytic activity and platelet function in vivo. *Am J Physiol* 1996; **271**: H528-H534 [PMID: 8770093]
  - 63 **Seidel H**, Molderings GJ, Oldenburg J, Meis K, Kolck UW, Homann J, Hertfelder HJ. Bleeding diathesis in patients with mast cell activation disease. *Thromb Haemost* 2011; **106**: 987-989 [PMID: 21901238 DOI: 10.1160/TH11-05-0351]
  - 64 **Kindblom LG**. Factor VIII related antigen and mast cells. *Acta Pathol Microbiol Immunol Scand A* 1982; **90**: 437-439 [PMID: 6187180]
  - 65 **Kyrle PA**, Minar E, Hirschl M, Bialonczyk C, Stain M, Schneider B, Weltermann A, Speiser W, Lechner K, Eichinger S. High plasma levels of factor VIII and the risk of recurrent venous thromboembolism. *N Engl J Med* 2000; **343**: 457-462 [PMID: 10950667 DOI: 10.1056/NEJM200008173430702]
  - 66 **Kiladjian JJ**, Cervantes F, Leebeek FW, Marzac C, Cassinat B, Chevret S, Cazals-Hatem D, Plessier A, Garcia-Pagan JC, Darwish Murad S, Raffa S, Janssen HL, Gardin C, Cereja S, Tonetti C, Giraudier S, Condat B, Casadevall N, Fenaux P, Valla DC. The impact of JAK2 and MPL mutations on diagnosis and prognosis of splanchic vein thrombosis: a report on 241 cases. *Blood* 2008; **111**: 4922-4929 [PMID: 18250227 DOI: 10.1182/blood-2007-11-125328]
  - 67 **Raithel M**, Zopf Y, Kimpel S, Naegel A, Molderings GJ, Buchwald F, Schultis HW, Kressel J, Hahn EG, Konturek P. The measurement of leukotrienes in urine as diagnostic option in systemic mastocytosis. *J Physiol Pharmacol* 2011; **62**: 469-472 [PMID: 22100848]
  - 68 **Prasad P**, Yanagihara AA, Small-Howard AL, Turner H, Stokes AJ. Secretogranin III directs secretory vesicle biogenesis in mast cells in a manner dependent upon interaction with chromogranin A. *J Immunol* 2008; **181**: 5024-5034 [PMID: 18802106]
  - 69 **Takiyuddin MA**, Cervenka JH, Hsiao RJ, Barbosa JA, Parmer RJ, O'Connor DT. Chromogranin A. Storage and release in hypertension. *Hypertension* 1990; **15**: 237-246 [PMID: 2406199 DOI: 10.1161/01.HYP.15.3.237]
  - 70 **Pregun I**, Herszényi L, Juhász M, Miheller P, Hritz I, Patócs A, Rác K, Tulassay Z. Effect of proton-pump inhibitor therapy on serum chromogranin A level. *Digestion* 2011; **84**: 22-28 [PMID: 21304238 DOI: 10.1159/000321535]

- 71 **Sanduleanu S**, Stridsberg M, Jonkers D, Hameeteman W, Biemond I, Lundqvist G, Lamers C, Stockbrügger RW. Serum gastrin and chromogranin A during medium- and long-term acid suppressive therapy: a case-control study. *Aliment Pharmacol Ther* 1999; **13**: 145-153 [PMID: 10102943 DOI: 10.1046/j.1365-2036.1999.00466.x]
- 72 **Fossmark R**, Jianu CS, Martinsen TC, Qvigstad G, Syversen U, Waldum HL. Serum gastrin and chromogranin A levels in patients with fundic gland polyps caused by long-term proton-pump inhibition. *Scand J Gastroenterol* 2008; **43**: 20-24 [PMID: 18938772 DOI: 10.1080/00365520701561959]
- 73 **Furuta T**, Shirai N, Sugimoto M, Ohashi K, Ishizaki T. Pharmacogenomics of proton pump inhibitors. *Pharmacogenomics* 2004; **5**: 181-202 [PMID: 15016609 DOI: 10.1517/phgs.5.2.181.27483]
- 74 **Sanduleanu S**, De Bruïne A, Stridsberg M, Jonkers D, Biemond I, Hameeteman W, Lundqvist G, Stockbrügger RW. Serum chromogranin A as a screening test for gastric enterochromaffin-like cell hyperplasia during acid-suppressive therapy. *Eur J Clin Invest* 2001; **31**: 802-811 [PMID: 11589723 DOI: 10.1046/j.1365-2362.2001.00890.x]
- 75 **Holmgren H**, Wilander O. Beiträge zur Kenntnis der Chemie und Funktion der Ehrlichschen Mastzellen. *Z Mikroskop Anat Forsch* 1937; **42**: 242-278
- 76 **Jorpes E**, Holmgren H, Wilander O. Über das Vorkommen von Heparin in den Gefäßwänden und in den Augen. *Z Mikroskop Anat Forsch* 1937; **42**: 279-300
- 77 **Hirsh J**, Anand SS, Halperin JL, Fuster V. Guide to anti-coagulant therapy: Heparin : a statement for healthcare professionals from the American Heart Association. *Circulation* 2001; **103**: 2994-3018 [PMID: 11413093 DOI: 10.1161/01.CIR.103.24.2994]
- 78 **Emborg J**, Laursen BG, Dalgaard P. Significant histamine formation in tuna (*Thunnus albacares*) at 2 degrees C--effect of vacuum- and modified atmosphere-packaging on psychrotolerant bacteria. *Int J Food Microbiol* 2005; **101**: 263-279 [PMID: 15925710 DOI: 10.1016/j.ijfoodmicro.2004.12.001]
- 79 **Laroche D**, Vergnaud MC, Sillard B, Soufarapis H, Bricard H. Biochemical markers of anaphylactoid reactions to drugs. Comparison of plasma histamine and tryptase. *Anesthesiology* 1991; **75**: 945-949 [PMID: 1741515]
- 80 **Takeda J**, Ueda E, Takahashi J, Fukushima K. Plasma N-methylhistamine concentration as an indicator of histamine release by intravenous d-tubocurarine in humans: preliminary study in five patients by radioimmunoassay kits. *Anesth Analg* 1995; **80**: 1015-1017 [PMID: 7537026]
- 81 **Lake AM**, Kagey-Sobotka A, Jakubowicz T, Lichtenstein LM. Histamine release in acute anaphylactic enteropathy of the rat. *J Immunol* 1984; **133**: 1529-1534 [PMID: 6205084]
- 82 **Laroche D**, Dubois F, Gérard JL, Lefrançois C, André B, Vergnaud MC, Dubus L, Bricard H. Radioimmunoassay for plasma histamine: a study of false positive and false negative values. *Br J Anaesth* 1995; **74**: 430-437 [PMID: 7734264 DOI: 10.1093/bja/74.4.430]
- 83 **Meyers CD**, Liu P, Kamanna VS, Kashyap ML. Nicotinic acid induces secretion of prostaglandin D2 in human macrophages: an in vitro model of the niacin flush. *Atherosclerosis* 2007; **192**: 253-258 [PMID: 16945375 DOI: 10.1016/j.atherosclerosis.2006.07.014]
- 84 **Hsueh W**. Prostaglandin biosynthesis in pulmonary macrophages. *Am J Pathol* 1979; **97**: 137-148 [PMID: 495692]
- 85 **Decker K**. Biologically active products of stimulated liver macrophages (Kupffer cells). *Eur J Biochem* 1990; **192**: 245-261 [PMID: 2170121 DOI: 10.1111/j.1432-1033.1990.tb19222.x]
- 86 **Kuiper J**, Zijlstra FJ, Kamps JA, van Berkel TJ. Identification of prostaglandin D2 as the major eicosanoid from liver endothelial and Kupffer cells. *Biochim Biophys Acta* 1988; **959**: 143-152 [PMID: 3126817]
- 87 **Maciejewski-Lenoir D**, Richman JG, Hakak Y, Gaidarov I, Behan DP, Connolly DT. Langerhans cells release prostaglandin D2 in response to nicotinic acid. *J Invest Dermatol* 2006; **126**: 2637-2646 [PMID: 17008871 DOI: 10.1038/sj.jid.5700586]
- 88 **Ali M**, Cerskus AL, Zamecnik J, McDonald JW. Synthesis of prostaglandin D2 and thromboxane B2 by human platelets. *Thromb Res* 1977; **11**: 485-496 [PMID: 918907 DOI: 10.1016/0049-3848(77)90202-X]
- 89 **Tanaka K**, Ogawa K, Sugamura K, Nakamura M, Takano S, Nagata K. Cutting edge: differential production of prostaglandin D2 by human helper T cell subsets. *J Immunol* 2000; **164**: 2277-2280 [PMID: 10679060]
- 90 **Gallant MA**, Samadifam R, Hackett JA, Antoniou J, Parent JL, de Brum-Fernandes AJ. Production of prostaglandin D(2) by human osteoblasts and modulation of osteoprotegerin, RANKL, and cellular migration by DP and CRTH2 receptors. *J Bone Miner Res* 2005; **20**: 672-681 [PMID: 15765187 DOI: 10.1359/JBMR.041211]
- 91 **Jowsey IR**, Murdock PR, Moore GB, Murphy GJ, Smith SA, Hayes JD. Prostaglandin D2 synthase enzymes and PPAR-gamma are co-expressed in mouse 3T3-L1 adipocytes and human tissues. *Prostaglandins Other Lipid Mediat* 2003; **70**: 267-284 [PMID: 12611492]
- 92 **Bochenek G**, Nizankowska E, Gielicz A, Swierczyńska M, Szczeklik A. Plasma 9alpha,11beta-PGF2, a PGD2 metabolite, as a sensitive marker of mast cell activation by allergen in bronchial asthma. *Thorax* 2004; **59**: 459-464 [PMID: 15170023 DOI: 10.1136/thx.2003.013573]
- 93 **Dahlén SE**, Kumlin M. Monitoring mast cell activation by prostaglandin D2 in vivo. *Thorax* 2004; **59**: 453-455 [PMID: 15170020 DOI: 10.1136/thx.2004.026641]
- 94 **Morrow JD**, Guzzo C, Lazarus G, Oates JA, Roberts LJ. Improved diagnosis of mastocytosis by measurement of the major urinary metabolite of prostaglandin D2. *J Invest Dermatol* 1995; **104**: 937-940 [PMID: 7769262]
- 95 **Suzuki F**, Hayashi H, Hayaishi O. Transport of prostaglandin D2 into brain. *Brain Res* 1986; **385**: 321-328 [PMID: 3465420 DOI: 10.1016/0006-8993(86)91079-6]
- 96 **MacIouf J**, Corvazier E, Wang ZY. Development of a radioimmunoassay for prostaglandin D2 using an antiserum against 11-methoxime prostaglandin D2. *Prostaglandins* 1986; **31**: 123-132 [PMID: 3456623]
- 97 **Schuligoi R**, Schmidt R, Geisslinger G, Kollroser M, Peskar BA, Heinemann A. PGD2 metabolism in plasma: kinetics and relationship with bioactivity on DP1 and CRTH2 receptors. *Biochem Pharmacol* 2007; **74**: 107-117 [PMID: 17452035 DOI: 10.1016/j.bcp.2007.03.023]
- 98 **Haberl C**, Hültner L, Flügel A, Falk M, Geuenich S, Wilmanns W, Denzlinger C. Release of prostaglandin D2 by murine mast cells: importance of metabolite formation for antiproliferative activity. *Mediators Inflamm* 1998; **7**: 79-84 [PMID: 9836493 DOI: 10.1080/09629359891216]
- 99 **Zhang J**, Gong Y, Yu Y. PG F(2α) Receptor: A Promising Therapeutic Target for Cardiovascular Disease. *Front Pharmacol* 2010; **1**: 116 [PMID: 21607067 DOI: 10.3389/fphar.2010.00116]
- 100 **Liston TE**, Roberts LJ. Metabolic fate of radiolabeled prostaglandin D2 in a normal human male volunteer. *J Biol Chem* 1985; **260**: 13172-13180 [PMID: 3863815]
- 101 **Akiyama M**, Watanabe Y, Nishikawa T. Immunohistochemical characterization of human cutaneous mast cells in urticaria pigmentosa (cutaneous mastocytosis). *Acta Pathol Jpn* 1991; **41**: 344-349 [PMID: 1651041]
- 102 **Freeman JG**, Ryan JJ, Shelburne CP, Bailey DP, Bouton LA, Narasimhachari N, Domen J, Siméon N, Couderc F, Stewart JK. Catecholamines in murine bone marrow derived mast cells. *J Neuroimmunol* 2001; **119**: 231-238 [PMID: 11585626 DOI: 10.1016/S0165-5728(01)00384-8]
- 103 **Azzolina A**, Bongiovanni A, Lampiasi N. Substance P induces TNF-alpha and IL-6 production through NF kappa B in peritoneal mast cells. *Biochim Biophys Acta* 2003; **1643**: 75-83 [PMID: 14654230 DOI: 10.1016/j.bbamer.2003.09.003]

- 104 **Gordon JR**, Galli SJ. Mast cells as a source of both preformed and immunologically inducible TNF-alpha/cachectin. *Nature* 1990; **346**: 274-276 [PMID: 2374592 DOI: 10.1038/346274a0]
- 105 **Echtenacher B**, Männel DN, Hültner L. Critical protective role of mast cells in a model of acute septic peritonitis. *Nature* 1996; **381**: 75-77 [PMID: 8609992 DOI: 10.1038/381075a0]
- 106 **Kaartinen M**, Penttilä A, Kovanen PT. Mast cells in rupture-prone areas of human coronary atheromas produce and store TNF-alpha. *Circulation* 1996; **94**: 2787-2792 [PMID: 8941103 DOI: 10.1161/01.CIR.94.11.2787]
- 107 **Suto H**, Nakae S, Kakurai M, Sedgwick JD, Tsai M, Galli SJ. Mast cell-associated TNF promotes dendritic cell migration. *J Immunol* 2006; **176**: 4102-4112 [PMID: 16547246]
- 108 **Nakae S**, Suto H, Kakurai M, Sedgwick JD, Tsai M, Galli SJ. Mast cells enhance T cell activation: Importance of mast cell-derived TNF. *Proc Natl Acad Sci USA* 2005; **102**: 6467-6472 [PMID: 15840716 DOI: 10.1073/pnas.0501912102]
- 109 **Bradding P**, Mediawake R, Feather IH, Madden J, Church MK, Holgate ST, Howarth PH. TNF alpha is localized to nasal mucosal mast cells and is released in acute allergic rhinitis. *Clin Exp Allergy* 1995; **25**: 406-415 [PMID: 7553243]
- 110 **Walsh LJ**, Trinchieri G, Waldorf HA, Whitaker D, Murphy GF. Human dermal mast cells contain and release tumor necrosis factor alpha, which induces endothelial leukocyte adhesion molecule 1. *Proc Natl Acad Sci USA* 1991; **88**: 4220-4224 [PMID: 1709737 DOI: 10.1073/pnas.88.10.4220]
- 111 **Nakae S**, Ho LH, Yu M, Monteforte R, Iikura M, Suto H, Galli SJ. Mast cell-derived TNF contributes to airway hyper-reactivity, inflammation, and TH2 cytokine production in an asthma model in mice. *J Allergy Clin Immunol* 2007; **120**: 48-55 [PMID: 17482668 DOI: 10.1016/j.jaci.2007.02.046]
- 112 **Sampson AP**, Castling DP, Green CP, Price JF. Persistent increase in plasma and urinary leukotrienes after acute asthma. *Arch Dis Child* 1995; **73**: 221-225 [PMID: 7492159]
- 113 **Brightling CE**, Bradding P. The re-emergence of the mast cell as a pivotal cell in asthma pathogenesis. *Curr Allergy Asthma Rep* 2005; **5**: 130-135 [PMID: 15683613 DOI: 10.1007/s11882-005-0086-9]
- 114 **Tanaka S**, Tanaka H, Abe S. High dose of inhaled fluticasone reduces high levels of urinary leukotriene E4 in the early morning in mild and moderate nocturnal asthma. *Chest* 2003; **124**: 1768-1773 [PMID: 14605047 DOI: 10.1378/chest.124.5.1768]
- 115 **Leukotriene B4**. Human metabolome database, 2013, May 29. Available from: URL: <http://www.hmdb.ca/metabolites/HMDB01085>
- 116 **Longley J**, Duffy TP, Kohn S. The mast cell and mast cell disease. *J Am Acad Dermatol* 1995; **32**: 545-651; quiz 562-564 [PMID: 7896943 DOI: 10.1016/0190-9622(95)90336-4]
- 117 **Rosbotham JL**, Malik NM, Syrris P, Jeffery S, Bedlow A, Gharraie S, Murday VA, Holden CA, Carter ND. Lack of c-kit mutation in familial urticaria pigmentosa. *Br J Dermatol* 1999; **140**: 849-852 [PMID: 10354021 DOI: 10.1046/j.1365-2133.1999.02814.x]
- 118 **Lim KH**, Tefferi A, Lasho TL, Finke C, Patnaik M, Butterfield JH, McClure RF, Li CY, Pardanani A. Systemic mastocytosis in 342 consecutive adults: survival studies and prognostic factors. *Blood* 2009; **113**: 5727-5736 [PMID: 19363219 DOI: 10.1182/blood-2009-02-205237]
- 119 **Molderings GJ**, Kolck U, Scheurlen C, Brüss M, Frieling T, Raithe M, Homann J. Systemic mast cell disease with gastrointestinal symptoms--a diagnostic questionnaire. *Dtsch Med Wochenschr* 2006; **131**: 2095-2100 [PMID: 16981082 DOI: 10.1055/s-2006-951337]

**P- Reviewers:** Fozza C, Imashuku S, Takahashi M

**S- Editor:** Ma YJ **L- Editor:** A **E- Editor:** Wu HL





## INSTRUCTIONS TO AUTHORS

### GENERAL INFORMATION

*World Journal of Hematology* (*World J Hematol*, *WJH*, online ISSN 2218-6204, DOI: 10.5315) is a peer-reviewed open access (OA) academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

#### Aim and scope

*WJH* covers topics concerning experimental, clinical, oncological and transplant hematology, transfusion science, hemostasis and thrombosis, evidence-based medicine, epidemiology and nursing. The current columns of *WJH* include editorial, frontier, diagnostic advances, therapeutics advances, field of vision, mini-reviews, review, topic highlight, medical ethics, original articles, case report, clinical case conference (Clinicopathological conference), and autobiography. Priority publication will be given to articles concerning diagnosis and treatment of hematological diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJH*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

*WJH* is edited and published by Baishideng Publishing Group (BPG). BPG has a strong professional editorial team composed of science editors, language editors and electronic editors. BPG currently publishes 42 OA clinical medical journals, including 41 in English, has a total of 15 471 editorial board members or peer reviewers, and is a world first-class publisher.

#### Columns

The columns in the issues of *WJH* will include: (1) Editorial: The editorial board members are invited to make comments on an important topic in their field in terms of its current research status and future directions to lead the development of this discipline; (2) Frontier: The editorial board members are invited to select a highly cited cutting-edge original paper of his/her own to summarize major findings, the problems that have been resolved and remain to be resolved, and future research directions to help readers understand his/her important academic point of view and future research directions in the field; (3) Diagnostic Advances: The editorial board members are invited to write high-quality diagnostic advances in their field to improve the diagnostic skills of readers. The topic covers general clinical diagnosis, differential diagnosis, pathological diagnosis, laboratory diagnosis, imaging diagnosis, endoscopic diagnosis, biotechnological diagnosis, functional diagnosis, and physical diagnosis; (4) Therapeutics Advances: The editorial board members are invited to write high-quality therapeutic advances in their field to help improve the therapeutic skills of readers. The topic covers medication therapy, psychotherapy, physical therapy, replacement therapy, interventional therapy, minimally invasive therapy, endoscopic therapy, transplantation therapy, and surgical therapy; (5) Field of Vision: The editorial board members are invited to write commentaries on classic articles, hot topic articles, or latest articles to keep readers at the forefront of research and increase their levels of clinical research. Classic articles refer to papers that are included

in Web of Knowledge and have received a large number of citations (ranking in the top 1%) after being published for more than years, reflecting the quality and impact of papers. Hot topic articles refer to papers that are included in Web of Knowledge and have received a large number of citations after being published for no more than 2 years, reflecting cutting-edge trends in scientific research. Latest articles refer to the latest published high-quality papers that are included in PubMed, reflecting the latest research trends. These commentary articles should focus on the status quo of research, the most important research topics, the problems that have now been resolved and remain to be resolved, and future research directions. Basic information about the article to be commented (including authors, article title, journal name, year, volume, and inclusive page numbers); (6) Minireviews: The editorial board members are invited to write short reviews on recent advances and trends in research of molecular biology, genomics, and related cutting-edge technologies to provide readers with the latest knowledge and help improve their diagnostic and therapeutic skills; (7) Review: To make a systematic review to focus on the status quo of research, the most important research topics, the problems that have now been resolved and remain to be resolved, and future research directions; (8) Topic Highlight: The editorial board members are invited to write a series of articles (7-10 articles) to comment and discuss a hot topic to help improve the diagnostic and therapeutic skills of readers; (9) Medical Ethics: The editorial board members are invited to write articles about medical ethics to increase readers' knowledge of medical ethics. The topic covers international ethics guidelines, animal studies, clinical trials, organ transplantation, etc.; (10) Clinical Case Conference or Clinicopathological Conference: The editorial board members are invited to contribute high-quality clinical case conference; (11) Original Articles: To report innovative and original findings in hematology; (12) Brief Articles: To briefly report the novel and innovative findings in hematology; (13) Meta-Analysis: To evaluate the clinical effectiveness in hematology by using data from two or more randomised control trials; (14) Case Report: To report a rare or typical case; (15) Letters to the Editor: To discuss and make reply to the contributions published in *WJH*, or to introduce and comment on a controversial issue of general interest; (16) Book Reviews: To introduce and comment on quality monographs of hematology; and (17) Autobiography: The editorial board members are invited to write their autobiography to provide readers with stories of success or failure in their scientific research career. The topic covers their basic personal information and information about when they started doing research work, where and how they did research work, what they have achieved, and their lessons from success or failure.

#### Name of journal

*World Journal of Hematology*

#### ISSN

ISSN 2218-6204 (online)

#### Launch date

June 6, 2012

#### Frequency

Quarterly

#### Editor-in-Chief

Xiaoyan Jiang, MD, PhD, Associate Professor, Medical Gen-

## Instructions to authors

etics, University of British Columbia, Terry Fox Laboratory, British Columbia Cancer Agency, 675 West 10th Ave, Vancouver, BC, V5Z 1L3, Canada

**Thomas J Kipps, MD, PhD, Professor** of Medicine, University of California, San Diego, Moores Cancer Center, 3855 Health Sciences Drive, MC 0820, La Jolla, CA 92093-0820, United States

### Editorial office

Jin-Lei Wang, Director

Xiu-Xia Song, Vice Director

*World Journal of Hematology*

Room 903, Building D, Ocean International Center,

No. 62 Dongsihuan Zhonglu, Chaoyang District,

Beijing 100025, China

Telephone: +86-10-59080039

Fax: +86-10-85381893

E-mail: [wjhematol@wjnet.com](mailto:wjhematol@wjnet.com)

<http://www.wjnet.com>

### Publisher

Baishideng Publishing Group Co., Limited

Flat C, 23/F, Lucky Plaza, 315-321 Lockhart Road,

Wan Chai, Hong Kong, China

Telephone: +852-58042046

Fax: +852-31158812

E-mail: [bpgoffice@wjnet.com](mailto:bpgoffice@wjnet.com)

<http://www.wjnet.com>

### Production center

Beijing Baishideng BioMed Scientific Co., Limited

Room 903, Building D, Ocean International Center,

No. 62 Dongsihuan Zhonglu, Chaoyang District,

Beijing 100025, China

Telephone: +86-10-85381892

Fax: +86-10-85381893

### Representative office

USA Office

8226 Regency Drive,

Pleasanton, CA 94588-3144, United States

### Instructions to authors

Full instructions are available online at [http://www.wjnet.com/2218-6204/g\\_info\\_20100722173604.htm](http://www.wjnet.com/2218-6204/g_info_20100722173604.htm).

### Indexed and Abstracted in

Digital Object Identifier.

## SPECIAL STATEMENT

All articles published in this journal represent the viewpoints of the authors except where indicated otherwise.

### Biostatistical editing

Statistical review is performed after peer review. We invite an expert in Biomedical Statistics to evaluate the statistical method used in the paper, including *t*-test (group or paired comparisons), chi-squared test, Redit, probit, logit, regression (linear, curvilinear, or stepwise), correlation, analysis of variance, analysis of covariance, *etc.* The reviewing points include: (1) Statistical methods should be described when they are used to verify the results; (2) Whether the statistical techniques are suitable or correct; (3) Only homogeneous data can be averaged. Standard deviations are preferred to standard errors. Give the number of observations and subjects (*n*). Losses in observations, such as drop-outs from the study should be reported; (4) Values such as ED50, LD50, IC50 should have their 95% confidence limits calculated and compared by weighted probit analysis (Bliss and Finney); and (5) The word 'significantly' should be replaced by its synonyms (if it indicates extent) or the *P* value (if it indicates statistical significance).

### Conflict-of-interest statement

In the interests of transparency and to help reviewers assess any potential bias, *WJH* requires authors of all papers to declare any competing commercial, personal, political, intellectual, or religious interests in relation to the submitted work. Referees are also asked to indicate any potential conflict they might have reviewing a particular paper. Before submitting, authors are suggested to read "Uniform Requirements for Manuscripts Submitted to Biomedical Journals: Ethical Considerations in the Conduct and Reporting of Research: Conflicts of Interest" from International Committee of Medical Journal Editors (ICMJE), which is available at: [http://www.icmje.org/ethical\\_4conflicts.html](http://www.icmje.org/ethical_4conflicts.html).

Sample wording: [Name of individual] has received fees for serving as a speaker, a consultant and an advisory board member for [names of organizations], and has received research funding from [names of organization]. [Name of individual] is an employee of [name of organization]. [Name of individual] owns stocks and shares in [name of organization]. [Name of individual] owns patent [patent identification and brief description].

### Statement of informed consent

Manuscripts should contain a statement to the effect that all human studies have been reviewed by the appropriate ethics committee or it should be stated clearly in the text that all persons gave their informed consent prior to their inclusion in the study. Details that might disclose the identity of the subjects under study should be omitted. Authors should also draw attention to the Code of Ethics of the World Medical Association (Declaration of Helsinki, 1964, as revised in 2004).

### Statement of human and animal rights

When reporting the results from experiments, authors should follow the highest standards and the trial should conform to Good Clinical Practice (for example, US Food and Drug Administration Good Clinical Practice in FDA-Regulated Clinical Trials; UK Medicines Research Council Guidelines for Good Clinical Practice in Clinical Trials) and/or the World Medical Association Declaration of Helsinki. Generally, we suggest authors follow the lead investigator's national standard. If doubt exists whether the research was conducted in accordance with the above standards, the authors must explain the rationale for their approach and demonstrate that the institutional review body explicitly approved the doubtful aspects of the study.

Before submitting, authors should make their study approved by the relevant research ethics committee or institutional review board. If human participants were involved, manuscripts must be accompanied by a statement that the experiments were undertaken with the understanding and appropriate informed consent of each. Any personal item or information will not be published without explicit consents from the involved patients. If experimental animals were used, the materials and methods (experimental procedures) section must clearly indicate that appropriate measures were taken to minimize pain or discomfort, and details of animal care should be provided.

## SUBMISSION OF MANUSCRIPTS

Manuscripts should be typed in 1.5 line spacing and 12 pt. Book Antiqua with ample margins. Number all pages consecutively, and start each of the following sections on a new page: Title Page, Abstract, Introduction, Materials and Methods, Results, Discussion, Acknowledgements, References, Tables, Figures, and Figure Legends. Neither the editors nor the publisher are responsible for the opinions expressed by contributors. Manuscripts formally accepted for publication become the permanent property of Baishideng Publishing Group Co., Limited, and may not be reproduced by any means, in whole or in part, without the written permission of both the authors and the publisher. We reserve the right to copyedit and put onto our website accepted manuscripts. Authors should follow the relevant guidelines for the care and use of laboratory animals of their institution or national animal welfare committee. For the sake of transparency in regard to the performance and reporting of clinical trials, we endorse the policy of the ICMJE to refuse to publish papers on clinical trial results if the trial was not recorded in a publicly-accessible registry at its outset. The only register now

available, to our knowledge, is <http://www.clinicaltrials.gov> sponsored by the United States National Library of Medicine and we encourage all potential contributors to register with it. However, in the case that other registers become available you will be duly notified. A letter of recommendation from each author's organization should be provided with the contributed article to ensure the privacy and secrecy of research is protected.

Authors should retain one copy of the text, tables, photographs and illustrations because rejected manuscripts will not be returned to the author(s) and the editors will not be responsible for loss or damage to photographs and illustrations sustained during mailing.

### Online submissions

Manuscripts should be submitted through the Online Submission System at: <http://www.wjnet.com/esps/>. Authors are highly recommended to consult the ONLINE INSTRUCTIONS TO AUTHORS ([http://www.wjnet.com/2218-6204/g\\_info\\_20100722173604.htm](http://www.wjnet.com/2218-6204/g_info_20100722173604.htm)) before attempting to submit online. For assistance, authors encountering problems with the Online Submission System may send an email describing the problem to [wjhematol@wjnet.com](mailto:wjhematol@wjnet.com), or by telephone: +86-10-85381892. If you submit your manuscript online, do not make a postal contribution. Repeated online submission for the same manuscript is strictly prohibited.

## MANUSCRIPT PREPARATION

All contributions should be written in English. All articles must be submitted using word-processing software. All submissions must be typed in 1.5 line spacing and 12 pt. Book Antiqua with ample margins. Style should conform to our house format. Required information for each of the manuscript sections is as follows:

### Title page

**Title:** Title should be less than 12 words.

**Running title:** A short running title of less than 6 words should be provided.

**Authorship:** Authorship credit should be in accordance with the standard proposed by ICMJE, based on (1) substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; (2) drafting the article or revising it critically for important intellectual content; and (3) final approval of the version to be published. Authors should meet conditions 1, 2, and 3.

**Institution:** Author names should be given first, then the complete name of institution, city, province and postcode. For example, Xu-Chen Zhang, Li-Xin Mei, Department of Pathology, Chengde Medical College, Chengde 067000, Hebei Province, China. One author may be represented from two institutions, for example, George Sgourakis, Department of General, Visceral, and Transplantation Surgery, Essen 45122, Germany; George Sgourakis, 2nd Surgical Department, Korgialenio-Benakio Red Cross Hospital, Athens 15451, Greece

**Author contributions:** The format of this section should be: Author contributions: Wang CL and Liang L contributed equally to this work; Wang CL, Liang L, Fu JF, Zou CC, Hong F and Wu XM designed the research; Wang CL, Zou CC, Hong F and Wu XM performed the research; Xue JZ and Lu JR contributed new reagents/analytic tools; Wang CL, Liang L and Fu JF analyzed the data; and Wang CL, Liang L and Fu JF wrote the paper.

**Supportive foundations:** The complete name and number of supportive foundations should be provided, e.g., Supported by National Natural Science Foundation of China, No. 30224801

**Correspondence to:** Only one corresponding address should be provided. Author names should be given first, then author title, affiliation, the complete name of institution, city, postcode, province, country, and email. All the letters in the email should be in lower case. A space interval should be inserted between country name and email address. For example, Montgomery Bissell, MD, Profes-

sor of Medicine, Chief, Liver Center, Gastroenterology Division, University of California, Box 0538, San Francisco, CA 94143, United States. [montgomery.bissell@ucsf.edu](mailto:montgomery.bissell@ucsf.edu)

**Telephone and fax:** Telephone and fax should consist of +, country number, district number and telephone or fax number, e.g., Telephone: +86-10-85381892 Fax: +86-10-85381893

**Peer reviewers:** All articles received are subject to peer review. Normally, three experts are invited for each article. Decision on acceptance is made only when at least two experts recommend publication of an article. All peer-reviewers are acknowledged on Express Submission and Peer-review System website.

### Abstract

There are unstructured abstracts (no less than 200 words) and structured abstracts. The specific requirements for structured abstracts are as follows:

An informative, structured abstract should accompany each manuscript. Abstracts of original contributions should be structured into the following sections: AIM (no more than 20 words; Only the purpose of the study should be included. Please write the Aim in the form of "To investigate/study/..."), METHODS (no less than 140 words for Original Articles; and no less than 80 words for Brief Articles), RESULTS (no less than 150 words for Original Articles and no less than 120 words for Brief Articles; You should present *P* values where appropriate and must provide relevant data to illustrate how they were obtained, e.g.,  $6.92 \pm 3.86$  vs  $3.61 \pm 1.67$ ,  $P < 0.001$ ), and CONCLUSION (no more than 26 words).

### Key words

Please list 5-10 key words, selected mainly from *Index Medicus*, which reflect the content of the study.

### Core tip

Please write a summary of less than 100 words to outline the most innovative and important arguments and core contents in your paper to attract readers.

### Text

For articles of these sections, original articles and brief articles, the main text should be structured into the following sections: INTRODUCTION, MATERIALS AND METHODS, RESULTS and DISCUSSION, and should include appropriate Figures and Tables. Data should be presented in the main text or in Figures and Tables, but not in both.

### Illustrations

Figures should be numbered as 1, 2, 3, etc., and mentioned clearly in the main text. Provide a brief title for each figure on a separate page. Detailed legends should not be provided under the figures. This part should be added into the text where the figures are applicable. Keeping all elements compiled is necessary in line-art image. Scale bars should be used rather than magnification factors, with the length of the bar defined in the legend rather than on the bar itself. File names should identify the figure and panel. Avoid layering type directly over shaded or textured areas. Please use uniform legends for the same subjects. For example: Figure 1 Pathological changes in atrophic gastritis after treatment. A: ...; B: ...; C: ...; D: ...; E: ...; F: ...; G: ...etc. It is our principle to publish high resolution-figures for the E-versions.

### Tables

Three-line tables should be numbered 1, 2, 3, etc., and mentioned clearly in the main text. Provide a brief title for each table. Detailed legends should not be included under tables, but rather added into the text where applicable. The information should complement, but not duplicate the text. Use one horizontal line under the title, a second under column heads, and a third below the Table, above any footnotes. Vertical and italic lines should be omitted.

### Notes in tables and illustrations

Data that are not statistically significant should not be noted. \**P* <

## Instructions to authors

0.05, <sup>b</sup> $P < 0.01$  should be noted ( $P > 0.05$  should not be noted). If there are other series of  $P$  values, <sup>c</sup> $P < 0.05$  and <sup>d</sup> $P < 0.01$  are used. A third series of  $P$  values can be expressed as <sup>e</sup> $P < 0.05$  and <sup>f</sup> $P < 0.01$ . Other notes in tables or under illustrations should be expressed as <sup>1</sup>F, <sup>2</sup>F, <sup>3</sup>F; or sometimes as other symbols with a superscript (Arabic numerals) in the upper left corner. In a multi-curve illustration, each curve should be labeled with ●, ○, ■, □, ▲, △, etc., in a certain sequence.

### Acknowledgments

Brief acknowledgments of persons who have made genuine contributions to the manuscript and who endorse the data and conclusions should be included. Authors are responsible for obtaining written permission to use any copyrighted text and/or illustrations.

## REFERENCES

### Coding system

The author should number the references in Arabic numerals according to the citation order in the text. Put reference numbers in square brackets in superscript at the end of citation content or after the cited author's name. For citation content which is part of the narration, the coding number and square brackets should be typeset normally. For example, "Crohn's disease (CD) is associated with increased intestinal permeability<sup>[1,2]</sup>". If references are cited directly in the text, they should be put together within the text, for example, "From references<sup>[19,22-24]</sup>, we know that..."

When the authors write the references, please ensure that the order in text is the same as in the references section, and also ensure the spelling accuracy of the first author's name. Do not list the same citation twice.

### PMID and DOI

Please provide PubMed citation numbers to the reference list, e.g., PMID and DOI, which can be found at <http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed> and <http://www.crossref.org/SimpleTextQuery/>, respectively. The numbers will be used in E-version of this journal.

### Style for journal references

Authors: the name of the first author should be typed in bold-faced letters. The family name of all authors should be typed with the initial letter capitalized, followed by their abbreviated first and middle initials. (For example, Lian-Sheng Ma is abbreviated as Ma LS, Bo-Rong Pan as Pan BR). The title of the cited article and italicized journal title (journal title should be in its abbreviated form as shown in PubMed), publication date, volume number (in black), start page, and end page [PMID: 11819634 DOI: 10.3748/wjg.13.5396].

### Style for book references

Authors: the name of the first author should be typed in bold-faced letters. The surname of all authors should be typed with the initial letter capitalized, followed by their abbreviated middle and first initials. (For example, Lian-Sheng Ma is abbreviated as Ma LS, Bo-Rong Pan as Pan BR) Book title. Publication number. Publication place: Publication press, Year: start page and end page.

### Format

#### Journals

English journal article (list all authors and include the PMID where applicable)

- 1 **Jung EM**, Clevert DA, Schreyer AG, Schmitt S, Rennert J, Kubale R, Feuerbach S, Jung F. Evaluation of quantitative contrast harmonic imaging to assess malignancy of liver tumors: A prospective controlled two-center study. *World J Gastroenterol* 2007; **13**: 6356-6364 [PMID: 18081224 DOI: 10.3748/wjg.13.6356]

Chinese journal article (list all authors and include the PMID where applicable)

- 2 **Lin GZ**, Wang XZ, Wang P, Lin J, Yang FD. Immunologic effect of Jianpi Yishen decoction in treatment of Pixu-diarhoea. *Shijie Huaren Xiaohua Zazhi* 1999; **7**: 285-287

In press

- 3 **Tian D**, Araki H, Stahl E, Bergelson J, Kreitman M. Signature

of balancing selection in Arabidopsis. *Proc Natl Acad Sci USA* 2006; In press

Organization as author

- 4 **Diabetes Prevention Program Research Group**. Hypertension, insulin, and proinsulin in participants with impaired glucose tolerance. *Hypertension* 2002; **40**: 679-686 [PMID: 12411462 PMID:2516377 DOI:10.1161/01.HYP.0000035706.28494.09]

Both personal authors and an organization as author

- 5 **Vallancien G**, Emberton M, Harving N, van Moorselaar RJ; Alf-One Study Group. Sexual dysfunction in 1, 274 European men suffering from lower urinary tract symptoms. *J Urol* 2003; **169**: 2257-2261 [PMID: 12771764 DOI:10.1097/01.ju.0000067940.76090.73]

No author given

- 6 21st century heart solution may have a sting in the tail. *BMJ* 2002; **325**: 184 [PMID: 12142303 DOI:10.1136/bmj.325.7357.184]

Volume with supplement

- 7 **Geraud G**, Spierings EL, Keywood C. Tolerability and safety of frovatriptan with short- and long-term use for treatment of migraine and in comparison with sumatriptan. *Headache* 2002; **42** Suppl 2: S93-99 [PMID: 12028325 DOI:10.1046/j.1526-4610.42.s2.7.x]

Issue with no volume

- 8 **Banit DM**, Kaufer H, Hartford JM. Intraoperative frozen section analysis in revision total joint arthroplasty. *Clin Orthop Relat Res* 2002; (**401**): 230-238 [PMID: 12151900 DOI:10.1097/00003086-200208000-00026]

No volume or issue

- 9 Outreach: Bringing HIV-positive individuals into care. *HRS-A Careaction* 2002; 1-6 [PMID: 12154804]

### Books

Personal author(s)

- 10 **Sherlock S**, Dooley J. Diseases of the liver and biliary system. 9th ed. Oxford: Blackwell Sci Pub, 1993: 258-296

Chapter in a book (list all authors)

- 11 **Lam SK**. Academic investigator's perspectives of medical treatment for peptic ulcer. In: Swabb EA, Azabo S. Ulcer disease: investigation and basis for therapy. New York: Marcel Dekker, 1991: 431-450

Author(s) and editor(s)

- 12 **Breedlove GK**, Schorffheide AM. Adolescent pregnancy. 2nd ed. Wiczorek RR, editor. White Plains (NY): March of Dimes Education Services, 2001: 20-34

Conference proceedings

- 13 **Harnden P**, Joffe JK, Jones WG, editors. Germ cell tumours V. Proceedings of the 5th Germ cell tumours Conference; 2001 Sep 13-15; Leeds, UK. New York: Springer, 2002: 30-56

Conference paper

- 14 **Christensen S**, Oppacher F. An analysis of Koza's computational effort statistic for genetic programming. In: Foster JA, Lutton E, Miller J, Ryan C, Tettamanzi AG, editors. Genetic programming. EuroGP 2002: Proceedings of the 5th European Conference on Genetic Programming; 2002 Apr 3-5; Kinsdale, Ireland. Berlin: Springer, 2002: 182-191

Electronic journal (list all authors)

- 15 Morse SS. Factors in the emergence of infectious diseases. *Emerg Infect Dis* serial online, 1995-01-03, cited 1996-06-05; 1(1): 24 screens. Available from: URL: <http://www.cdc.gov/ncidod/eid/index.htm>

Patent (list all authors)

- 16 **Pagedas AC**, inventor; Ancel Surgical R&D Inc., assignee. Flexible endoscopic grasping and cutting device and positioning tool assembly. United States patent US 20020103498. 2002 Aug 1

### Statistical data

Write as mean  $\pm$  SD or mean  $\pm$  SE.

### Statistical expression

Express *t* test as *t* (in italics), *F* test as *F* (in italics), chi square test as  $\chi^2$  (in Greek), related coefficient as *r* (in italics), degree of freedom as *ν* (in Greek), sample number as *n* (in italics), and probability as *P* (in italics).

### Units

Use SI units. For example: body mass, *m* (B) = 78 kg; blood pressure, *p* (B) = 16.2/12.3 kPa; incubation time, *t* (incubation) = 96 h; blood glucose concentration, *c* (glucose)  $6.4 \pm 2.1$  mmol/L; blood CEA mass concentration, *p* (CEA) = 8.6  $24.5 \mu\text{g/L}$ ; CO<sub>2</sub> volume fraction, 50 mL/L CO<sub>2</sub>, not 5% CO<sub>2</sub>; likewise for 40 g/L formaldehyde, not 10% formalin; and mass fraction, 8 ng/g, *etc.* Arabic numerals such as 23, 243, 641 should be read 23 243 641.

The format for how to accurately write common units and quantum numbers can be found at: [http://www.wjgnet.com/2218-6204/g\\_info\\_20100723213202.htm](http://www.wjgnet.com/2218-6204/g_info_20100723213202.htm).

### Abbreviations

Standard abbreviations should be defined in the abstract and on first mention in the text. In general, terms should not be abbreviated unless they are used repeatedly and the abbreviation is helpful to the reader. Permissible abbreviations are listed in Units, Symbols and Abbreviations: A Guide for Biological and Medical Editors and Authors (Ed. Baron DN, 1988) published by The Royal Society of Medicine, London. Certain commonly used abbreviations, such as DNA, RNA, HIV, LD50, PCR, HBV, ECG, WBC, RBC, CT, ESR, CSF, IgG, ELISA, PBS, ATP, EDTA, mAb, can be used directly without further explanation.

### Italics

Quantities: *t* time or temperature, *c* concentration, *A* area, *l* length, *m* mass, *V* volume.

Genotypes: *gyrA*, *arg 1*, *c myc*, *c fos*, *etc.*

Restriction enzymes: *EcoRI*, *HindI*, *BamHI*, *Kho I*, *Kpn I*, *etc.*

Biology: *H. pylori*, *E. coli*, *etc.*

### Examples for paper writing

All types of articles' writing style and requirement will be found in the link: <http://www.wjgnet.com/esps/NavigationInfo.aspx?id=15>

## RESUBMISSION OF THE REVISED MANUSCRIPTS

Authors must revise their manuscript carefully according to the revision policies of Baishideng Publishing Group Co., Limited. The revised version, along with the signed copyright transfer agreement, responses to the reviewers, and English language Grade A certifi-

cate (for non-native speakers of English), should be submitted to the online system *via* the link contained in the e-mail sent by the editor. If you have any questions about the revision, please send e-mail to [esps@wjgnet.com](mailto:esps@wjgnet.com).

### Language evaluation

The language of a manuscript will be graded before it is sent for revision. (1) Grade A: priority publishing; (2) Grade B: minor language polishing; (3) Grade C: a great deal of language polishing needed; and (4) Grade D: rejected. Revised articles should reach Grade A.

### Copyright assignment form

Please download a Copyright assignment form from [http://www.wjgnet.com/2218-6204/g\\_info\\_20100723213049.htm](http://www.wjgnet.com/2218-6204/g_info_20100723213049.htm).

### Responses to reviewers

Please revise your article according to the comments/suggestions provided by the reviewers. The format for responses to the reviewers' comments can be found at: [http://www.wjgnet.com/2218-6204/g\\_info\\_20100723212803.htm](http://www.wjgnet.com/2218-6204/g_info_20100723212803.htm).

### Proof of financial support

For papers supported by a foundation, authors should provide a copy of the approval document and serial number of the foundation.

### Links to documents related to the manuscript

WJH will be initiating a platform to promote dynamic interactions between the editors, peer reviewers, readers and authors. After a manuscript is published online, links to the PDF version of the submitted manuscript, the peer-reviewers' report and the revised manuscript will be put on-line. Readers can make comments on the peer reviewer's report, authors' responses to peer reviewers, and the revised manuscript. We hope that authors will benefit from this feedback and be able to revise the manuscript accordingly in a timely manner.

### Publication fee

WJH is an international, peer-reviewed, OA online journal. Articles published by this journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium and format, provided the original work is properly cited. The use is non-commercial and is otherwise in compliance with the license. Authors of accepted articles must pay a publication fee. Publication fee: 600 USD per article. All invited articles are published free of charge.



百世登  
**Baishideng**®

Published by **Baishideng Publishing Group Co., Limited**

Flat C, 23/F., Lucky Plaza, 315-321 Lockhart Road,

Wan Chai, Hong Kong, China

Fax: +852-31158812

Telephone: +852-58042046

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

<http://www.wjgnet.com>

