

World Journal of *Neurology*

World J Neurol 2014 September 28; 4(3): 12-22





Editorial Board

2011-2015

The *World Journal of Neurology* Editorial Board consists of 324 members, representing a team of worldwide experts in neurology. They are from 38 countries, including Australia (7), Austria (3), Belgium (3), Brazil (4), Canada (7), China (26), Croatia (1), Cuba (1), Cyprus (1), Czech Republic (1), Denmark (2), Egypt (4), Finland (1), France (2), Germany (17), Greece (5), Hungary (3), India (8), Iran (1), Israel (4), Italy (35), Japan (16), Mexico (5), Morocco (1), New Zealand (2), Nigeria (1), Poland (1), Portugal (2), Saudi Arabia (2), Singapore (6), Slovakia (1), South Korea (3), Spain (13), Switzerland (1), Thailand (3), Turkey (7), United Kingdom (11), and United States (113).

EDITORS-IN-CHIEF

Felipe Fregni, *Boston*
Vincenzo Solfrizzi, *Bari*

GUEST EDITORIAL BOARD MEMBERS

Fang-Chia Chang, *Taipei*
Chun-Chuan Chen, *Taoyuan*
San-Yuan Huang, *Taipei*
Suh-Hang Hank Juo, *Kaohsiung City*
Hsien-Yuan Lane, *Taichung*
Ching-Po Lin, *Taipei*
Hung-Chuan Pan, *Taichung*
Bing-wen Soong, *Taipei*
Kuo-Chen Wei, *Taichung*
Sheng-Nan Wu, *Tainan*

MEMBERS OF THE EDITORIAL BOARD



Australia

Roy G Beran, *Sydney*
Lucette Cysique, *Sydney*
John Gardiner, *Sydney*
Manuel B Graeber, *Sydney*
Xu-Feng Huang, *Wollongong*
George Damion Mellick, *Brisbane*
Sarah Spencer, *Victoria*



Austria

Andreas Gruber, *Vienna*
Joerg Kraus, *Salzburg*
Gerlig Widmann, *Anichstr*



Belgium

Anna C Jansen, *Brussels*

Steve Majerus, *Liege*
Rik RC Vandenberghe, *Leuven*



Brazil

Monica L Andersen, *São Paulo*
Elisa Brietzke, *São Paulo*
Paulo A Lotufo, *São Paulo*
Marcelo Rodrigues Masruha, *São Paulo*



Canada

Dave Ellemberg, *Montreal*
Shirley Fecteau, *Quebec*
Adel Gabriel, *Calgary*
Bernard Le Foll, *Toronto*
Laura-Ann Petitto, *Scarborough*
Mubeen Rafay, *Winnipeg*
José M Trigo, *Toronto*



China

Hui-Sheng Chen, *ShenYang*
Mei-Chun Cheung, *Hong Kong*
Wong Kwok Chu, *Hong Kong*
Wan Feng, *Wuhan*
Zhang Hao, *Shanghai*
De-Wen Hu, *Changsha*
Vincent Lai, *Hong Kong*
Xiao-Li Li, *Beijing*
Jian-Min Liu, *Shanghai*
Jun Liu, *Guangzhou*
Xiang-Dong Tang, *Chengdu*
Peng-Fei Yang, *Shanghai*
Xiang Zhang, *Xi'an*
Yan Zhang, *Beijing*
Zhi-Ren Zhang, *Chongqing*
Jin-Xia Zhu, *Beijing*



Croatia

Nela Pivac, *Zagreb*



Cuba

Elena R Cuspineda Bravo, *Habana*



Cyprus

Kyproula Christodoulou, *Nicosia*



Czech Republic

Robert Mikulik, *Brno*



Denmark

Hong-You Ge, *Aalborg*
Peter Johannsen, *Copenhagen*



Egypt

Sherifa Ahmad Hamed, *Dayton*
Taha M Mahmoud, *Zagazig*
Ahmed A K Abdel Razek, *Mansoura*
Sahar N Saleem, *Cairo*



Finland

Seppo Antero Kahkonen, *Helsinki*

**France**

Julien Dauguet, *Orsay*
Cyril Goudet, *Cardonville*

**Germany**

Boldizsar Czeh, *Munich*
Hans-Peter Hartung, *Dusseldorf*
Jens Haueisen, *Ilmenau*
Dirk M Hermann, *Essen*
Raimund Kleiser, *Linz*
Ulrich Muller, *Giessen*
Walter Paulus, *Göttingen*
Nikolaus Plesnila, *Munich*
Marc Röllinghoff, *Landsberg*
Jens Schittenhelm, *Tuebingen*
Jens Schmidt, *Göttingen*
Rudiger Seitz, *Dusseldorf*
Thomas Straube, *Jena*
Philipp A Thomann, *Heidelberg*
Johannes U V Thome, *Rostock*
Marcus Michael Unger, *Marburg*
Wolfgang Wick, *Heidelberg*

**Greece**

Constantoyannis Constantine, *Patras*
Kostas n Fountas, *Larissa*
Savas Grigoriadis, *Thessaloniki*
Ioannis N Mavridis, *Athens*
George Ntaios, *Larissa*

**Hungary**

Daniel Bereczki, *Budapest*
Xenia Gonda, *Budapest*
Norbert Kovacs, *Pecs*

**India**

Ravindra Kumar Garg, *Lucknow*
Ravi Gupta, *Dehradun*
Birendra Nath Mallick, *New Delhi*
Balraj Mittal, *Lucknow*
Hitesh N Modi, *Ahmedabad*
Mona Ragothaman, *Bangalore*
TN Sathyaprabha, *Bangalore*
Shirley Anne Telles, *Haridwar*

**Iran**

Seyyed Amirhossein Fazeli, *Gorgan*

**Israel**

Dimitrios M Karussis, *Jerusalem*
Asher Ornoy, *Hebrew*
Abraham Weizman, *Tikva*
Perla Werner, *Haifa*

**Italy**

Alberto Albanese, *Milan*
Claudia Altamura, *Rome*
Valeria Barresi, *Messina*
Emanuela Bartoccioni, *Rome*
Gabriella Bottini, *Milan*
Paolo Brambilla, *Udine*
Alfonso Cerase, *Siena*
Polezzi David, *Padova*
Luigi De Gennaro, *Rome*
Stefano Diciotti, *Florence*
Marina Fanin, *Padova*
Michele Ferrara, *L'Aquila*
Daniele Focosi, *Pisa*
Daniela Galimberti, *Milan*
Valentina Garibotto, *Geneva*
Marco Leonardi, *Bologna*
Maria Liguori, *Mangone*
Laura Mandelli, *Bologna*
Gianvito Martino, *Milan*
Giovanni Martinotti, *Rome*
Marianna Mazza, *Rome*
Antonio Orlacchio, *Rome*
Maurizio Paciaroni, *Perugia*
Marco Paoloni, *Rome*
Bernardo Perfetti, *Cheiti*
Alessandro Pezzini, *Brescia*
Gianfranco Puoti, *Naples*
Nilo Riva, *Milan*
Paolo Maria Rossini, *Rome*
Andrea Serino, *Bologna*
Gianfranco Spalletta, *Rome*
Fabrizio Stocchi, *Rome*
Pasquale Striano, *Genova*
Camillo Porcaro, *Rome*

**Japan**

Wataru Araki, *Tokyo*
Katsutoshi Furukawa, *Sendai*
Masafumi Ihara, *Kyoto*
Yuichi Inoue, *Tokyo*
Hiroshi Kadotani, *Kyoto*
Kazutaka Kobayashi, *Tokyo*
Yoshihiro Kokubo, *Suita*
Ryuichi Morishita, *Suita*
Tetsu Niwa, *Yokohama*
Katsunori Nonogaki, *Sendai*
Kunihiro Sakuma, *Aichi*
Katsuya Satoh, *Nagasaki*
Atsuyoshi Shimada, *Aichi*
Hiroshi Takahashi, *Tottori*
Naoyuki Takeuchi, *Sendai*
Kanato Yamagata, *Tokyo*

**Mexico**

Agnes Fleury, *Tlalpan*
Daniel San Juan, *Mexico City*
Julio Sotelo, *Mexico City*
Teresa Corona Vázquez, *Tlalpan*
Rodrigo Ramos Zuniga, *Guadalajara Jalisco*

**Morocco**

Samir Ahboucha, *Marrakesh*

**New Zealand**

Juan J Canales, *Christchurch*
Valery Feigin, *Auckland*

**Nigeria**

Mayowa O Owolabi, *Ibadan*

**Poland**

Pawel Piotr Liberski, *Lodz*

**Portugal**

Nuno Martins Marques Canas, *Lisbon*
Isaura Ferreira Tavares, *Porto*

**Saudi Arabia**

Ahmed S BaHammam, *Riyadh*
Essam A Elgamal, *Riyadh*

**Singapore**

Justin HG Dauwels, *Singapore*
Steven Graham, *Singapore*
Charanjit Kaur, *Singapore*
Vijay K Sharma, *Singapore*
Feng-Ru Tang, *Singapore*
Philip Lin Kiat Yap, *Singapore*

**Slovakia**

Peter Valkovič, *Bratislava*

**South Korea**

Ji Soo Kim, *Gyeonggi-do*
Myung Sik Lee, *Seoul*
Kyoungcho Suk, *Daegu*

**Spain**

Adria Arboix, *Barcelona*
Pedro Emilio Bermejo, *Madrid*
Ramon Cacabelos, *Coruna*
Isidro Ferrer, *Hospitalet de Llobregat*
Jesús A García-Sevilla, *Palma de Mallorca*
Bernardo Hontanilla, *Pamplona*
Esteban Pena Llamas, *Madrid*
Fernando Maestu, *Madrid*
Arturo Mangas, *Salamanca*
German Moris, *Gijon*
Jordi Perez-Tur, *Madrid*
Javier SCastresana, *Pamplona*
José M Trigo, *Toronto*

**Switzerland**

Marcel Arnold, *Bern*



Thailand

Suparerk Janjarasjitt, *Ubon Ratchathani*
Panitha Jindahra, *Bangkok*
Kittipan Rerkasem, *Chiang Mai*



Turkey

Ayşe Aralasmak, *Istanbul*
Isin Baral-Kulaksizoglu, *Istanbul*
Cengiz Cokluk, *Samsun*
Saygın Salih Eker, *Bursa*
Ateş Kadioglu, *Istanbul*
Suleyman Kaplan, *Samsun*
Rifat Nuri Sener, *Izmir*



United Kingdom

Zubair Ahmed, *Birmingham*
Chris John Bushe, *Basingstoke*
Andrea Eugenio Cavanna, *London*
White Gables, *Norwich*
Valentina Gallo, *London*
Sanjay Kumar, *Birmingham*
Tarik F Massoud, *Cambridge*
Mike Modo, *London*
Mario Alfredo Parra, *Edinburgh*
Annette Sterr, *Surrey*
Jan Stochl, *Cambridge*



United States

Herve Abdi, *Dallas*
Abhishek Agrawal, *New York*
Abass Alavi, *Philadelphia*
Quincy J Almeida, *Waterloo*
Alfredo Ardila, *Miami*
Carmel Armon, *Springfield*
Andrew D Barreto, *Houston*
Raymond T Bartus, *San Diego*
Archit Bhatt, *East Lansing*
Margit L Bleecker, *Baltimore*
Anna-Liisa Brownell, *Charlestown*
Ignazio Cali, *Cleveland*
Maren Carbon-Correll, *Manhasset*

Rudolph J Castellani, *Baltimore*
Carlos Cepeda, *Los Angeles*
Munmun Chattopadhyay, *Ann Arbor*
Rivka R Colen, *Boston*
James R Connor, *Hershey*
Li Cui, *Little Rock*
Yuri P Danilov, *Madison*
Mukeshwar Dhamala, *Atlanta*
David M Diamond, *Tampa*
David William Dodick, *Phoenix*
Richard I Doty, *Pennsylvania*
Christopher L Drake, *Detroit*
Timothy Q Duongthphd, *San Antonio*
Sherif M Elbasiouny, *Chicago*
Adam S Fleisher, *San Diego*
Robert Folmer, *Portland*
Majid Fotuhi, *Baltimore*
Dheeraj Gandhi, *Baltimore*
Yu-Lin Ge, *New York*
Hugo Geerts, *Berwyn*
Alexandros L Georgiadis, *Minneapolis*
Srikanth Givvmani, *Louisville*
Charles G Gross, *Princeton*
George T Grossberg, *Saint Louis*
Zhen He, *Jefferson*
Ming-Xiong Huang, *San Diego*
Sirous Jafarian, *North Vancouver*
Xiangning Jiang, *San Francisco*
Peter B Kang, *Boston*
Junghoon Kim, *Elkins Park*
David C Knight, *Birmingham*
Kennith F Layton, *Dallas*
Andrew G Lee, *Houston*
Walter S Lesley, *Temple*
David Sigmund Liebeskind, *Los Angeles*
Tianming Liu, *Athens*
Yahia M Lodi, *New York*
Edythe D London, *Los Angeles*
Jean-Pierre Louboutin, *Philadelphia*
Hanzhang Lu, *Dallas*
Kenneth Maiese, *Newark*
Silva Markovic-Plese, *Chapel Hill*
Marlon Stephen Mathews, *Orange*
Yousef Mohammad, *Chicago*
Amanda J Myers, *Miami*
Josef Novotny Jr, *Pittsburgh*
Arne M Nystuen, *Omaha*
Darin T Okuda, *Phoenix*
Wei-hong Pan, *Baton Rouge*
Juliann M Paolicchi, *Nashville*

Spyros Papapetropoulos, *Santa Ana*
Sergio Paradiso, *Iowa City*
Paul Park, *Ann Arbor*
Hemant A Parmar, *Ann Arbor*
George Perry, *San Antonio*
Marc Nicholas Potenza, *New Haven*
Yonglin Pu, *Chicago*
Haifa Qiao, *Tallahassee*
Liya Qiao, *Richmond*
Ralph Rahme, *Cincinnati*
Amit Ray, *New York*
Catherine M Roe, *Missouri*
Theodore H Schwartz, *New York*
Robert J Schwartzman, *Philadelphia*
Thomas F Scott, *Pittsburgh*
Souvik Sen, *Columbia*
KHEMA Sharma, *Miami*
Li Shen, *Indianapolis*
Gabriel A Silva, *La Jolla*
Elsayed Z Soliman, *Winston Salem*
Joshua Goh Oon Soo, *Baltimore*
Ashok Srinivasan, *Ann Arbor*
Massoud Stephane, *Minneapolis*
Shu-Wei Sun, *Loma Linda*
Emi Takahashi, *Boston*
Thomas Thannickal, *North Hills*
Timothy Adam Thrasher, *Houston*
Guochuan Emil Tsai, *Torrance*
Vassiliy Tsytarev, *Fairfax*
Tanya Nadine Turan, *Charleston*
Neetu Tyagi, *Louisville*
Denise A Valenti, *Boston*
Piero Verro, *Sacramento*
Marcela Votruba, *Cardiff*
Jian Wang, *Baltimore*
Kenneth L Weiss, *Jackson*
Harry T Whelan, *Milwaukee*
Keith D White, *Gainesville*
Peter Widdess-Walsh, *Livingston*
Zhongcong Xie, *Boston*
Midori Anne Yenari, *San Francisco*
Albert J Yoo, *Boston*
Robert J Young, *New York*
Brad Evan Zacharia, *New York*
T Thomas Zacharia, *Hershey*
Gabriel Zada, *Los Angeles*
Haoqian Zhang, *San Francisco*
Ming Zhang, *Philadelphia*
Yun Zhou, *Baltimore*

**REVIEW**

- 12 Cerebral ageing-the role of insulin and insulin-like growth factor signalling:
A review
Romain G, Opacka-Juffry J

Contents

World Journal of Neurology
Volume 4 Number 3 September 28, 2014

APPENDIX I-V Instructions to authors

ABOUT COVER Editorial Board Member of *World Journal of Neurology*, Paolo Maria Rossini, Professor, Department of Neurology, University campus Bio-Medico, Via Alvaro del Portillo, 21, 00128 Rome, Italy

AIM AND SCOPE

World Journal of Neurology (*World J Neurol*, *WJN*, online ISSN 2218-6212, DOI: 10.5316) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJN covers topics concerning neuro-oncology, electroneurophysiology, cerebrovascular diseases, epilepsy, cognitive impairment, myopathy and peripheral neuropathy, degenerative diseases, infectious diseases, demyelinating diseases, immunological diseases, genetic/metabolic diseases, affective disorders, headaches, sleep disorders, interventional neuroradiology, minimally invasive therapy, rehabilitation, diagnostic imaging, evidence-based medicine, epidemiology and nursing. Priority publication will be given to articles concerning diagnosis and treatment of neurological diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJN*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING *World Journal of Neurology* is now indexed in Digital Object Identifier.

FLYLEAF I-III Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Su-Qing Liu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*
Proofing Editorial Office Director: *Xin-Xia Song*

NAME OF JOURNAL
World Journal of Neurology

ISSN
ISSN 2218-6212 (online)

LAUNCH DATE
December 28, 2011

FREQUENCY
Quarterly

EDITORS-IN-CHIEF
Felipe Fregni, MD, PhD, MPH, Associate Professor, Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Harvard Medical School, 125 Nashua St. Room 725, Boston, MA 02114, United States

Vincenzo Solfrizzi, MD, PhD, Professor, Department of Internal Medicine, Immunology and Infectious Diseases, Section of Geriatric Medicine-Memory Unit,

University of Bari, Piazza G. Cesare, 11, 70124 Bari, Italy

EDITORIAL OFFICE
Jin-Lei Wang, Director
Xiu-Xia Song, Vice Director
World Journal of Neurology
Room 903, Building D, Ocean International Center,
No. 62 Dongsihuan Zhonglu, Chaoyang District,
Beijing 100025, China
Telephone: +86-10-59080039
Fax: +86-10-85381893
E-mail: editorialoffice@wjnet.com
Help Desk: <http://www.wjnet.com/esps/helpdesk.aspx>
<http://www.wjnet.com>

PUBLISHER
Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjnet.com
Help Desk: <http://www.wjnet.com/esps/helpdesk.aspx>
<http://www.wjnet.com>

PUBLICATION DATE
September 28, 2014

COPYRIGHT
© 2014 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at http://www.wjnet.com/2218-6212/g_info_20100722173918.htm

ONLINE SUBMISSION
<http://www.wjnet.com/esps/>

Cerebral ageing-the role of insulin and insulin-like growth factor signalling: A review

Georgia Romain, Jolanta Opacka-Juffry

Georgia Romain, Jolanta Opacka-Juffry, Department of Life Sciences, University of Roehampton, SW15 4JD London, United Kingdom

Author contributions: Romain G and Opacka-Juffry J both contributed to this paper.

Conflict-of-interest: No potential conflicts of interest relevant to this article were reported.

Open-Access: This article is an open-access article which selected by an in-house editor and fully peer-reviewed by external reviewers. It distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Jolanta Opacka-Juffry, Professor, Department of Life Sciences, University of Roehampton, SW15 4JD London, United Kingdom. j.opacka-juffry@roehampton.ac.uk
Telephone: +44-20-83923563

Fax: +44-20-83923527

Received: July 29, 2014

Peer-review started: July 29, 2014

First decision: August 14, 2014

Revised: August 27, 2014

Accepted: September 3, 2014

Article in press: September 4, 2014

Published online: September 28, 2014

Abstract

Cerebral ageing is a complex biological process associated with progressing cerebrovascular disease and neuronal death. It does not always, however, associate with a functional decline, as the ageing mammalian brain retains considerable functional plasticity which supports successful cerebral ageing where age-related cognitive decline is modest. On the contrary, pathological cerebral ageing results in memory impairment and cognitive deterioration, with Alzheimer's disease (AD) being a florid example. Trophic/growth factors promote brain plasticity; among them are peptides which belong to

the insulin family. Preclinical research suggests that the evolutionarily conserved brain insulin/insulin-like growth factor-1 (IGF-1) signalling system controls lifespan and protects against some features of AD such as neurodegeneration-related accumulation of toxic proteins and cognitive deficiencies, as observed in animal models. Insulin and IGF-1 activate cell signalling mechanisms which play protective and regenerative roles; abnormalities in the insulin/IGF-1 system may trigger a cascade of neurodegeneration in AD. AD patients show cerebral resistance to insulin which associates with IGF-I resistance and dysregulation of insulin/IGF-1 receptors as well as cognitive deterioration. This review is focused on the roles of the insulin/IGF-1 signalling system in cerebral ageing and its potential involvement in neurodegeneration in the human brain as seen against the background of preclinical evidence.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

Key words: Brain; Insulin; Insulin receptor; Insulin-like growth factor; Longevity; Alzheimer's; Diabetes mellitus; Inflammation

Core tip: Age itself is a major risk factor for the development of age-related cognitive decline, Alzheimer's and cerebrovascular diseases. Increased life expectancy necessitates the need to understand the processes that underlie successful vs pathological brain ageing in order to develop early interventions which may assist in delaying if not reversing the detrimental effects of brain ageing. This review focuses on the signalling system of insulin and insulin-like growth factor-1 (IIS) and its roles in cerebral ageing; it highlights some conflicting literature opinions and incomplete understandings of the roles and mechanisms of the IIS system.

Romain G, Opacka-Juffry J. Cerebral ageing-the role of insulin and insulin-like growth factor signalling: A review. *World J Neurol* 2014; 4(3): 12-22 Available from: URL: <http://www.wjnet.com>

INTRODUCTION

Ageing is a complex biological process that affects all species ranging from invertebrates through to non-human mammals and humans, being underpinned by alterations at molecular and cellular levels^[1,2] that compromise the organism's physiological homeostasis, induce susceptibility to disease and accelerate death^[3-5]. The cellular and anatomical changes that occur in the course of brain ageing contribute to cerebrovascular disease^[6], impact motor performance and learning and memory functions^[7,8], thus leading to cognitive decline and dementia^[9]. As human life span is growing and the proportion of the elderly is increasing in most societies, the prevalence of age-dependent diseases such as Alzheimer's disease (AD), Parkinson's disease and stroke is also on the increase world-wide. Consequently, understanding the biological and molecular mechanisms of what constitutes "successful" *vs* pathological cerebral ageing is critical to both delaying the ageing process, enhancing the quality of life in the elderly, and alleviating the already overburdened health care systems, from the cost component involved in treating age-related diseases.

In view of the above points, research on brain-ageing has generated a pool of information regarding the potential mechanisms, which underpin cognitive decline. Advancements in molecular and cell technology, have been instrumental in identifying factors such as oxidative stress^[10,11], epigenetic changes^[12], mitochondrial dysfunction, inflammatory response^[13], impaired cell signalling and gene expressions^[14], autophagy and protein turnover, target of rapamycin (TOR) and insulin/insulin-like growth factor (IGF) signalling as potential mechanisms that contribute to alterations observed in brain ageing^[9].

This review will first discuss what constitutes "successful" *vs* pathological brain ageing. Within this context, the review will then focus predominantly on the evolutionarily conserved insulin/IGF-signalling (IIS) system, and its roles in cognitive decline, dementia, AD and neuroinflammation.

It has been postulated that the IIS system plays a regulatory role in organismal ageing, lifespan and longevity^[15,16], and a reduced expression of its components under experimental conditions has been linked to amelioration of amyloid- β accumulation, the latter being one of the key features of AD^[17-19], and cognitive impairment^[20]. The IIS system merits an arduous and in depth investigation into the role and the extent of its involvement in cerebral aging, as targeting its components may pave the way to designing novel pharmacological approaches to early interventions and facilitate reversal or delay of cognitive decline in human patients.

SUCCESSFUL VS PATHOLOGICAL CEREBRAL AGEING

Defining successful cerebral ageing has been both challenging and controversial and to date there is no scientific consensus defining normal ageing in older age. Despite the lack of an operational definition, it is generally agreed that successful cerebral ageing is a multi-dimensional process, characterized by the absence of cognitive impairment and preservation of mental faculties, which allows for social functioning and independence in older age^[21]. This suggests that beyond the neurophysiological and psychological functions, equally vital are some esoteric elements such as wisdom and resilience, which together with lifestyle factors may contribute towards the variability, detected in cognitive abilities amongst "successful" *vs* "unsuccessful" elderly individuals and groups^[22,23].

Current research postulates that the ageing mammalian brain retains a considerable functional plasticity, which is activity-related and thus it depends on the lifestyle of the individual (*e.g.*,^[24]). There is evidence that human age-related diseases can be delayed by a healthy lifestyle which includes stress management, physical exercise and caloric restriction^[24,25]. Thus, although genes are important determinants of longevity, an individual's lifestyle is a powerful instrument that can delay the development of age-related diseases and lead to the path of ageing successfully^[25-27].

Functional imaging studies on ageing human brains, suggest that in the absence of pathology, age-related cognitive decline is rather modest and varies amongst individuals^[9]. It is characterized by anatomical and functional changes, which are associated with neuronal-synaptic molecular substrates specific to brain area^[28,29]. These changes may be attributed to synaptic connectivity rather than neuronal and white matter losses^[30,31].

In contrast, the pathologically ageing brain, as that in AD, exhibits marked cognitive decline, which is associated with a significant loss of synapses^[32]. Although the molecular mechanisms underlying this synaptic impairment are not fully understood, dysfunction of γ -secretase is evident in many cases of early onset of AD^[33], and the gamma-secretase-mediated EphA4 signalling system may be involved in the synaptic pathogenesis of AD^[32,33]. Equally, apolipoprotein E4 can increase the presence of amyloid beta (A β) oligomers in the brain, which in turn may increase the loss of dendritic spines and accelerate memory decline in AD^[34].

Of the regions associated with memory and learning, the hippocampal formation exhibit age-related decrease in volume, which may be a consequence of a decrease in neuronal and synaptic density^[28,30]; prefrontal cortex (PFC) also shows reductions in grey matter^[35,36]. The PFC is implicated in higher executive functions, involving explicit, implicit and spatial memories^[28,37,38]. Its decreased grey matter diffusivity may be a potential biomarker for early

AD^[39].

Similarly, the posterior cingulate, which is both anatomically and functionally connected to PFC and medial temporal lobe (MTL)^[40,41], and which plays a vital role in encoding and retrieval of information^[42,43] is also affected in ageing. Interestingly, it is one of the first structures to be affected in AD^[44,45] but has not yet been explicitly studied in conjunction with PFC to establish associations between task-related and cognitive task activation^[46].

These associations may be modifiable in healthy aged individuals^[46-48] in contrast to AD patients^[48-50]. PFC grey matter loss may trigger plasticity, which is dependent on the MTL function for memory tasks^[51], and consequently, those individuals with functionally intact grey matter/MTL ratio, may make a greater use of the PFC^[52].

The hippocampal region implicated in memory function shows age-related atrophy^[53], and a decline of working memory function is often observed in healthy ageing^[54]. In AD pathology, impaired hippocampal function is detectable even before the formation and accumulation of plaques^[55] and volume analysis by means of MRI has been used as diagnostic tool in distinguishing AD patients and healthy age-matched subjects by measuring the grey matter volumes in the lateral temporal and parietal cortices^[56]. Furthermore, the AD brain is characterized by ventricular enlargement^[57-60], consistent with a considerable loss of grey and white matter^[61].

The above alterations may be attributed to age-related neuronal loss, and/or compensatory plasticity, and future studies are needed to test how these three way structure-function-behaviour associations impact the grey matter loss and PFC activation in successful ageing^[46]. Equally, stress-related hormonal changes^[9] or compromised calcium homeostasis^[38,62] can play a role too as prolonged increases of intracellular calcium concentrations may cause neurite degeneration and cell death in ageing^[63].

In addition to the observed anatomical, functional and cellular changes, the brain's neurochemistry is also affected, with dopaminergic, noradrenergic and cholinergic systems exhibiting deficits^[60,64-69]. Studies on human and rhesus monkey PFC indicate that the balance between inhibitory and excitatory neurotransmission is decreased^[70] as an effect of reduced gene expression, which may compromise neural activity resulting in excitotoxicity and neurodegenerative pathology. Positron emission tomography scans in ageing humans show a reduction in dopamine synthesis in the striatum, of relevance to frontal lobe cognitive function^[71], and a marked decrease in dopamine receptor binding within caudate and putamen nuclei^[66,72].

Reductions in serotonin synthesis, reuptake and receptor binding have also been noted in the caudate nucleus, putamen and PFC of ageing brains^[67], and glutamate decreased levels in grey and white matter, basal ganglia have been reported^[69,73].

It is of significance that all the pathological features

of AD such as neuronal loss, neurofibrillary tangles and plaques may be present in the brains of elderly who may never show the full extent of cognitive deterioration observed in AD^[9]. This resilience to cognitive decline in the presence of AD pathology may be attributed to "cognitive reserve", which may reduce the risk of dementia in ageing^[74]. It further suggests that the hallmarks of AD may be secondary to ageing.

One of the cellular mechanisms regulating ageing processes is the insulin and IIS system, which is described below with regard to its role and those of its components in cerebral ageing. This system, extensively studied in model organisms, appears to underpin the innate resilience that is essential in successful ageing; it may also present therapeutic potential in the treatment of debilitating neurodegenerative and cerebrovascular diseases.

INSULIN, INSULIN GROWTH FACTORS AND THEIR RECEPTORS

Insulin and the IGF-1 and IGF-2 constitute a family of structurally similar peptides^[75,76], which have been preserved in most organisms through evolution^[77]. Peripherally, insulin is synthesized and secreted into blood by pancreatic cells, whereas IGF-1 and IGF-2 by the liver in response to the pituitary growth hormone^[78].

Insulin is a powerful player in glucose homeostasis, *e.g.*,^[79,80], which targets the liver, muscle, and adipose tissue^[81], and also the vasculature and the brain^[82]. The IGF-1 in contrast, is implicated in foetal and postnatal development, with a role in cellular survival of adult tissues^[82]. The circulation and delivery of IGF-1 to the tissues is aided by IGF-1 binding proteins 1-6 in contrast to insulin, which circulates freely^[82].

The transportation of insulin and IGF 1 into the brain is achieved through a saturable mechanism within the blood-brain barrier (BBB)^[78,82,83], although there is evidence of their *de novo* synthesis in the central nervous system (CNS)^[84-86]. Insulin's ability to cross the BBB^[83,87-89] depends on a number of factors such as age, fasting or obesity^[88]. Under experimental conditions, insulin administered directly into the CNS, decreases body weight by suppressing appetite, lowers serum insulin levels and increases serum glucose^[90,91]. An increase in peripheral insulin levels leads to increased cerebrospinal fluid (CSF) insulin, whereas chronic insulin resistance impairs cerebral transportation by down regulating insulin receptors (IR) at BBB^[92]. Brain activity in healthy individuals subjected to direct determination of insulin sensitivity with the hyperinsulinemic-euglycemic clamp technique, has been shown to be affected by increased levels of circulating insulin^[93].

BBB uptake of IGF involves a lipoprotein receptor-related protein 1, the respective receptor (IGF-1R) and other transport mechanisms, enabling IGF access to




Figure 1 Insulin and insulin-like growth factor signalling pathways-a simplified outline as based on ref. [77]. Low affinity binding between insulin and IGF-1R, and IGF-1 and IR, and IGF-2 and IR is represented by dashed lines. ACh: Acetylcholine; ERK: Extracellular signal-related kinase; GRB: Growth factor receptor-bound protein; IGF-1: Insulin-like growth factor 1; IR: Insulin receptor; IGF-1R: IGF-1 receptor; mTOR: Mammalian target of rapamycin; PLC: Phospholipase C; PKC: Protein kinase C; PI3K: Phosphoinositide 3-kinase; Ras: Signalling proteins involved in cell proliferation (name derived from rat sarcoma).

CSF, and hypothalamic and hippocampal regions^[82]. Insulin and IGFs activate signalling systems through their respective receptors, which belong to the tyrosine kinase receptor family^[77]: IR, IGF-1R and IGF-2R (Figure 1). Low affinity binding can take place between insulin and IGF-1R, and IGF-1 and IR using the phosphoinositide 3-kinase (PI3K)/AKT pathway, while IGF-2 signals downstream not only *via* IGF-2R but also IGF-1R^[77]. It should be mentioned that IGF-2R mainly controls the uptake and activation/destruction of extracellular of IGF-1/2 IGF2^[94]; the present review is focused on IGF-1R (Figure 1).

IR and IGF-1R differ in their respective functions and tissue expressions^[82] being present in the brain within neuron rich structures^[95-97] and glial cells^[18,98,99]; they are different entities expressed in diverse brain regions. The IR is highly expressed in the anterior thalamic and hypothalamic nuclei, olfactory bulb, hippocampus, cerebral cortex^[100,101] and promotes plasticity through supporting synaptogenesis and synaptic remodelling^[102,103], and metabolic homeostasis^[75,104]. IGF-1R signalling supports normal brain development with its neurogenesis, and successful ageing, consistent with the roles of its agonist neurotrophins (see review^[77]). A genetic modification, which results in IGF-1R deletion, causes microcephaly and death in experimental animals^[105,106].

INSULIN AND IGF-1 IIS AND ITS ROLES IN COGNITIVE FUNCTION, BRAIN AGEING AND LONGEVITY

The IIS system has been probably the most widely studied conserved mechanism that extends lifespan in worms, flies and mammals^[16], linking longevity and successful ageing across the species^[107,108]. It has been suggested that although insulin does not directly influence cerebral glucose transport, it appears to influence regionally the distribution of glucose transporter (GLUT) isoforms such as GLUT 4 and GLUT 8; the former being expressed in the cerebellum, hippocampus, pituitary and hypothalamus^[109] whereas the latter in the hippocampus and hypothalamus^[110]. This selective stimulation of glucose uptake in the brain areas implicated in learning and memory renders the hormone a potent player in potential therapeutic use to restore or enhance impaired cognitive functions^[111]. In addition, insulin's indirect role in hippocampal functioning *via* the long-term potentiation cascade, involving the N-methyl-D-aspartate receptor^[112] further suggests that insulin can be implicated in synaptic remodelling which is vital for the formation of new memories.

Similarly, insulin's ability to modulate CNS levels of acetylcholine and norepinephrine, known to influence

cognitive function^[113,114], may further substantiate its role in neural activity and potential neural protection against the effects of oxidative stress^[115].

Studies in animals and humans have suggested that deficiency or reduced effectiveness of insulin is a contributing factor to cognitive decline, impacting memory functions and brain ageing^[116-118]. Insulin resistance has been positively correlated with neurodegeneration observed in AD and mild cognitive impairment^[118,119]. Patients with type II diabetes mellitus, hypertension and chronic hyperinsulinemia show impaired verbal memory while enhanced memory function has been observed upon intranasal insulin administration^[120,121].

In preclinical studies, liver-specific IGF-1-deficient (LID) mice, deficient in liver-derived IGF-1 exhibit impaired spatial learning and memory functions^[122], demonstrating that this peptide is vital in mediating exercise-induced effects on the adult brain, thus suggesting promotion of neurogenesis^[123,124]. In addition, IGF-1 appears to possess neurotrophic properties and plays a role in the amelioration of age-related reduction in hippocampal neurogenesis and behavioural deficits^[125,126]; it also improves regional cerebral blood flow in normal rats^[127]. It targets brain neurones and glia implying a trophic action on glutamatergic synapses, modulating hippocampal circuitries that are involved in learning and memory^[116].

Within this context, research on brain ageing suggests that IGF-1 has potent effects on brain function, and that its reduced signalling during ageing may contribute to cognitive deterioration and compromise the organism's ability to deal with age-associated cerebral pathologies^[119]. Similarly, impaired insulin signalling in the brain has been linked to cognitive decline associated with pathological brain ageing^[111,128]. Epidemiological studies suggest that individuals with type II diabetes and obesity may be at higher risk of developing vascular dementia^[129].

On the other hand, centenarian studies provide evidence of correlation between reduced IIS activity and extreme human longevity. Ashkenazi centenarians have been found to have mutations in the IGF-1R that leads to lower activity of the respective signalling pathway^[107], and centenarians' offspring had lower peripheral IGF-1 activity when compared with appropriately matched-controls^[130]. Of significance, in the same sample of centenarians' offspring, IGF-1 was inversely related to insulin sensitivity^[130], and an Italian cohort study reported an IR variant to be associated with longevity^[131]. The above findings suggest that in humans the IIS system is a complex determinant of lifespan^[132].

PRECLINICAL VS CLINICAL STUDIES ON THE ROLE OF THE IIS SYSTEM IN AD

The importance of IR, IGF-1R and their respective agonist peptides has been underscored by data postulating the involvement of the IIS system in AD, which is characterized by amyloid-dependent neurodegeneration and late onset progressive cognitive decline. AD sufferers

display impaired cerebral glucoregulation^[133], reduced brain insulin receptor activity, reduced insulin concentrations in cerebrospinal fluid, peripheral hyperinsulinemia^[134] and reduced insulin and IGF-1 expression^[135], together with synaptic loss associated with the accumulation and formation of aggregate amyloid plaques (a β) and neurofibrillary tangles (tau protein) as measured post mortem^[136-138]. The above suggests that the IIS system may play a significant role in the loss of memory functions associated with AD and a reduction of its activity may reduce toxicity, delay a β accumulation and improve cognitive functions^[18,139-143].

In the preclinical approaches, mouse models of AD with a knockout of IGF-1 receptor exhibit reduced cognitive impairment, neurodegeneration and longer lifespan. The findings from the above studies, point to a β oligomers as the toxic species associated with AD, and the ageing process to be associated with the organism's exposure to their toxicity, leading to age-related neurodegenerative diseases^[132]. Consequently, the IIS system was mechanistically linked to neurone-associated toxic protein accumulation and ageing, as reduced signalling is thought to protect the brain and slow the progression of AD^[132]. It has been shown to activate the PI3K/AKT and Ras pathways (Figure 1). The former leads to the activation of mammalian TOR (mTOR), and rapamycin-treated mice have been shown to increase their lifespan and ameliorate age-related cognitive deficits^[141,144]. The latter activates extracellular signal-regulated kinase-1/-2, which has been implicated in plasticity, including long-term potentiation, and memory formation in the CNS^[145]. Although no data as yet exist to support the involvement of both pathways in the pathogenesis of AD, it may present an interesting direction for future research.

A number of current preclinical studies on the animal models of AD suggest that genetic reduction of the signalling pathway may protect against the AD pathology^[18,20] while older studies seem to postulate that its reduction is associated with the age-related pathologies^[146]. These conflicting views may arise from the use of different experimental approaches^[19], or as it has been suggested, reduced IGF-1 peripheral bioactivity may not necessarily induce the same results in brain IGF-1 levels^[147], postulating an independent regulatory activity.

It is important to appreciate the complexity of this relationship as human studies which try to correlate peripheral levels of IGF-1/2 with cognitive functioning in health and disease, report disparate finding, as illustrated below. Of the most recent publications in this area, a large scale long-term study on a community sample of over 3500 participants of middle and old ages has demonstrated that lower serum levels of IGF-1 associate with an increased risk of developing AD dementia while higher levels of IGF-1 associate with greater brain volumes in middle-aged participants free of stroke and dementia. The authors conclude that elevated levels of

IGF-1 may protect against neurodegeneration^[148]. On the other hand, the Caerphilly Prospective Study on 746 men has not found associations between age-related cognitive decline and IGF-1, contrary to IGF-2 which was associated with both normal age-related and pathological cognitive decline^[149]. Furthermore, offspring from families with a parental history of AD appear to have higher serum IGF-1 levels in middle age when compared with appropriate controls, leading to a conclusion that elevated peripheral IGF-1 associates with an increased risk of AD^[150].

Despite the existence of opposing views in the field, epidemiological evidence postulates a strong association between type II diabetes (T2D) and AD occurrence as AD patients exhibit higher rates of diabetes and impaired fasting glucose levels^[151-153], and although the molecular mechanisms underlying this association are not yet clearly understood^[82] impaired insulin signalling, amyloid-genesis and inflammation appear to be heavily implicated in the aetiology of diabetes, AD and consequently cerebral ageing^[154].

IIS SYSTEM AND INFLAMMATION

Inflammation is seen a key player in obesity, insulin resistance and diabetes, as based on elevated levels of pro-inflammatory cytokines in the circulation and pancreatic islets of T2D patients^[155]. Similarly, elevated levels of pro-inflammatory proteins and chemokines have been detected in post-mortem AD patients' brains, *e.g.*,^[156]. This was further substantiated by findings from studies on AD mouse models suggestive of inflammation as key to early and/or intermediate stages of the neurodegenerative condition^[157]. There is consensus that cerebrovascular inflammation and neuroinflammation, along with an increased accumulation of toxic $\alpha\beta$, all result in a disruption of synaptic activity, which according to some theories is a trigger in AD pathophysiology, *e.g.*,^[158].

Studies investigating IGF-1 and IGF-2 peptides' expression in human microglia *in vivo* and *in vitro* suggest that both peptides are expressed in microglia, conferring vital protection against cytokine-mediated neuronal death. It should be mentioned here that microglial activation is associated with increased activities of inflammatory cytokines, *e.g.*, interleukin (IL)-1 β and IL-6 which itself can disrupt neural signalling, *e.g.*,^[159].

Chronic inflammation increases the production of inflammatory cytokines in the long-term, which contributes to the suppression of neurotrophic factors, including the IGFs, and leads to progressive tissue damage, thus accelerating the onset of clinical manifestations of AD and metabolic disorders including T2D^[160,161], and may contribute to neurodegeneration^[162].

CONCLUSION

Research on brain ageing suggests that age itself is a major risk factor for the development of age-related cognitive

decline, Alzheimer's and cerebrovascular diseases. The increased life expectancy observed in most societies has further necessitated the need to understand the processes that underlie successful *vs* pathological brain ageing such that early interventions through lifestyle modifications or pharmacological agents may assist in delaying if not reversing the detrimental effects on brain pathology.

Within this context, this review examined the role of an evolutionarily conserved signalling pathway, IIS, with the focus on insulin and insulin-like growth factor IGF-1 and their roles in cerebral ageing. Translation of data derived from animal models allow for linking the IIS pathway with its supporting longevity, protein homeostasis, learning and memory, and delayed ageing. The above is also consistent with the human studies, which find evidence of reduced messaging for insulin, IGF-1 and their receptors in post mortem brains of patients with AD. While the link between insulin as such and brain ageing has been recognised, the IIS pathway in its entirety deserves more attention; our still incomplete understanding of the roles and mechanisms of this pathway calls for more translational research to explore novel treatments for cognitive decline through delaying cerebral ageing.

Some conflicting literature opinions and incomplete understanding of the roles and mechanisms of the IIS system demand novel approaches and directions in this field. The IIS system clearly lends itself to the ongoing search for modifiable physiological factors which may delay the onset of cognitive decline and cerebral ageing.

REFERENCES

- 1 **de Magalhães JP.** From cells to ageing: a review of models and mechanisms of cellular senescence and their impact on human ageing. *Exp Cell Res* 2004; **300**: 1-10 [PMID: 15383309 DOI: 10.1016/j.yexcr.2004.07.006]
- 2 **de Magalhães JP, Faragher RG.** Cell divisions and mammalian aging: integrative biology insights from genes that regulate longevity. *Bioessays* 2008; **30**: 567-578 [PMID: 18478536 DOI: 10.1002/bies.20760]
- 3 **Kirkwood TB.** The origins of human ageing. *Philos Trans R Soc Lond B Biol Sci* 1997; **352**: 1765-1772 [PMID: 9460059 DOI: 10.1098/rstb.1997.0160]
- 4 **Partridge L, Mangel M.** Messages from mortality: the evolution of death rates in the old. *Trends Ecol Evol* 1999; **14**: 438-442 [PMID: 10511720 DOI: 10.1016/S0169-5347(99)01646-8]
- 5 **Sharpless NE, DePinho RA.** How stem cells age and why this makes us grow old. *Nat Rev Mol Cell Biol* 2007; **8**: 703-713 [PMID: 17717515 DOI: 10.1038/nrm2241]
- 6 **Sonntag WE, Lynch C, Thornton P, Khan A, Bennett S, Ingram R.** The effects of growth hormone and IGF-1 deficiency on cerebrovascular and brain ageing. *J Anat* 2000; **197** Pt 4: 575-585 [PMID: 11197531 DOI: 10.1017/S002187829900713X]
- 7 **Mattay VS, Fera F, Tessitore A, Hariri AR, Das S, Callicott JH, Weinberger DR.** Neurophysiological correlates of age-related changes in human motor function. *Neurology* 2002; **58**: 630-635 [PMID: 11865144 DOI: 10.1212/WNL.58.4.630]
- 8 **Koivisto K, Reinikainen KJ, Hänninen T, Vanhanen M, Helkala EL, Mykkanen L, Laakso M, Pyörälä K, Riekkinen PJ.** Prevalence of age-associated memory impairment in a randomly selected population from eastern Finland. *Neurology*

- 1995; **45**: 741-747 [PMID: 7723964 DOI: 10.1212/WNL.45.4.741]
- 9 **Bishop NA**, Lu T, Yankner BA. Neural mechanisms of ageing and cognitive decline. *Nature* 2010; **464**: 529-535 [PMID: 20336135 DOI: 10.1038/nature08983]
- 10 **Muller FL**, Lustgarten MS, Jang Y, Richardson A, Van Remmen H. Trends in oxidative aging theories. *Free Radic Biol Med* 2007; **43**: 477-503 [PMID: 17640558 DOI: 10.1016/j.freeradbiomed.2007.03.034]
- 11 **Wolf FI**, Fasanella S, Tedesco B, Cavallini G, Donati A, Bergamini E, Cittadini A. Peripheral lymphocyte 8-OHdG levels correlate with age-associated increase of tissue oxidative DNA damage in Sprague-Dawley rats. Protective effects of caloric restriction. *Exp Gerontol* 2005; **40**: 181-188 [PMID: 15763395 DOI: 10.1016/j.exger.2004.11.002]
- 12 **Fischer A**, Sananbenesi F, Wang X, Dobbin M, Tsai LH. Recovery of learning and memory is associated with chromatin remodelling. *Nature* 2007; **447**: 178-182 [PMID: 17468743 DOI: 10.1038/nature05772]
- 13 **McGeer EG**, McGeer PL. Neuroinflammation in Alzheimer's disease and mild cognitive impairment: a field in its infancy. *J Alzheimers Dis* 2010; **19**: 355-361 [PMID: 20061650 DOI: 10.3233/JAD-2010-1219]
- 14 **Freitas AA**, de Magalhães JP. A review and appraisal of the DNA damage theory of ageing. *Mutat Res* 2011; **728**: 12-22 [PMID: 21600302 DOI: 10.1016/j.mrrev.2011.05.001]
- 15 **Blalock EM**, Geddes JW, Chen KC, Porter NM, Markesbery WR, Landfield PW. Incipient Alzheimer's disease: microarray correlation analyses reveal major transcriptional and tumor suppressor responses. *Proc Natl Acad Sci USA* 2004; **101**: 2173-2178 [PMID: 14769913 DOI: 10.1073/pnas.0308512100]
- 16 **Broughton S**, Partridge L. Insulin/IGF-like signalling, the central nervous system and aging. *Biochem J* 2009; **418**: 1-12 [PMID: 19159343 DOI: 10.1042/BJ20082102]
- 17 **Moloney AM**, Griffin RJ, Timmons S, O'Connor R, Ravid R, O'Neill C. Defects in IGF-1 receptor, insulin receptor and IRS-1/2 in Alzheimer's disease indicate possible resistance to IGF-1 and insulin signalling. *Neurobiol Aging* 2010; **31**: 224-243 [PMID: 18479783 DOI: 10.1016/j.neurobiolaging.2008.04.002]
- 18 **Freude S**, Hettich MM, Schumann C, Stöhr O, Koch L, Köhler C, Udelhoven M, Leiser U, Müller M, Kubota N, Kadowaki T, Krone W, Schröder H, Brüning JC, Schubert M. Neuronal IGF-1 resistance reduces Abeta accumulation and protects against premature death in a model of Alzheimer's disease. *FASEB J* 2009; **23**: 3315-3324 [PMID: 19487308 DOI: 10.1096/fj.09-132043]
- 19 **Piriz J**, Muller A, Trejo JL, Torres-Aleman I. IGF-I and the aging mammalian brain. *Exp Gerontol* 2011; **46**: 96-99 [PMID: 20863877 DOI: 10.1016/j.exger.2010.08.022]
- 20 **Cohen E**, Paulsson JF, Blinder P, Burstyn-Cohen T, Du D, Estepa G, Adame A, Pham HM, Holzenberger M, Kelly JW, Masliah E, Dillin A. Reduced IGF-1 signaling delays age-associated proteotoxicity in mice. *Cell* 2009; **139**: 1157-1169 [PMID: 20005808 DOI: 10.1016/j.cell.2009.11.014]
- 21 **Hendrie HC**, Albert MS, Butters MA, Gao S, Knopman DS, Launer LJ, Yaffe K, Cuthbert BN, Edwards E, Wagster MV. The NIH Cognitive and Emotional Health Project. Report of the Critical Evaluation Study Committee. *Alzheimers Dement* 2006; **2**: 12-32 [PMID: 19595852 DOI: 10.1016/j.jalz.2005.11.004]
- 22 **Lamond AJ**, Depp CA, Allison M, Langer R, Reichstadt J, Moore DJ, Golshan S, Ganiats TG, Jeste DV. Measurement and predictors of resilience among community-dwelling older women. *J Psychiatr Res* 2008; **43**: 148-154 [PMID: 18455190 DOI: 10.1016/j.jpsychires.2008.03.007]
- 23 **Jeste DV**, Ardelt M, Blazer D, Kraemer HC, Vaillant G, Meeks TW. Expert consensus on characteristics of wisdom: a Delphi method study. *Gerontologist* 2010; **50**: 668-680 [PMID: 20233730 DOI: 10.1093/geront/gnq022]
- 24 **Stranahan AM**, Mattson MP. Recruiting adaptive cellular stress responses for successful brain ageing. *Nat Rev Neurosci* 2012; **13**: 209-216 [PMID: 22251954 DOI: 10.1038/nrn3151]
- 25 **Gitler AD**. Another reason to exercise. *Science* 2011; **334**: 606-607 [DOI: 10.1126/science.1214714]
- 26 **Abbott A**. Ageing: growing old gracefully. *Nature* 2004; **428**: 116-118 [PMID: 15014466 DOI: 10.1038/428116a]
- 27 **Mora F**, Segovia G, Del Arco A, de Blas M, Garrido P. Stress, neurotransmitters, corticosterone and body-brain integration. *Brain Res* 2012; **1476**: 71-85 [PMID: 22285436 DOI: 10.1016/j.brainres.2011.12.049]
- 28 **Hedden T**, Gabrieli JD. Insights into the ageing mind: a view from cognitive neuroscience. *Nat Rev Neurosci* 2004; **5**: 87-96 [PMID: 14735112 DOI: 10.1038/nrn1323]
- 29 **Burke SN**, Barnes CA. Neural plasticity in the ageing brain. *Nat Rev Neurosci* 2006; **7**: 30-40 [PMID: 16371948 DOI: 10.1038/nrn1809]
- 30 **Morrison JH**, Baxter MG. The ageing cortical synapse: hallmarks and implications for cognitive decline. *Nat Rev Neurosci* 2012; **13**: 240-250 [PMID: 22395804 DOI: 10.1038/nrn3200]
- 31 **Balietti M**, Tamagnini F, Fattoretti P, Burattini C, Casoli T, Platano D, Lattanzio F, Aicardi G. Impairments of synaptic plasticity in aged animals and in animal models of Alzheimer's disease. *Rejuvenation Res* 2012; **15**: 235-238 [PMID: 22533439 DOI: 10.1089/rej.2012.1318]
- 32 **Matsui C**, Inoue E, Kakita A, Arita K, Deguchi-Tawarada M, Togawa A, Yamada A, Takai Y, Takahashi H. Involvement of the γ -secretase-mediated EphA4 signaling pathway in synaptic pathogenesis of Alzheimer's disease. *Brain Pathol* 2012; **22**: 776-787 [PMID: 22404518 DOI: 10.1111/j.1750-3639.2012.00587.x]
- 33 **Inoue E**, Deguchi-Tawarada M, Togawa A, Matsui C, Arita K, Katahira-Tayama S, Sato T, Yamauchi E, Oda Y, Takai Y. Synaptic activity prompts gamma-secretase-mediated cleavage of EphA4 and dendritic spine formation. *J Cell Biol* 2009; **185**: 551-564 [PMID: 19414612 DOI: 10.1083/jcb.200809151]
- 34 **Hashimoto T**, Serrano-Pozo A, Hori Y, Adams KW, Takeda S, Banerji AO, Mitani A, Joyner D, Thyssen DH, Bacskai BJ, Frosch MP, Spires-Jones TL, Finn MB, Holtzman DM, Hyman BT. Apolipoprotein E, especially apolipoprotein E4, increases the oligomerization of amyloid beta peptide. *J Neurosci* 2012; **32**: 1581-1592 [DOI: 10.1523/JNEUROSCI.1542-12.2012]
- 35 **Raz N**, Gunning FM, Head D, Dupuis JH, McQuain J, Briggs SD, Loken WJ, Thornton AE, Acker JD. Selective aging of the human cerebral cortex observed in vivo: differential vulnerability of the prefrontal gray matter. *Cereb Cortex* 1997; **7**: 268-282 [PMID: 9143446 DOI: 10.1093/cercor/7.3.268]
- 36 **Salat DH**, Buckner RL, Snyder AZ, Greve DN, Desikan RS, Busa E, Morris JC, Dale AM, Fischl B. Thinning of the cerebral cortex in aging. *Cereb Cortex* 2004; **14**: 721-730 [PMID: 15054051 DOI: 10.1093/cercor/bhh032]
- 37 **Woodruff-Pak DS**, Thompson RF. Classical conditioning of the eyeblink response in the delay paradigm in adults aged 18-83 years. *Psychol Aging* 1988; **3**: 219-229 [PMID: 3268262 DOI: 10.1037/0882-7974.3.3.219]
- 38 **Kenessary A**, Zhumadilov Z, Nurgozhin T, Kipling D, Yeoman M, Cox L, Ostler E, Faragher R. Biomarkers, interventions and healthy ageing. *N Biotechnol* 2013; **30**: 373-377 [PMID: 23201073 DOI: 10.1016/j.nbt.2012.11.018]
- 39 **Jacobs HI**, van Bostel MP, Gronenschild EH, Uyilings HB, Jolles J, Verhey FR. Decreased gray matter diffusivity: a potential early Alzheimer's disease biomarker? *Alzheimers Dement* 2013; **9**: 93-97 [PMID: 22651939 DOI: 10.1016/j.jalz.2011.11.004]
- 40 **Greicius MD**, Supekar K, Menon V, Dougherty RF. Resting-state functional connectivity reflects structural connectivity in the default mode network. *Cereb Cortex* 2009; **19**: 72-78 [PMID: 18403396 DOI: 10.1093/cercor/bhn059]
- 41 **Dörfel D**, Werner A, Schaefer M, von Kummer R, Karl A. Distinct brain networks in recognition memory share

- a defined region in the precuneus. *Eur J Neurosci* 2009; **30**: 1947-1959 [PMID: 19895564 DOI: 10.1111/j.1460-9568.2009.06973.x]
- 42 **Vannini P**, O'Brien J, O'Keefe K, Pihlajamäki M, Laviolette P, Sperling RA. What goes down must come up: role of the posteromedial cortices in encoding and retrieval. *Cereb Cortex* 2011; **21**: 22-34 [PMID: 20363808 DOI: 10.1093/cercor/bhq051]
- 43 **Huijbers W**, Vannini P, Sperling RA, C M P, Cabeza R, Daselaar SM. Explaining the encoding/retrieval flip: memory-related deactivations and activations in the posteromedial cortex. *Neuropsychologia* 2012; **50**: 3764-3774 [PMID: 22982484 DOI: 10.1016/j.neuropsychologia.2012.08.021]
- 44 **Minoshima S**, Giordani B, Berent S, Frey KA, Foster NL, Kuhl DE. Metabolic reduction in the posterior cingulate cortex in very early Alzheimer's disease. *Ann Neurol* 1997; **42**: 85-94 [PMID: 9225689]
- 45 **Pengas G**, Hodges JR, Watson P, Nestor PJ. Focal posterior cingulate atrophy in incipient Alzheimer's disease. *Neurobiol Aging* 2010; **31**: 25-33 [PMID: 18455838 DOI: 10.1016/j.neurobiolaging.2008.03.014]
- 46 **Maillet D**, Rajah MN. Association between prefrontal activity and volume change in prefrontal and medial temporal lobes in aging and dementia: a review. *Ageing Res Rev* 2013; **12**: 479-489 [PMID: 23183352 DOI: 10.1016/j.arr.2012.11.001]
- 47 **Brassen S**, Büchel C, Weber-Fahr W, Lehmbeck JT, Sommer T, Braus DF. Structure-function interactions of correct retrieval in healthy elderly women. *Neurobiol Aging* 2009; **30**: 1147-1156 [PMID: 18023505]
- 48 **Trivedi MA**, Stoub TR, Murphy CM, George S, deToledo-Morrell L, Shah RC, Whitfield-Gabrieli S, Gabrieli JD, Stebbins GT. Entorhinal cortex volume is associated with episodic memory related brain activation in normal aging and amnesic mild cognitive impairment. *Brain Imaging Behav* 2011; **5**: 126-136 [PMID: 21328083 DOI: 10.1007/s11682-011-9117-4]
- 49 **Rémy F**, Mirrashed F, Campbell B, Richter W. Verbal episodic memory impairment in Alzheimer's disease: a combined structural and functional MRI study. *Neuroimage* 2005; **25**: 253-266 [PMID: 15734360]
- 50 **Meulenbroek O**, Rijpkema M, Kessels RP, Rikkert MG, Fernández G. Autobiographical memory retrieval in patients with Alzheimer's disease. *Neuroimage* 2010; **53**: 331-340 [PMID: 20570740 DOI: 10.1016/j.neuroimage.2010.05.082]
- 51 **Greenwood PM**. Functional plasticity in cognitive aging: review and hypothesis. *Neuropsychology* 2007; **21**: 657-673 [PMID: 17983277 DOI: 10.1037/0894-4105.21.6.657]
- 52 **Rosen AC**, Gabrieli JD, Stoub T, Prull MW, O'Hara R, Yesavage J, deToledo-Morrell L. Relating medial temporal lobe volume to frontal fMRI activation for memory encoding in older adults. *Cortex* 2005; **41**: 595-602 [PMID: 16042035 DOI: 10.1016/S0010-9452(08)70199-0]
- 53 **Du AT**, Schuff N, Chao LL, Kornak J, Jagust WJ, Kramer JH, Reed BR, Miller BL, Norman D, Chui HC, Weiner MW. Age effects on atrophy rates of entorhinal cortex and hippocampus. *Neurobiol Aging* 2006; **27**: 733-740 [PMID: 15961190 DOI: 10.1016/j.neurobiolaging.2005.03.021]
- 54 **Wang M**, Gambo NJ, Yang Y, Jin LE, Wang XJ, Laubach M, Mazer JA, Lee D, Arnsten AF. Neuronal basis of age-related working memory decline. *Nature* 2011; **476**: 210-213 [PMID: 21796118 DOI: 10.1038/nature10243]
- 55 **Marchetti C**, Marie H. Hippocampal synaptic plasticity in Alzheimer's disease: what have we learned so far from transgenic models? *Rev Neurosci* 2011; **22**: 373-402 [PMID: 21732714 DOI: 10.1515/RNS.2011.035]
- 56 **Hänggi J**, Streffer J, Jäncke L, Hock C. Volumes of lateral temporal and parietal structures distinguish between healthy aging, mild cognitive impairment, and Alzheimer's disease. *J Alzheimers Dis* 2011; **26**: 719-734 [PMID: 21709375 DOI: 10.3233/JAD-2011-101260]
- 57 **Carmichael OT**, Kuller LH, Lopez OL, Thompson PM, Dutton RA, Lu A, Lee SE, Lee JY, Aizenstein HJ, Meltzer CC, Liu Y, Toga AW, Becker JT. Ventricular volume and dementia progression in the Cardiovascular Health Study. *Neurobiol Aging* 2007; **28**: 389-397 [PMID: 16504345 DOI: 10.1016/j.neurobiolaging.2006.01.006]
- 58 **Carmichael OT**, Kuller LH, Lopez OL, Thompson PM, Dutton RA, Lu A, Lee SE, Lee JY, Aizenstein HJ, Meltzer CC, Liu Y, Toga AW, Becker JT. Cerebral ventricular changes associated with transitions between normal cognitive function, mild cognitive impairment, and dementia. *Alzheimer Dis Assoc Disord* 2007; **21**: 14-24 [PMID: 17334268 DOI: 10.1093/brain/awn146]
- 59 **Nestor SM**, Rupsingh R, Borrie M, Smith M, Accomazzi V, Wells JL, Fogarty J, Bartha R; Alzheimer's Disease Neuroimaging Initiative. Ventricular enlargement as a possible measure of Alzheimer's disease progression validated using the Alzheimer's disease neuroimaging initiative database. *Brain* 2008; **131**: 2443-2454
- 60 **Apostolova LG**, Mosconi L, Thompson PM, Green AE, Hwang KS, Ramirez A, Mistur R, Tsui WH, de Leon MJ. Subregional hippocampal atrophy predicts Alzheimer's dementia in the cognitively normal. *Neurobiol Aging* 2010; **31**: 1077-1088 [PMID: 18814937 DOI: 10.1016/j.neurobiolaging.2008.08.008]
- 61 **Salat DH**, Kaye JA, Janowsky JS. Prefrontal gray and white matter volumes in healthy aging and Alzheimer disease. *Arch Neurol* 1999; **56**: 338-344 [PMID: 10190825 DOI: 10.1001/archneur.56.3.338]
- 62 **Toescu EC**, Verkhatsky A, Landfield PW. Ca²⁺ regulation and gene expression in normal brain aging. *Trends Neurosci* 2004; **27**: 614-620 [PMID: 15374673 DOI: 10.1016/j.tins.2004.07.010]
- 63 **Kapogiannis D**, Mattson MP. Disrupted energy metabolism and neuronal circuit dysfunction in cognitive impairment and Alzheimer's disease. *Lancet Neurol* 2011; **10**: 187-198 [PMID: 21147038 DOI: 10.1016/S1474-4422(10)70277-5]
- 64 **Rinne JO**, Lönnberg P, Marjamäki P. Age-dependent decline in human brain dopamine D1 and D2 receptors. *Brain Res* 1990; **508**: 349-352 [PMID: 2407314 DOI: 10.1016/0006-8993(90)90423-9]
- 65 **Iyo M**, Yamasaki T. The detection of age-related decrease of dopamine D1, D2 and serotonin 5-HT₂ receptors in living human brain. *Prog Neuropsychopharmacol Biol Psychiatry* 1993; **17**: 415-421 [PMID: 8475323 DOI: 10.1016/0278-5846(93)9007-5]
- 66 **Wang Y**, Chan GL, Holden JE, Dobko T, Mak E, Schulzer M, Huser JM, Snow BJ, Ruth TJ, Calne DB, Stoessl AJ. Age-dependent decline of dopamine D1 receptors in human brain: a PET study. *Synapse* 1998; **30**: 56-61 [PMID: 9704881 DOI: 10.1002/(SICI)1098-2396(199809)30:1<56::AID-SYN7>3.0.CO;2-J]
- 67 **Yamamoto M**, Suhara T, Okubo Y, Ichimiya T, Sudo Y, Inoue M, Takano A, Yasuno F, Yoshikawa K, Tanada S. Age-related decline of serotonin transporters in living human brain of healthy males. *Life Sci* 2002; **71**: 751-757 [PMID: 12074934 DOI: 10.1016/S0024-3205(02)01745-9]
- 68 **Ota M**, Yasuno F, Ito H, Seki C, Nozaki S, Asada T, Suhara T. Age-related decline of dopamine synthesis in the living human brain measured by positron emission tomography with L-[beta-11C]DOPA. *Life Sci* 2006; **79**: 730-736 [PMID: 16580023 DOI: 10.1016/j.lfs.2006.02.017]
- 69 **Chang L**, Jiang CS, Ernst T. Effects of age and sex on brain glutamate and other metabolites. *Magn Reson Imaging* 2009; **27**: 142-145 [PMID: 18687554 DOI: 10.1016/j.mri.2008.06.002]
- 70 **Loerch PM**, Lu T, Dakin KA, Vann JM, Isaacs A, Geula C, Wang J, Pan Y, Gabuzda DH, Li C, Prolla TA, Yankner BA. Evolution of the aging brain transcriptome and synaptic regulation. *PLoS One* 2008; **3**: e3329 [PMID: 18830410 DOI: 10.1371/journal.pone.0003329]

- 71 **Braskie MN**, Wilcox CE, Landau SM, O'Neil JP, Baker SL, Madison CM, Kluth JT, Jagust WJ. Relationship of striatal dopamine synthesis capacity to age and cognition. *J Neurosci* 2008; **28**: 14320-14328 [PMID: 19109513 DOI: 10.1523/JNEUROSCI.3729-08.2008]
- 72 **Kaasinen V**, Vilkinen H, Hietala J, Nägren K, Helenius H, Olsson H, Farde L, Rinne J. Age-related dopamine D2/D3 receptor loss in extrastriatal regions of the human brain. *Neurobiol Aging* 2000; **21**: 683-688 [PMID: 11016537 DOI: 10.1016/S0197-4580(00)00149-4]
- 73 **Sailasuta N**, Ernst T, Chang L. Regional variations and the effects of age and gender on glutamate concentrations in the human brain. *Magn Reson Imaging* 2008; **26**: 667-675 [PMID: 17692491 DOI: 10.1016/j.mri.2007.06.007]
- 74 **Honer WG**, Barr AM, Sawada K, Thornton AE, Morris MC, Leurgans SE, Schneider JA, Bennett DA. Cognitive reserve, presynaptic proteins and dementia in the elderly. *Transl Psychiatry* 2012; **2**: e114 [PMID: 22832958 DOI: 10.1038/tp.2012.38]
- 75 **Bondy CA**, Cheng CM. Signaling by insulin-like growth factor 1 in brain. *Eur J Pharmacol* 2004; **490**: 25-31 [PMID: 15094071 DOI: 10.1016/j.ejphar.2004.02.042]
- 76 **Schulingkamp RJ**, Pagano TC, Hung D, Raffa RB. Insulin receptors and insulin action in the brain: review and clinical implications. *Neurosci Biobehav Rev* 2000; **24**: 855-872 [PMID: 11118610 DOI: 10.1016/S0149-7634(00)00040-3]
- 77 **Fernandez AM**, Torres-Alemán I. The many faces of insulin-like peptide signalling in the brain. *Nat Rev Neurosci* 2012; **13**: 225-239 [PMID: 22430016 DOI: 10.1038/nrn3209]
- 78 **Skottner A**. Biosynthesis of Growth Hormone and Insulin-Like Growth Factor-I and the Regulation of their Secretion. *Open Endocrinol J* 2012; **6** (Suppl 1): 3-12
- 79 **Banting FG**, Best CH, Collip JB, Campbell WR, Fletcher AA. Pancreatic Extracts in the Treatment of Diabetes Mellitus. *Can Med Assoc J* 1922; **12**: 141-146 [PMID: 20314060]
- 80 **Zhao WQ**, Chen H, Quon MJ, Alkon DL. Insulin and the insulin receptor in experimental models of learning and memory. *Eur J Pharmacol* 2004; **490**: 71-81 [PMID: 15094074 DOI: 10.1016/j.ejphar.2004.02.045]
- 81 **Fulop T**, Larbi A, Douziech N. Insulin receptor and ageing. *Pathol Biol (Paris)* 2003; **51**: 574-580 [PMID: 14622948]
- 82 **Werner H**, LeRoith D. Insulin and insulin-like growth factor receptors in the brain: Physiological and pathological aspects. *Eur Neuropsychopharmacol* 2014; **24**: 1947-1953 [PMID: 24529663 DOI: 10.1016/j.euroneuro.2014.01.020]
- 83 **Schechter R**, Whitmire J, Holtzclaw L, George M, Harlow R, Devaskar SU. Developmental regulation of insulin in the mammalian central nervous system. *Brain Res* 1992; **582**: 27-37 [PMID: 1482442 DOI: 10.1016/0006-8993(92)90313-X]
- 84 **Lupu F**, Terwilliger JD, Lee K, Segre GV, Efstratiadis A. Roles of growth hormone and insulin-like growth factor 1 in mouse postnatal growth. *Dev Biol* 2001; **229**: 141-162 [PMID: 11133160 DOI: 10.1016/j.patbio.2003.09.007]
- 85 **Sun LY**, Al-Regaiey K, Masternak MM, Wang J, Bartke A. Local expression of GH and IGF-1 in the hippocampus of GH-deficient long-lived mice. *Neurobiol Aging* 2005; **26**: 929-937 [PMID: 15718052]
- 86 **Costantini C**, Scrable H, Puglielli L. An aging pathway controls the TrkA to p75NTR receptor switch and amyloid beta-peptide generation. *EMBO J* 2006; **25**: 1997-2006 [PMID: 16619032]
- 87 **Margolis RU**, Altszuler N. Insulin in the cerebrospinal fluid. *Nature* 1967; **215**: 1375-1376 [PMID: 6055448]
- 88 **Banks WA**. The source of cerebral insulin. *Eur J Pharmacol* 2004; **490**: 5-12 [PMID: 15094069 DOI: 10.1016/j.ejphar.2004.02.040]
- 89 **Burns JM**, Donnelly JE, Anderson HS, Mayo MS, Spencer-Gardner L, Thomas G, Cronk BB, Haddad Z, Klima D, Hansen D, Brooks WM. Peripheral insulin and brain structure in early Alzheimer disease. *Neurology* 2007; **69**: 1094-1104 [PMID: 17846409 DOI: 10.1212/01.wnl.0000276952.91704.af]
- 90 **Florant GL**, Singer L, Scheurink AJ, Park CR, Richardson RD, Woods SC. Intraventricular insulin reduces food intake and body weight of marmosets during the summer feeding period. *Physiol Behav* 1991; **49**: 335-338 [PMID: 1905822 DOI: 10.1016/0031-9384(91)90053-Q]
- 91 **Foster LA**, Ames NK, Emery RS. Food intake and serum insulin responses to intraventricular infusions of insulin and IGF-I. *Physiol Behav* 1991; **50**: 745-749 [PMID: 1663628 DOI: 10.1016/0031-9384(91)90012-D]
- 92 **Moreira PI**, Duarte AI, Santos MS, Rego AC, Oliveira CR. An integrative view of the role of oxidative stress, mitochondria and insulin in Alzheimer's disease. *J Alzheimers Dis* 2009; **16**: 741-761 [PMID: 19387110 DOI: 10.3233/JAD-2009-0972]
- 93 **Freude S**, Plum L, Schnitker J, Leeser U, Udelhoven M, Krone W, Bruning JC, Schubert M. Peripheral hyperinsulinemia promotes tau phosphorylation in vivo. *Diabetes* 2005; **54**: 3343-3348 [PMID: 16306348 DOI: 10.2337/diabetes.54.12.3343]
- 94 **Hawkes C**, Kar S. The insulin-like growth factor-II/mannose-6-phosphate receptor: structure, distribution and function in the central nervous system. *Brain Res Brain Res Rev* 2004; **44**: 117-140 [PMID: 15003389 DOI: 10.1016/j.brainresrev.2003.11.002]
- 95 **de la Monte SM**, Wands JR. Review of insulin and insulin-like growth factor expression, signaling, and malfunction in the central nervous system: relevance to Alzheimer's disease. *J Alzheimers Dis* 2005; **7**: 45-61 [PMID: 15750214]
- 96 **Hill JM**, Lesniak MA, Pert CB, Roth J. Autoradiographic localization of insulin receptors in rat brain: prominence in olfactory and limbic areas. *Neuroscience* 1986; **17**: 1127-1138 [PMID: 3520377 DOI: 10.1016/0306-4522(86)90082-5]
- 97 **Banks WA**, Jaspan JB, Huang W, Kastin AJ. Transport of insulin across the blood-brain barrier: saturability at euglycemic doses of insulin. *Peptides* 1997; **18**: 1423-1429 [PMID: 9392846 DOI: 10.1016/S0196-9781(97)00231-3]
- 98 **Broughton SK**, Chen H, Riddle A, Kuhn SE, Nagalla S, Roberts CT, Back SA. Large-scale generation of highly enriched neural stem-cell-derived oligodendroglial cultures: maturation-dependent differences in insulin-like growth factor-mediated signal transduction. *J Neurochem* 2007; **100**: 628-638 [PMID: 17263792 DOI: 10.1111/j.1471-4159.2006.04171.x]
- 99 **Zeger M**, Popken G, Zhang J, Xuan S, Lu QR, Schwab MH, Nave KA, Rowitch D, D'Ercolo AJ, Ye P. Insulin-like growth factor type 1 receptor signaling in the cells of oligodendrocyte lineage is required for normal in vivo oligodendrocyte development and myelination. *Glia* 2007; **55**: 400-411 [PMID: 17186502 DOI: 10.1002/glia.20469]
- 100 **Havrankova J**, Roth J, Brownstein M. Insulin receptors are widely distributed in the central nervous system of the rat. *Nature* 1978; **272**: 827-829 [PMID: 205798 DOI: 10.1038/272827a0]
- 101 **Unger JW**, Livingston JN, Moss AM. Insulin receptors in the central nervous system: localization, signalling mechanisms and functional aspects. *Prog Neurobiol* 1991; **36**: 343-362 [PMID: 1887067 DOI: 10.1016/0301-0082(91)90015-S]
- 102 **Chiu SL**, Cline HT. Insulin receptor signaling in the development of neuronal structure and function. *Neural Dev* 2010; **5**: 7 [PMID: 20230616 DOI: 10.1186/1749-8104-5-7]
- 103 **Abbott MA**, Wells DG, Fallon JR. The insulin receptor tyrosine kinase substrate p58/53 and the insulin receptor are components of CNS synapses. *J Neurosci* 1999; **19**: 7300-7308 [PMID: 10460236]
- 104 **McNay EC**, Ong CT, McCrimmon RJ, Cresswell J, Bogan JS, Sherwin RS. Hippocampal memory processes are modulated by insulin and high-fat-induced insulin resistance. *Neurobiol Learn Mem* 2010; **93**: 546-553 [PMID: 20176121 DOI: 10.1016/j.nlm.2010.02.002]
- 105 **Baker J**, Liu JP, Robertson EJ, Efstratiadis A. Role of insulin-like growth factors in embryonic and postnatal growth. *Cell* 1993; **75**: 73-82 [PMID: 8402902]
- 106 **Vicario-Abejón C**, Yusta-Boyo MJ, Fernández-Moreno C, de

- Pablo F. Locally born olfactory bulb stem cells proliferate in response to insulin-related factors and require endogenous insulin-like growth factor-I for differentiation into neurons and glia. *J Neurosci* 2003; **23**: 895-906 [PMID: 12574418]
- 107 **Suh Y**, Atzmon G, Cho MO, Hwang D, Liu B, Leahy DJ, Barzilai N, Cohen P. Functionally significant insulin-like growth factor I receptor mutations in centenarians. *Proc Natl Acad Sci USA* 2008; **105**: 3438-3442 [PMID: 18316725 DOI: 10.1073/pnas.0705467105]
- 108 **Flachsbarth F**, Caliebe A, Kleindorp R, Blanché H, von Eller-Eberstein H, Nikolaus S, Schreiber S, Nebel A. Association of FOXO3A variation with human longevity confirmed in German centenarians. *Proc Natl Acad Sci USA* 2009; **106**: 2700-2705 [PMID: 19196970 DOI: 10.1073/pnas.0809594106]
- 109 **Reagan LP**, Gorovits N, Hoskin EK, Alves SE, Katz EB, Grillo CA, Piroli GG, McEwen BS, Charron MJ. Localization and regulation of GLUTx1 glucose transporter in the hippocampus of streptozotocin diabetic rats. *Proc Natl Acad Sci USA* 2001; **98**: 2820-2825 [PMID: 11226324 DOI: 10.1073/pnas.051629798]
- 110 **Brant AM**, Jess TJ, Milligan G, Brown CM, Gould GW. Immunological analysis of glucose transporters expressed in different regions of the rat brain and central nervous system. *Biochem Biophys Res Commun* 1993; **192**: 1297-1302 [PMID: 8507199 DOI: 10.1006/bbrc.1993.1557]
- 111 **Cholerton B**, Baker LD, Craft S. Insulin resistance and pathological brain ageing. *Diabet Med* 2011; **28**: 1463-1475 [PMID: 21974744 DOI: 10.1111/j.1464-5491.2011.03464.x]
- 112 **Skeberdis VA**, Lan J, Zheng X, Zukin RS, Bennett MV. Insulin promotes rapid delivery of N-methyl-D- aspartate receptors to the cell surface by exocytosis. *Proc Natl Acad Sci USA* 2001; **98**: 3561-3566 [PMID: 11248117 DOI: 10.1073/pnas.051634698]
- 113 **Figlewicz DP**, Szot P, Israel PA, Payne C, Dorsa DM. Insulin reduces norepinephrine transporter mRNA in vivo in rat locus coeruleus. *Brain Res* 1993; **602**: 161-164 [PMID: 8448653 DOI: 10.1016/0006-8993(93)90258-O]
- 114 **Kopf D**, Baratti M. Effects of posttraining administration of insulin on retention of a habituation response in mice: participation of a central cholinergic mechanism. *Neurobiol Learn Mem* 1999; **71**: 50-61 [DOI: 10.1006/nlme.1998.3831]
- 115 **Duarte AI**, Santos MS, Seica R, Oliveira CR. Oxidative stress affects synaptosomal gamma-aminobutyric acid and glutamate transport in diabetic rats: the role of insulin. *Diabetes* 2004; **53**: 2110-2116 [PMID: 15277393 DOI: 10.2337/diabetes.53.8.2110]
- 116 **Trejo JL**, Carro E, Torres-Aleman I. Circulating insulin-like growth factor I mediates exercise-induced increases in the number of new neurons in the adult hippocampus. *J Neurosci* 2001; **21**: 1628-1634 [PMID: 11222653]
- 117 **Zhao WQ**, Alkon DL. Role of insulin and insulin receptor in learning and memory. *Mol Cell Endocrinol* 2001; **177**: 125-134 [PMID: 11377828 DOI: 10.1016/S0303-7207(01)00455-5]
- 118 **Talbot K**, Wang HY, Kazi H, Han LY, Bakshi KP, Stucky A, Fuino RL, Kawaguchi KR, Samoyedny AJ, Wilson RS, Arvanitakis Z, Schneider JA, Wolf BA, Bennett DA, Trojanowski JQ, Arnold SE. Demonstrated brain insulin resistance in Alzheimer's disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. *J Clin Invest* 2012; **122**: 1316-1338 [PMID: 22476197 DOI: 10.1172/JCI59903]
- 119 **Frölich L**, Blum-Degen D, Bernstein HG, Engelsberger S, Humrich J, Laufer S, Muschner D, Thalheimer A, Türk A, Hoyer S, Zöchling R, Boissl KW, Jellinger K, Riederer P. Brain insulin and insulin receptors in aging and sporadic Alzheimer's disease. *J Neural Transm* 1998; **105**: 423-438 [PMID: 9720972 DOI: 10.1007/s007020050068]
- 120 **Wallum BJ**, Taborsky GJ, Porte D, Figlewicz DP, Jacobson L, Beard JC, Ward WK, Dorsa D. Cerebrospinal fluid insulin levels increase during intravenous insulin infusions in man. *J Clin Endocrinol Metab* 1987; **64**: 190-194 [PMID: 3536982 DOI: 10.1210/jcem-64-1-190]
- 121 **Reger MA**, Watson GS, Frey WH, Baker LD, Cholerton B, Keeling ML, Belongia DA, Fishel MA, Plymate SR, Schellenberg GD, Cherrier MM, Craft S. Effects of intranasal insulin on cognition in memory-impaired older adults: modulation by APOE genotype. *Neurobiol Aging* 2006; **27**: 451-458 [PMID: 15964100 DOI: 10.1016/j.neurobiolaging.2005.03.016]
- 122 **Svensson J**, Diez M, Engel J, Wass C, Tivesten A, Jansson JO, Isaksson O, Archer T, Hökfelt T, Ohlsson C. Endocrine, liver-derived IGF-I is of importance for spatial learning and memory in old mice. *J Endocrinol* 2006; **189**: 617-627 [PMID: 16731792 DOI: 10.1677/joe.1.06631]
- 123 **O'Kusky JR**, Ye P, D'Ercole AJ. Insulin-like growth factor-I promotes neurogenesis and synaptogenesis in the hippocampal dentate gyrus during postnatal development. *J Neurosci* 2000; **20**: 8435-8442 [PMID: 11069951]
- 124 **Torres-Aleman I**. Serum growth factors and neuroprotective surveillance: focus on IGF-I. *Mol Neurobiol* 2000; **21**: 153-160 [PMID: 11379797 DOI: 10.1385/MN:21:3:153]
- 125 **Lichtenwalner RJ**, Forbes ME, Bennett SA, Lynch CD, Sonntag WE, Riddle DR. Intracerebroventricular infusion of insulin-like growth factor-I ameliorates the age-related decline in hippocampal neurogenesis. *Neuroscience* 2001; **107**: 603-613 [PMID: 11720784 DOI: 10.1016/S0306-4522(01)00378-5]
- 126 **Markowska AL**, Mooney M, Sonntag WE. Insulin-like growth factor-1 ameliorates age-related behavioral deficits. *Neuroscience* 1998; **87**: 559-569 [PMID: 9758223 DOI: 10.1016/S0306-4522(98)00143-2]
- 127 **Gillespie CM**, Merkel AL, Martin AA. Effects of insulin-like growth factor-I and LR3IGF-I on regional blood flow in normal rats. *J Endocrinol* 1997; **155**: 351-358 [PMID: 9415069 DOI: 10.1677/joe.0.1550351]
- 128 **Hölscher C**. Diabetes as a risk factor for Alzheimer's disease: insulin signalling impairment in the brain as an alternative model of Alzheimer's disease. *Biochem Soc Trans* 2011; **39**: 891-897 [PMID: 21787319 DOI: 10.1042/BST0390891]
- 129 **MacKnight C**, Rockwood K, Awalt E, McDowell I. Diabetes mellitus and the risk of dementia, Alzheimer's disease and vascular cognitive impairment in the Canadian Study of Health and Aging. *Dement Geriatr Cogn Disord* 2002; **14**: 77-83 [PMID: 12145454 DOI: 10.1159/000064928]
- 130 **Vitale G**, Brugts MP, Ogliari G, Castaldi D, Fatti LM, Varewijck AJ, Lamberts SW, Monti D, Bucci L, Cevenini E, Cavagnini F, Franceschi C, Hofland LJ, Mari D, Janssen J. Low circulating IGF-I bioactivity is associated with human longevity: findings in centenarians' offspring. *Aging (Albany NY)* 2012; **4**: 580-589 [PMID: 22983440]
- 131 **Barbieri M**, Rizzo MR, Papa M, Boccardi V, Esposito A, White MF, Paolisso G. The IRS2 Gly1057Asp variant is associated with human longevity. *J Gerontol A Biol Sci Med Sci* 2010; **65**: 282-286 [PMID: 19887537 DOI: 10.1093/gerona/glp154]
- 132 **Dillin A**, Cohen E. Ageing and protein aggregation-mediated disorders: from invertebrates to mammals. *Philos Trans R Soc Lond B Biol Sci* 2011; **366**: 94-98 [PMID: 21115535 DOI: 10.1098/rstb.2010.0271]
- 133 **Mosconi L**, Mistur R, Switalski R, Brys M, Glodzik L, Rich K, Pirraglia E, Tsui W, De Santi S, de Leon MJ. Declining brain glucose metabolism in normal individuals with a maternal history of Alzheimer disease. *Neurology* 2009; **72**: 513-520 [PMID: 19005175 DOI: 10.1212/01.wnl.0000333247.51383.43]
- 134 **Zhao L**, Teter B, Morihara T, Lim GP, Ambegaokar SS, Ubeda OJ, Frautschy SA, Cole GM. Insulin-degrading enzyme as a downstream target of insulin receptor signaling cascade: implications for Alzheimer's disease intervention. *J Neurosci* 2004; **24**: 11120-11126 [PMID: 15590928]
- 135 **Steen E**, Terry BM, Rivera EJ, Cannon JL, Neely TR, Tavares R, Xu XJ, Wands JR, de la Monte SM. Impaired insulin

- and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease--is this type 3 diabetes? *J Alzheimers Dis* 2005; **7**: 63-80 [PMID: 15750215]
- 136 **Braak H**, Braak E. Frequency of stages of Alzheimer-related lesions in different age categories. *Neurobiol Aging* 1997; **18**: 351-357 [PMID: 9330961]
 - 137 **Benilova I**, Karran E, De Strooper B. The toxic A β oligomer and Alzheimer's disease: an emperor in need of clothes. *Nat Neurosci* 2012; **15**: 349-357 [PMID: 22286176 DOI: 10.1038/nn.3028]
 - 138 **Haass C**. Initiation and propagation of neurodegeneration. *Nat Med* 2010; **16**: 1201-1204 [PMID: 21052073 DOI: 10.1038/nm.2223]
 - 139 **Franke TF**. PI3K/Akt: getting it right matters. *Oncogene* 2008; **27**: 6473-6488 [PMID: 18955974 DOI: 10.1038/onc.2008.313]
 - 140 **Cohen E**, Bieschke J, Perciavalle RM, Kelly JW, Dillin A. Opposing activities protect against age-onset proteotoxicity. *Science* 2006; **313**: 1604-1610 [PMID: 16902091]
 - 141 **Caccamo A**, Majumder S, Richardson A, Strong R, Oddo S. Molecular interplay between mammalian target of rapamycin (mTOR), amyloid-beta, and Tau: effects on cognitive impairments. *J Biol Chem* 2010; **285**: 13107-13120 [PMID: 20178983 DOI: 10.1074/jbc.M110.100420]
 - 142 **Majumder S**, Caccamo A, Medina DX, Benavides AD, Javors MA, Kraig E, Strong R, Richardson A, Oddo S. Lifelong rapamycin administration ameliorates age-dependent cognitive deficits by reducing IL-1 β and enhancing NMDA signaling. *Aging Cell* 2012; **11**: 326-335 [PMID: 22212527 DOI: 10.1111/j.1474-9726.2011.00791.x]
 - 143 **Killick R**, Scales G, Leroy K, Causevic M, Hooper C, Irvine EE, Choudhury AI, Drinkwater L, Kerr F, Al-Qassab H, Stephenson J, Yilmaz Z, Giese KP, Brion JP, Withers DJ, Lovestone S. Deletion of Irs2 reduces amyloid deposition and rescues behavioural deficits in APP transgenic mice. *Biochem Biophys Res Commun* 2009; **386**: 257-262 [PMID: 19523444 DOI: 10.1016/j.bbrc.2009.06.032]
 - 144 **Harrison DE**, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, Nadon NL, Wilkinson JE, Frenkel K, Carter CS, Pahor M, Javors MA, Fernandez E, Miller RA. Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature* 2009; **460**: 392-395 [PMID: 19587680 DOI: 10.1038/nature08221]
 - 145 **Sweatt JD**. The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory. *J Neurochem* 2001; **76**: 1-10 [PMID: 11145972]
 - 146 **Carro E**, Trejo JL, Gomez-Isla T, LeRoith D, Torres-Aleman I. Serum insulin-like growth factor I regulates brain amyloid-beta levels. *Nat Med* 2002; **8**: 1390-1397 [PMID: 12415260]
 - 147 **Adams MM**, Elizabeth Forbes M, Constance Linville M, Riddle DR, Sonntag WE, Brunso-Bechtold JK. Stability of local brain levels of insulin-like growth factor-I in two well-characterized models of decreased plasma IGF-I. *Growth Factors* 2009; **27**: 181-188 [PMID: 19343576 DOI: 10.1080/08977190902863639]
 - 148 **Westwood AJ**, Beiser A, Decarli C, Harris TB, Chen TC, He XM, Roubenoff R, Pikula A, Au R, Braverman LE, Wolf PA, Vasan RS, Seshadri S. Insulin-like growth factor-1 and risk of Alzheimer dementia and brain atrophy. *Neurology* 2014; **82**: 1613-1619 [PMID: 24706014 DOI: 10.1212/WNL.0000000000000382]
 - 149 **Green CJ**, Holly JM, Bayer A, Fish M, Ebrahim S, Gallacher J, Ben-Shlomo Y. The role of IGF-I, IGF-II, and IGFBP-3 in male cognitive aging and dementia risk: the Caerphilly Prospective Study. *J Alzheimers Dis* 2014; **41**: 867-875 [PMID: 24705546 DOI: 10.3233/JAD-132183]
 - 150 **van Exel E**, Eikelenboom P, Comijs H, Deeg DJ, Stek ML, Westendorp RG. Insulin-like growth factor-1 and risk of late-onset Alzheimer's disease: findings from a family study. *Neurobiol Aging* 2014; **35**: 725.e7-725.10 [PMID: 24054991 DOI: 10.1016/j.neurobiolaging.2013.08.014]
 - 151 **Kopf D**, Frölich L. Risk of incident Alzheimer's disease in diabetic patients: a systematic review of prospective trials. *J Alzheimers Dis* 2009; **16**: 677-685 [PMID: 19387104 DOI: 10.3233/JAD-2009-1011]
 - 152 **Bielsels GJ**, De Leeuw FE, Lindeboom J, Barkhof F, Scheltens P. Increased cortical atrophy in patients with Alzheimer's disease and type 2 diabetes mellitus. *J Neurol Neurosurg Psychiatry* 2006; **77**: 304-307 [PMID: 16484636 DOI: 10.1136/jnnp.2005.06.9583]
 - 153 **Janson J**, Laedtke T, Parisi JE, O'Brien P, Petersen RC, Butler PC. Increased risk of type 2 diabetes in Alzheimer disease. *Diabetes* 2004; **53**: 474-481 [PMID: 14747300]
 - 154 **Yang Y**, Song W. Molecular links between Alzheimer's disease and diabetes mellitus. *Neuroscience* 2013; **250**: 140-150 [PMID: 23867771 DOI: 10.1016/j.neuroscience.2013.07.009]
 - 155 **Osborn O**, Olefsky JM. The cellular and signaling networks linking the immune system and metabolism in disease. *Nat Med* 2012; **18**: 363-374 [PMID: 22395709 DOI: 10.1038/nm.2627]
 - 156 **Akiyama H**, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, Cooper NR, Eikelenboom P, Emmerling M, Fiebich BL, Finch CE, Frautschy S, Griffin WS, Hampel H, Hull M, Landreth G, Lue L, Mrak R, Mackenzie IR, McGeer PL, O'Banion MK, Pachter J, Pasinetti G, Plata-Salaman C, Rogers J, Rydel R, Shen Y, Streit W, Strohmeyer R, Tooyoma I, Van Muiswinkel FL, Veerhuis R, Walker D, Webster S, Wegrzyniak B, Wenk G, Wyss-Coray T. Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000; **21**: 383-421 [PMID: 10858586]
 - 157 **de la Monte SM**, Tong M. Brain metabolic dysfunction at the core of Alzheimer's disease. *Biochem Pharmacol* 2014; **88**: 548-559 [PMID: 24380887 DOI: 10.1016/j.bcp.2013.12.012]
 - 158 **Wyss-Coray T**. Inflammation in Alzheimer disease: driving force, bystander or beneficial response? *Nat Med* 2006; **12**: 1005-1015 [PMID: 16960575]
 - 159 **Hanisch UK**, Kettenmann H. Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 2007; **10**: 1387-1394 [PMID: 17965659]
 - 160 **Ferreira ST**, Clarke JR, Bomfim TR, De Felice FG. Inflammation, defective insulin signaling, and neuronal dysfunction in Alzheimer's disease. *Alzheimers Dement* 2014; **10**: S76-S83 [PMID: 24529528 DOI: 10.1016/j.jalz.2013.12.010]
 - 161 **De Felice FG**. Alzheimer's disease and insulin resistance: translating basic science into clinical applications. *J Clin Invest* 2013; **123**: 531-539 [PMID: 23485579 DOI: 10.1172/JCI64595]
 - 162 **Suh HS**, Zhao ML, Derico L, Choi N, Lee SC. Insulin-like growth factor 1 and 2 (IGF1, IGF2) expression in human microglia: differential regulation by inflammatory mediators. *J Neuroinflammation* 2013; **10**: 37 [PMID: 23497056 DOI: 10.1186/1742-2094-10-37]

P- Reviewer: Cavanna AE, Lee YJ, Lichter T **S- Editor:** Qi Y
L- Editor: A **E- Editor:** Liu SQ



GENERAL INFORMATION

World Journal of Neurology (*World J Neurol*, *WJN*, online ISSN 2218-6212, DOI: 10.5316) is a peer-reviewed open access (OA) academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

Aim and scope

WJN covers topics concerning neuro-oncology, electroneurophysiology, cerebrovascular diseases, epilepsy, cognitive impairment, myopathy and peripheral neuropathy, degenerative diseases, infectious diseases, demyelinating diseases, immunological diseases, genetic/metabolic diseases, affective disorders, headaches, sleep disorders, interventional neuroradiology, minimally invasive therapy, rehabilitation, diagnostic imaging, evidence-based medicine, epidemiology and nursing. The current columns of *WJN* include editorial, frontier, diagnostic advances, therapeutics advances, field of vision, mini-reviews, review, topic highlight, medical ethics, original articles, case report, clinical case conference (Clinicopathological conference), and autobiography. Priority publication will be given to articles concerning diagnosis and treatment of neurological diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJN*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

WJN is edited and published by Baishideng Publishing Group (BPG). BPG has a strong professional editorial team composed of science editors, language editors and electronic editors. BPG currently publishes 43 OA clinical medical journals, including 42 in English, has a total of 15471 editorial board members or peer reviewers, and is a world first-class publisher.

Columns

The columns in the issues of *WJN* will include: (1) Editorial: The editorial board members are invited to make comments on an important topic in their field in terms of its current research status and future directions to lead the development of this discipline; (2) Frontier: The editorial board members are invited to select a highly cited cutting-edge original paper of his/her own to summarize major findings, the problems that have been resolved and remain to be resolved, and future research directions to help readers understand his/her important academic point of view and future research directions in the field; (3) Diagnostic Advances: The editorial board members are invited to write high-quality diagnostic advances in their field to improve the diagnostic skills of readers. The topic covers general clinical diagnosis, differential diagnosis, pathological diagnosis, laboratory diagnosis, imaging diagnosis, endoscopic diagnosis, biotechnological diagnosis, functional diagnosis, and physical diagnosis; (4) Therapeutics Advances: The editorial board members are invited to write high-quality therapeutic advances in their field to help improve the therapeutic skills of readers. The topic covers medication therapy, psychotherapy, physical therapy, replacement therapy, interventional therapy, minimally invasive therapy, endo-

scopic therapy, transplantation therapy, and surgical therapy; (5) Field of Vision: The editorial board members are invited to write commentaries on classic articles, hot topic articles, or latest articles to keep readers at the forefront of research and increase their levels of clinical research. Classic articles refer to papers that are included in Web of Knowledge and have received a large number of citations (ranking in the top 1%) after being published for more than years, reflecting the quality and impact of papers. Hot topic articles refer to papers that are included in Web of Knowledge and have received a large number of citations after being published for no more than 2 years, reflecting cutting-edge trends in scientific research. Latest articles refer to the latest published high-quality papers that are included in PubMed, reflecting the latest research trends. These commentary articles should focus on the status quo of research, the most important research topics, the problems that have now been resolved and remain to be resolved, and future research directions. Basic information about the article to be commented (including authors, article title, journal name, year, volume, and inclusive page numbers); (6) Minireviews: The editorial board members are invited to write short reviews on recent advances and trends in research of molecular biology, genomics, and related cutting-edge technologies to provide readers with the latest knowledge and help improve their diagnostic and therapeutic skills; (7) Review: To make a systematic review to focus on the status quo of research, the most important research topics, the problems that have now been resolved and remain to be resolved, and future research directions; (8) Topic Highlight: The editorial board members are invited to write a series of articles (7-10 articles) to comment and discuss a hot topic to help improve the diagnostic and therapeutic skills of readers; (9) Medical Ethics: The editorial board members are invited to write articles about medical ethics to increase readers' knowledge of medical ethics. The topic covers international ethics guidelines, animal studies, clinical trials, organ transplantation, etc.; (10) Clinical Case Conference or Clinicopathological Conference: The editorial board members are invited to contribute high-quality clinical case conference; (11) Original Articles: To report innovative and original findings in neurology; (12) Research Report: To briefly report the novel and innovative findings in neurology; (13) Meta-Analysis: To summarize a given quantitative effect, e.g., the clinical effectiveness and safety of clinical treatments by combining data from two or more randomized controlled trials, thereby providing more precise and externally valid estimates than those which would stem from each individual dataset if analyzed separately from the others; (14) Case Report: To report a rare or typical case; (15) Letters to the Editor: To discuss and make reply to the contributions published in *WJN*, or to introduce and comment on a controversial issue of general interest; (16) Book Reviews: To introduce and comment on quality monographs of neurology; and (17) Autobiography: The editorial board members are invited to write their autobiography to provide readers with stories of success or failure in their scientific research career. The topic covers their basic personal information and information about when they started doing research work, where and how they did research work, what they have achieved, and their lessons from success or failure.

Name of journal

World Journal of Neurology

ISSN

ISSN 2218-6212 (online)

Instructions to authors

Launch date

December 28, 2011

Frequency

Quarterly

Editors-in-Chief

Felipe Fregni, MD, PhD, MPH, Associate Professor, Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Harvard Medical School, Room 725, 125 Nashua St. Boston, MA 02114, United States

Vincenzo Solfrizzi, MD, PhD, Professor, Department of Internal Medicine, Immunology and Infectious Diseases, Section of Geriatric Medicine-Memory Unit, University of Bari, P.zza G. Cesare, 11, 70124 Bari, Italy

Editorial office

Jin-Lei Wang, Director
Xiu-Xia Song, Vice Director
World Journal of Neurology
Room 903, Building D, Ocean International Center,
No. 62 Dongsihuan Zhonglu, Chaoyang District,
Beijing 100025, China
Telephone: +86-10-59080039
Fax: +86-10-85381893
E-mail: editorialoffice@wjnet.com
Help Desk: <http://www.wjnet.com/esps/helpdesk.aspx>
<http://www.wjnet.com>

Publisher

Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjnet.com
Help Desk: <http://www.wjnet.com/esps/helpdesk.aspx>
<http://www.wjnet.com>

Instructions to authors

Full instructions are available online at http://www.wjnet.com/2218-6212/g_info_20100722173918.htm.

Indexed and Abstracted in

Digital Object Identifier.

SPECIAL STATEMENT

All articles published in journals owned by the BPG represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

Biostatistical editing

Statistical review is performed after peer review. We invite an expert in Biomedical Statistics to evaluate the statistical method used in the paper, including *t*-test (group or paired comparisons), chi-squared test, Redit, probit, logit, regression (linear, curvilinear, or stepwise), correlation, analysis of variance, analysis of covariance, *etc.* The reviewing points include: (1) Statistical methods should be described when they are used to verify the results; (2) Whether the statistical techniques are suitable or correct; (3) Only homogeneous data can be averaged. Standard deviations are preferred to standard errors. Give the number of observations and subjects (*n*). Losses in observations, such as drop-outs from the study should be reported; (4) Values such as ED50, LD50, IC50 should have their 95% confidence limits calculated and compared by weighted probit analysis (Bliss and Finney); and (5) The word 'significantly' should be replaced by its synonyms (if it indicates extent) or the *P* value (if it indicates statistical significance).

Conflict-of-interest statement

In the interests of transparency and to help reviewers assess any potential bias, *WJN* requires authors of all papers to declare any competing commercial, personal, political, intellectual, or religious interests in relation to the submitted work. Referees are also asked to indicate any potential conflict they might have reviewing a particular paper. Before submitting, authors are suggested to read "Uniform Requirements for Manuscripts Submitted to Biomedical Journals: Ethical Considerations in the Conduct and Reporting of Research: Conflicts of Interest" from International Committee of Medical Journal Editors (ICMJE), which is available at: http://www.icmje.org/ethical_4conflicts.html.

Sample wording: [Name of individual] has received fees for serving as a speaker, a consultant and an advisory board member for [names of organizations], and has received research funding from [names of organization]. [Name of individual] is an employee of [name of organization]. [Name of individual] owns stocks and shares in [name of organization]. [Name of individual] owns patent [patent identification and brief description].

Statement of informed consent

Manuscripts should contain a statement to the effect that all human studies have been reviewed by the appropriate ethics committee or it should be stated clearly in the text that all persons gave their informed consent prior to their inclusion in the study. Details that might disclose the identity of the subjects under study should be omitted. Authors should also draw attention to the Code of Ethics of the World Medical Association (Declaration of Helsinki, 1964, as revised in 2004).

Statement of human and animal rights

When reporting the results from experiments, authors should follow the highest standards and the trial should conform to Good Clinical Practice (for example, US Food and Drug Administration Good Clinical Practice in FDA-Regulated Clinical Trials; UK Medicines Research Council Guidelines for Good Clinical Practice in Clinical Trials) and/or the World Medical Association Declaration of Helsinki. Generally, we suggest authors follow the lead investigator's national standard. If doubt exists whether the research was conducted in accordance with the above standards, the authors must explain the rationale for their approach and demonstrate that the institutional review body explicitly approved the doubtful aspects of the study.

Before submitting, authors should make their study approved by the relevant research ethics committee or institutional review board. If human participants were involved, manuscripts must be accompanied by a statement that the experiments were undertaken with the understanding and appropriate informed consent of each. Any personal item or information will not be published without explicit consents from the involved patients. If experimental animals were used, the materials and methods (experimental procedures) section must clearly indicate that appropriate measures were taken to minimize pain or discomfort, and details of animal care should be provided.

SUBMISSION OF MANUSCRIPTS

Manuscripts should be typed in 1.5 line spacing and 12 pt. Book Antiqua with ample margins. Number all pages consecutively, and start each of the following sections on a new page: Title Page, Abstract, Introduction, Materials and Methods, Results, Discussion, Acknowledgements, References, Tables, Figures, and Figure Legends. Neither the editors nor the publisher are responsible for the opinions expressed by contributors. Manuscripts formally accepted for publication become the permanent property of BPG, and may not be reproduced by any means, in whole or in part, without the written permission of both the authors and the publisher. We reserve the right to copy-edit and put onto our website accepted manuscripts. Authors should follow the relevant guidelines for the care and use of laboratory animals of their institution or national animal welfare committee. For the sake of transparency in regard to the performance and reporting of clinical trials, we endorse the policy of the ICMJE to refuse to publish papers on clinical trial results if the trial was not recorded in a publicly-accessible registry.

at its outset. The only register now available, to our knowledge, is <http://www.clinicaltrials.gov> sponsored by the United States National Library of Medicine and we encourage all potential contributors to register with it. However, in the case that other registers become available you will be duly notified. A letter of recommendation from each author's organization should be provided with the contributed article to ensure the privacy and secrecy of research is protected.

Authors should retain one copy of the text, tables, photographs and illustrations because rejected manuscripts will not be returned to the author(s) and the editors will not be responsible for loss or damage to photographs and illustrations sustained during mailing.

Online submissions

Manuscripts should be submitted through the Online Submission System at: <http://www.wjgnet.com/esp/>. Authors are highly recommended to consult the ONLINE INSTRUCTIONS TO AUTHORS (http://www.wjgnet.com/2218-6212/g_info_20100722173918.htm) before attempting to submit online. For assistance, authors encountering problems with the Online Submission System may send an email describing the problem to bpgoffice@wjgnet.com, or by telephone: +86-10-85381892. If you submit your manuscript online, do not make a postal contribution. Repeated online submission for the same manuscript is strictly prohibited.

MANUSCRIPT PREPARATION

All contributions should be written in English. All articles must be submitted using word-processing software. All submissions must be typed in 1.5 line spacing and 12 pt. Book Antiqua with ample margins. Style should conform to our house format. Required information for each of the manuscript sections is as follows:

Title page

Title: Title should be less than 12 words.

Running title: A short running title of less than 6 words should be provided.

Authorship: Authorship credit should be in accordance with the standard proposed by ICMJE, based on (1) substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; (2) drafting the article or revising it critically for important intellectual content; and (3) final approval of the version to be published. Authors should meet conditions 1, 2, and 3.

Institution: Author names should be given first, then the complete name of institution, city, province and postcode. For example, Xu-Chen Zhang, Li-Xin Mei, Department of Pathology, Chengde Medical College, Chengde 067000, Hebei Province, China. One author may be represented from two institutions, for example, George Sgourakis, Department of General, Visceral, and Transplantation Surgery, Essen 45122, Germany; George Sgourakis, 2nd Surgical Department, Korgialenio-Benakio Red Cross Hospital, Athens 15451, Greece

Author contributions: The format of this section should be: Author contributions: Wang CL and Liang L contributed equally to this work; Wang CL, Liang L, Fu JF, Zou CC, Hong F and Wu XM designed the research; Wang CL, Zou CC, Hong F and Wu XM performed the research; Xue JZ and Lu JR contributed new reagents/analytic tools; Wang CL, Liang L and Fu JF analyzed the data; and Wang CL, Liang L and Fu JF wrote the paper.

Supportive foundations: The complete name and number of supportive foundations should be provided, e.g., Supported by National Natural Science Foundation of China, No. 30224801

Correspondence to: Only one corresponding address should be

provided. Author names should be given first, then author title, affiliation, the complete name of institution, city, postcode, province, country, and email. All the letters in the email should be in lower case. A space interval should be inserted between country name and email address. For example, Montgomery Bissell, MD, Professor of Medicine, Chief, Liver Center, Gastroenterology Division, University of California, Box 0538, San Francisco, CA 94143, United States. montgomerybissell@ucsf.edu

Telephone and fax: Telephone and fax should consist of +, country number, district number and telephone or fax number, e.g., Telephone: +86-10-85381892 Fax: +86-10-85381893

Peer reviewers: All articles received are subject to peer review. Normally, three experts are invited for each article. Decision on acceptance is made only when at least two experts recommend publication of an article. All peer-reviewers are acknowledged on Express Submission and Peer-review System website.

Abstract

There are unstructured abstracts (no less than 200 words) and structured abstracts. The specific requirements for structured abstracts are as follows:

An informative, structured abstract should accompany each manuscript. Abstracts of original contributions should be structured into the following sections: AIM (no more than 20 words; Only the purpose of the study should be included. Please write the Aim in the form of "To investigate/study/..."), METHODS (no less than 140 words for Original Articles; and no less than 80 words for Brief Articles), RESULTS (no less than 150 words for Original Articles and no less than 120 words for Brief Articles; You should present *P* values where appropriate and must provide relevant data to illustrate how they were obtained, e.g., 6.92 ± 3.86 vs 3.61 ± 1.67 , $P < 0.001$), and CONCLUSION (no more than 26 words).

Key words

Please list 5-10 key words, selected mainly from *Index Medicus*, which reflect the content of the study.

Core tip

Please write a summary of less than 100 words to outline the most innovative and important arguments and core contents in your paper to attract readers.

Text

For articles of these sections, original articles and brief articles, the main text should be structured into the following sections: INTRODUCTION, MATERIALS AND METHODS, RESULTS and DISCUSSION, and should include appropriate Figures and Tables. Data should be presented in the main text or in Figures and Tables, but not in both.

Illustrations

Figures should be numbered as 1, 2, 3, etc., and mentioned clearly in the main text. Provide a brief title for each figure on a separate page. Detailed legends should not be provided under the figures. This part should be added into the text where the figures are applicable. Keeping all elements compiled is necessary in line-art image. Scale bars should be used rather than magnification factors, with the length of the bar defined in the legend rather than on the bar itself. File names should identify the figure and panel. Avoid layering type directly over shaded or textured areas. Please use uniform legends for the same subjects. For example: Figure 1 Pathological changes in atrophic gastritis after treatment. A: ...; B: ...; C: ...; D: ...; E: ...; F: ...; G: ...etc. It is our principle to publish high resolution-figures for the E-versions.

Tables

Three-line tables should be numbered 1, 2, 3, etc., and mentioned clearly in the main text. Provide a brief title for each table. Detailed legends should not be included under tables, but rather added into

Instructions to authors

the text where applicable. The information should complement, but not duplicate the text. Use one horizontal line under the title, a second under column heads, and a third below the Table, above any footnotes. Vertical and italic lines should be omitted.

Notes in tables and illustrations

Data that are not statistically significant should not be noted. ^a $P < 0.05$, ^b $P < 0.01$ should be noted ($P > 0.05$ should not be noted). If there are other series of P values, ^c $P < 0.05$ and ^d $P < 0.01$ are used. A third series of P values can be expressed as ^e $P < 0.05$ and ^f $P < 0.01$. Other notes in tables or under illustrations should be expressed as ¹F, ²F, ³F; or sometimes as other symbols with a superscript (Arabic numerals) in the upper left corner. In a multi-curve illustration, each curve should be labeled with ●, ○, ■, □, ▲, △, etc., in a certain sequence.

Acknowledgments

Brief acknowledgments of persons who have made genuine contributions to the manuscript and who endorse the data and conclusions should be included. Authors are responsible for obtaining written permission to use any copyrighted text and/or illustrations.

REFERENCES

Coding system

The author should number the references in Arabic numerals according to the citation order in the text. Put reference numbers in square brackets in superscript at the end of citation content or after the cited author's name. For citation content which is part of the narration, the coding number and square brackets should be typeset normally. For example, "Crohn's disease (CD) is associated with increased intestinal permeability^[12]". If references are cited directly in the text, they should be put together within the text, for example, "From references^[19,22-24], we know that..."

When the authors write the references, please ensure that the order in text is the same as in the references section, and also ensure the spelling accuracy of the first author's name. Do not list the same citation twice.

PMID and DOI

Please provide PubMed citation numbers to the reference list, e.g., PMID and DOI, which can be found at <http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed> and <http://www.crossref.org/SimpleTextQuery/>, respectively. The numbers will be used in E-version of this journal.

Style for journal references

Authors: the name of the first author should be typed in bold-faced letters. The family name of all authors should be typed with the initial letter capitalized, followed by their abbreviated first and middle initials. (For example, Lian-Sheng Ma is abbreviated as Ma LS, Bo-Rong Pan as Pan BR). The title of the cited article and italicized journal title (journal title should be in its abbreviated form as shown in PubMed), publication date, volume number (in black), start page, and end page [PMID: 11819634 DOI: 10.3748/wjg.13.5396].

Style for book references

Authors: the name of the first author should be typed in bold-faced letters. The surname of all authors should be typed with the initial letter capitalized, followed by their abbreviated middle and first initials. (For example, Lian-Sheng Ma is abbreviated as Ma LS, Bo-Rong Pan as Pan BR). Book title. Publication number. Publication place: Publication press, Year: start page and end page.

Format

Journals

English journal article (list all authors and include the PMID where applicable)

- 1 **Jung EM**, Clevert DA, Schreyer AG, Schmitt S, Rennett J, Kubale R, Feuerbach S, Jung F. Evaluation of quantitative contrast harmonic imaging to assess malignancy of liver tumors: A prospective controlled two-center study. *World J Gastroenterol* 2007; **13**: 6356-6364 [PMID: 18081224 DOI: 10.3748/wjg.13.6356]

Chinese journal article (list all authors and include the PMID where applicable)

- 2 **Lin GZ**, Wang XZ, Wang P, Lin J, Yang FD. Immunologic

effect of Jianpi Yishen decoction in treatment of Pixu-diarhoea. *Shijie Huaren Xiaobua Zazhi* 1999; **7**: 285-287

In press

- 3 **Tian D**, Araki H, Stahl E, Bergelson J, Kreitman M. Signature of balancing selection in Arabidopsis. *Proc Natl Acad Sci USA* 2006; In press

Organization as author

- 4 **Diabetes Prevention Program Research Group**. Hypertension, insulin, and proinsulin in participants with impaired glucose tolerance. *Hypertension* 2002; **40**: 679-686 [PMID: 12411462 PMID:2516377 DOI:10.1161/01.HYP.0000035706.28494.09]

Both personal authors and an organization as author

- 5 **Vallancien G**, Emberton M, Harving N, van Moorselaar RJ; Alf-One Study Group. Sexual dysfunction in 1, 274 European men suffering from lower urinary tract symptoms. *J Urol* 2003; **169**: 2257-2261 [PMID: 12771764 DOI:10.1097/01.ju.0000067940.76090.73]

No author given

- 6 21st century heart solution may have a sting in the tail. *BMJ* 2002; **325**: 184 [PMID: 12142303 DOI:10.1136/bmj.325.7357.184]

Volume with supplement

- 7 **Geraud G**, Spierings EL, Keywood C. Tolerability and safety of frovatriptan with short- and long-term use for treatment of migraine and in comparison with sumatriptan. *Headache* 2002; **42** Suppl 2: S93-99 [PMID: 12028325 DOI:10.1046/j.1526-4610.42.s2.7.x]

Issue with no volume

- 8 **Banit DM**, Kaufer H, Hartford JM. Intraoperative frozen section analysis in revision total joint arthroplasty. *Clin Orthop Relat Res* 2002; **(401)**: 230-238 [PMID: 12151900 DOI:10.1097/00003086-200208000-00026]

No volume or issue

- 9 Outreach: Bringing HIV-positive individuals into care. *HRS-A Careaction* 2002; 1-6 [PMID: 12154804]

Books

Personal author(s)

- 10 **Sherlock S**, Dooley J. Diseases of the liver and biliary system. 9th ed. Oxford: Blackwell Sci Pub, 1993: 258-296

Chapter in a book (list all authors)

- 11 **Lam SK**. Academic investigator's perspectives of medical treatment for peptic ulcer. In: Swabb EA, Azabo S. Ulcer disease: investigation and basis for therapy. New York: Marcel Dekker, 1991: 431-450

Author(s) and editor(s)

- 12 **Breedlove GK**, Schorheide AM. Adolescent pregnancy. 2nd ed. Wiczorek RR, editor. White Plains (NY): March of Dimes Education Services, 2001: 20-34

Conference proceedings

- 13 **Harnden P**, Joffe JK, Jones WG, editors. Germ cell tumours V. Proceedings of the 5th Germ cell tumours Conference; 2001 Sep 13-15; Leeds, UK. New York: Springer, 2002: 30-56

Conference paper

- 14 **Christensen S**, Oppacher F. An analysis of Koza's computational effort statistic for genetic programming. In: Foster JA, Lutton E, Miller J, Ryan C, Tettamanzi AG, editors. Genetic programming. EuroGP 2002: Proceedings of the 5th European Conference on Genetic Programming; 2002 Apr 3-5; Kinsdale, Ireland. Berlin: Springer, 2002: 182-191

Electronic journal (list all authors)

- 15 Morse SS. Factors in the emergence of infectious diseases. Emerg Infect Dis serial online, 1995-01-03, cited 1996-06-05; 1(1): 24 screens. Available from: URL: <http://www.cdc.gov/ncidod/eid/index.htm>

Patent (list all authors)

- 16 **Pagedas AC**, inventor; Ancel Surgical R&D Inc., assignee. Flexible endoscopic grasping and cutting device and positioning tool assembly. United States patent US 20020103498.

2002 Aug 1

Statistical dataWrite as mean \pm SD or mean \pm SE.**Statistical expression**

Express *t* test as *t* (in italics), *F* test as *F* (in italics), chi square test as χ^2 (in Greek), related coefficient as *r* (in italics), degree of freedom as *v* (in Greek), sample number as *n* (in italics), and probability as *P* (in italics).

Units

Use SI units. For example: body mass, *m* (B) = 78 kg; blood pressure, *p* (B) = 16.2/12.3 kPa; incubation time, *t* (incubation) = 96 h; blood glucose concentration, *c* (glucose) 6.4 \pm 2.1 mmol/L; blood CEA mass concentration, *p* (CEA) = 8.6 24.5 μ g/L; CO₂ volume fraction, 50 mL/L CO₂, not 5% CO₂; likewise for 40 g/L formaldehyde, not 10% formalin; and mass fraction, 8 ng/g, *etc.* Arabic numerals such as 23, 243, 641 should be read 23 243 641.

The format for how to accurately write common units and quantum numbers can be found at: http://www.wjgnet.com/2218-6212/g_info_20100723222255.htm.

Abbreviations

Standard abbreviations should be defined in the abstract and on first mention in the text. In general, terms should not be abbreviated unless they are used repeatedly and the abbreviation is helpful to the reader. Permissible abbreviations are listed in Units, Symbols and Abbreviations: A Guide for Biological and Medical Editors and Authors (Ed. Baron DN, 1988) published by The Royal Society of Medicine, London. Certain commonly used abbreviations, such as DNA, RNA, HIV, LD50, PCR, HBV, ECG, WBC, RBC, CT, ESR, CSF, IgG, ELISA, PBS, ATP, EDTA, mAb, can be used directly without further explanation.

Italics

Quantities: *t* time or temperature, *c* concentration, *A* area, *l* length, *m* mass, *V* volume.

Genotypes: *gyrA*, *arg 1*, *c myc*, *c fos*, *etc.*

Restriction enzymes: *EcoRI*, *HindI*, *BamHI*, *Kho I*, *Kpn I*, *etc.*

Biology: *H. pylori*, *E. coli*, *etc.*

Examples for paper writing

All types of articles' writing style and requirement will be found in the link: <http://www.wjgnet.com/esps/NavigationInfo.aspx?id=15>

RESUBMISSION OF THE REVISED MANUSCRIPTS

Authors must revise their manuscript carefully according to the

revision policies of BPG. The revised version, along with the signed copyright transfer agreement, responses to the reviewers, and English language Grade A certificate (for non-native speakers of English), should be submitted to the online system *via* the link contained in the e-mail sent by the editor. If you have any questions about the revision, please send e-mail to esps@wjgnet.com.

Language evaluation

The language of a manuscript will be graded before it is sent for revision. (1) Grade A: priority publishing; (2) Grade B: minor language polishing; (3) Grade C: a great deal of language polishing needed; and (4) Grade D: rejected. Revised articles should reach Grade A.

Copyright assignment form

Please download a Copyright assignment form from http://www.wjgnet.com/2218-6212/g_info_20100723222139.htm.

Responses to reviewers

Please revise your article according to the comments/suggestions provided by the reviewers. The format for responses to the reviewers' comments can be found at: http://www.wjgnet.com/2218-6212/g_info_20100723222041.htm.

Proof of financial support

For papers supported by a foundation, authors should provide a copy of the approval document and serial number of the foundation.

STATEMENT ABOUT ANONYMOUS PUBLICATION OF THE PEER REVIEWERS' COMMENTS

In order to increase the quality of peer review, push authors to carefully revise their manuscripts based on the peer reviewers' comments, and promote academic interactions among peer reviewers, authors and readers, we decide to anonymously publish the reviewers' comments and author's responses at the same time the manuscript is published online.

PUBLICATION FEE

WJN is an international, peer-reviewed, OA online journal. Articles published by this journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium and format, provided the original work is properly cited. The use is non-commercial and is otherwise in compliance with the license. Authors of accepted articles must pay a publication fee. Publication fee: 698 USD per article. All invited articles are published free of charge.



Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

