

World Journal of *Methodology*

World J Methodol 2015 March 26; 5(1): 1-9





Editorial Board

2011-2015

The *World Journal of Methodology* Editorial Board consists of 322 members, representing a team of worldwide experts in methodology. They are from 45 countries, including Argentina (4), Australia (11), Austria (3), Belgium (3), Bosnia and Herzegovina (1), Brazil (4), Canada (12), China (39), Croatia (1), Cuba (1), Czech Republic (4), Denmark (2), Egypt (1), France (8), Germany (5), Greece (6), Hungary (3), India (9), Iran (3), Israel (1), Italy (25), Japan (14), Lithuania (1), Malaysia (1), Mexico (4), Netherlands (3), New Zealand (1), Norway (2), Pakistan (2), Poland (2), Portugal (3), Romania (5), Russia (2), Senegal (1), Singapore (1), South Africa (1), South Korea (4), Spain (18), Sweden (2), Thailand (3), Turkey (4), United Arab Emirates (1), United Kingdom (14), United States (86), and Uruguay (1).

EDITOR-IN-CHIEF

Yicheng Ni, *Leuven*

STRATEGY ASSOCIATE

EDITORS-IN-CHIEF

António Vaz Carneiro, *Lisboa*
Guido Gainotti, *Rome*
Val J GebSKI, *Sydney*
Bo Hang, *Berkeley*
George A Kelley, *Morgantown*
Sang-Soo Lee, *Chuncheon*
Gerhard Litscher, *Graz*
Laurentiu M Popescu, *Bucharest*

GUEST EDITORIAL BOARD MEMBERS

Wen-Hsiung Chan, *Chung Li*
Long-Sen Chang, *Kaohsiung*
Yuh-Shan Ho, *Wufeng*
Shih-Chang Lin, *Taipei*
Hung-Jen Liu, *Taichung*
Ko-Huang Lue, *Taichung*
Rong-Jong Wai, *Chung Li*
Chin-Tsan Wang, *I Lan*
Yau-Huei Wei, *Taipei*
Ching-Feng Weng, *Hualien*

MEMBERS OF THE EDITORIAL BOARD



Argentina

Paula Abate, *Córdoba*
José Miguel Belizán, *Buenos Aires*
Enrique Roberto Soriano, *Buenos Aires*
Rodolfo G Wuilloud, *Mendoza*



Australia

Felix Acker, *Melbourne*
Seetal Dodd, *Geelong*
Guy D Eslick, *Sydney*
Adrian J Gibbs, *Canberra*
Phillipa Jane Hay, *Sydney*
Moyez Jiwa, *Bentley*
Sanjay Patole, *Perth*
Clive Julian Christie Phillips, *Gatton*
Shuhong Wang, *Adelaide*
Jiake Xu, *Perth*



Austria

Gerwin A Bernhardt, *Graz*
Martin Voracek, *Vienna*



Belgium

Zeger Debyser, *Leuven*
Piet K Vanhoenacker, *Aalst*



Bosnia and Herzegovina

Abdülhamit Subasi, *Sarajevo*



Brazil

Monica L Andersen, *São Paulo*
Mariana de Andrea Hacker, *Rio de Janeiro*
Delfim Soares Júnior, *Juiz de Fora*
Moacyr A Rebello, *Rio de Janeiro*



Canada

Ahmed M Abou-Setta, *Edmonton*
Amir Azarpazhooh, *Toronto*
Kenneth R Chapman, *Toronto*
Elijah Dixon, *Calgary*
Martin A Katzman, *Toronto*
Alejandro Lazo-Langner, *London*
Richard WJ Neufeld, *London*
Louis Perrault, *Montreal*
Prakesh S Shah, *Toronto*
Léon C van Kempen, *Montreal*
Yuzhuo Wang, *Vancouver*
Haishan Zeng, *Vancouver*



China

Deng-Feng Cao, *Beijing*
Gilbert Y S Chan, *Hong Kong*
George G Chen, *Hong Kong*
William CS Cho, *Hong Kong*
Raymond Chuen-Chung, *Hong Kong*
Meng-Jie Dong, *Hangzhou*
Zhi-Sheng Duan, *Beijing*
Hani El-Nezami, *Hong Kong*
Rajiv Kumar Jha, *Xi'an*
Gang Jin, *Beijing*
Huang-Xian Ju, *Nanjing*
Hui Li, *Zhejiang*
Yun-Feng Lin, *Chengdu*
Wing-Yee Lui, *Hong Kong*
Feng-Ming Luo, *Chengdu*
Jing-Yun Ma, *Beijing*
Hong-Xiang Sun, *Hangzhou*
Ji-Bo Wang, *Shenyang*
Zhi-Ming Wang, *Chengdu*

Tong-Wen Xu, *Hefei*
 Shi-Ying Xuan, *Qingdao*
 Xi-Lin Yang, *Hong Kong*
 Bang-Ce Ye, *Shanghai*
 Wen-Wu Yu, *Nanjing*
 Yue-Hong Zhang, *Hangzhou*
 Zhong-Ying Zhao, *Hong Kong*
 Chun-Fu Zheng, *Wuhan*
 Ma Zheng, *Beijing*
 Jun-Jie Zhu, *Nanjing*



Croatia

Marijeta Kralj, *Zagreb*



Cuba

Mariano R Ricard, *Habana*



Czech Republic

Kamil Kuca, *Hradec Kralove*
 Bozena Novotna, *Prague*
 Jiri Sedy, *Prague*
 Miroslav Sip, *Ceske Budejovice*



Denmark

Morten Mørup, *Lyngby*
 Hans Sanderson, *Roskilde*



Egypt

Nervana S Bayoumi, *Cairo*



France

Marc Y Bardou, *Dijon*
 Mohammed M Bettahar, *Nancy*
 Olivier David, *Grenoble*
 Guido Kroemer, *Paris*
 Florian Lesage, *Sophia Antipolis*
 Patrick Maison, *Creteil*
 Sandrine Marquet, *Marseille*
 Jean Yves Rotge, *Bordeaux*



Germany

Harald Hampel, *Frankfurt*
 Frank Peinemann, *Cologne*
 M Lienhard Schmitz, *Giessen*
 Alfons Schnitzler, *Duesseldorf*
 Frank Werner, *Magdeburg*



Greece

Konstantinos P Economopoulos, *Athens*
 Georgios A Koumantakis, *Aegion*
 Michael Koutsilieris, *Athens*
 Demosthenes Panagiotakos, *Athens*
 Issidora Papassideri, *Athens*
 Falaras Polycarpus, *Pallini Attikis*



Hungary

Péter Halász, *Budapest*
 András Komócsi, *Pécs*
 László Vécsei, *Szeged*



India

Dipshikha Chakravorty, *Bangalore*
 DK Dhawan, *Chandigarh*
 R Jayakumar, *Cochin*
 Abdul Viqar Khan, *Aligarh*
 Geetha Manivasagam, *Vellore*
 Jacob Peedicayil, *Vellore*
 YS Prabhakar, *Lucknow*
 Debasish Sarkar, *Orissa*
 Rakesh Kumar Sinha, *Ranchi*



Iran

Mehran Javanbakht, *Tehran*
 Enayat Kalantar, *Sanandaj*
 Shekoufeh Nikfar, *Tehran*



Israel

Dan Frenkel, *Tel Aviv*



Italy

Giuseppe Biondi-Zoccai, *Latina*
 Carlo Bonanno, *Vicenza*
 Paolo Borrione, *Turin*
 Filippo Cademartiri, *Monastier di Treviso*
 Alberto Chiesa, *Bologna*
 Annamaria Cimini, *L'Aquila*
 Giovanni Di Leo, *San Donato Milanese*
 Rosario Donato, *Via del Giochetto*
 Alfio Ferlito, *Udine*
 Giovanna Ferraioli, *Milan*
 Irene Floriani, *Milan*
 Landoni Giovanni, *Milano*
 Stefano Girotti, *Bologna*
 Paola Irato, *Padova*
 Giovanni Martinotti, *Rome*
 Mario Mascalchi, *Florence*
 Patrizia Mecocci, *Perugia*
 Germano Orrù, *Cagliari*
 Maurizio Pompili, *Rome*
 Carlo Riccardi, *Perugia*
 Domenico Rubello, *Rovigo*
 Gianfranco Spalletta, *Rome*
 Gambardella Stefano, *Rome*
 Mauro Valtieri, *Rome*



Japan

Kohei Akazawa, *Niigata*
 Subash CB Gopinath, *Tsukuba*
 Masafumi Goto, *Miyagi*
 Koichi Hattori, *Tokyo*
 Satoshi Hirohata, *Okayama*
 Yukihiko Ikeda, *Osaka-sayama*

Masahiro Kohzuki, *Sendai*
 Yoshinori Marunaka, *Kyoto*
 Kenji Miura, *Tokorozawa*
 Ryuichi Morishita, *Suita*
 Mitsuhiro Noda, *Tokyo*
 Yurai Okaji, *Tokyo*
 Hirosato Seki, *Osaka*
 Hisanori Umehara, *Kahoku-gun*



Lithuania

Giedrius Barauskas, *Kaunas*



Malaysia

Iis Sopyan, *Kuala Lumpur*



Mexico

Javier Camacho, *Mexico City*
 Mejía Aranguré Juan Manuel, *Col Doctores*
 Martha Rodríguez-Moran, *Durango*
 Julio Sotelo, *Mexico City*



Netherlands

Kristien Hens, *Maastricht*
 Bart J Polder, *Emmeloord*
 Frank Twisk, *Limmen*



New Zealand

Valery Feigin, *Auckland*



Norway

David F Mota, *Oslo*
 Tore Syversen, *Trondheim*



Pakistan

Muhammad A Noor, *Islamabad*
 Yasir Waheed, *Islamabad*



Poland

Piotr Dziegiel, *Wroclaw*
 Tadeusz Robak, *Lodz*



Portugal

Nuno Lunet, *Porto*
 Hugo Sousa, *Porto*



Romania

Elena Moldoveanu, *Bucharest*
 Monica Neagu, *Bucharest*
 Florin-Dan Popescu, *Bucharest*

Eugen Rusu, *Galati*



Russia

Galina B Bolshakova, *Moscow*
Sergey V Dorozhkin, *Moscow*



Senegal

Badara Cissé, *Dakar*



Singapore

Zhang Yong, *Singapore*



South Africa

Robin Alexander Emsley, *Cape Town*



South Korea

Sang Soo Hah, *Seoul*
Chang-Yong Lee, *Kongju*
Kwan Sik Lee, *Seoul*



Spain

Salvador F Aliño, *Valencia*
Mohamed Farouk Allam, *Cordoba*
Alejandro Cifuentes, *Madrid*
Miren Lopez de Alda, *Barcelona*
Joaquin de Haro, *Madrid*
M de la Guardia, *Valencia*
Emma Garcia-Meca, *Cartagena*
Mónica H Giménez, *Zaragoza*
Josep M Guerrero, *Barcelona*
Fernando Marin, *Madrid*
José A Orosa, *A Coruña*
Jesús Osada, *Zaragoza*
Soledad Rubio, *Córdoba*
Helmut Schröder, *Barcelona*
Jesus Simal-Gandara, *Ourense*
Bahi Takkouche, *Santiago de Compostela*
Gabriela Topa, *Madrid*
Miguel A Vallejo, *Madrid*



Sweden

Stefan Karlsson, *Lund*
Jenny Selander, *Stockholm*



Thailand

Amporn Jariyapongskul, *Bangkok*

Apiwat Mutirangura, *Bangkok*
Bungorn Sripanidkulchai, *Khon Kaen*



Turkey

Mehmet Doğan, *Çağış-Balikesir*
Ferda E Percin, *Ankara*
Ahmet Yildirim, *Bornova-Izmir*
Avsegiul Yildiz, *Izmir*



United Arab Emirates

Hassib Narchi, *Al Ain*



United Kingdom

Richard H Barton, *London*
Paul Evans, *London*
Giuseppe Garcea, *Leicester*
Marta I Garrido, *London*
Sinead Keeney, *Belfast*
Maurice J O'Kane, *Londonderry*
Abdullah Pandor, *Sheffield*
Susan Pang, *Teddington*
Pankaj Sharma, *London*
Andrew Harvey Sims, *Edinburgh*
David E Whitworth, *Aberystwyth*
Sorrel E Wolowacz, *Manchester*
Feng Wu, *Headington*
Shangming Zhou, *Swansea*



United States

Nasar U Ahmed, *Miami*
Mike Allen, *Milwaukee*
Srinivas Ayyadevara, *Little Rock*
Charles F Babbs, *West Lafayette*
Janet Barletta, *Baltimore*
Lawrence T Bish, *Philadelphia*
Richard W Bohannon, *Storrs*
M Ahmad Chaudhry, *Burlington*
Pei Chen, *Beltville*
Tao Chen, *Jefferson*
Yong Q Chen, *Winston-Salem*
Machado Christian, *Southfield*
Patricia Ann D'Amore, *Boston*
Undurti N Das, *Shaker Heights*
Feng Ding, *Chapel Hill*
Mary E Edgerton, *Houston*
D Mark Estes, *Athens*
Bingliang Fang, *Houston*
Ronnie Fass, *Tucson*
Vesna D Garovic, *Rochester*
Alexandros Georgakilas, *Greenville*
Ronald Gillam, *Logan*
Shannon S Glaser, *Temple*
Ga Nagana Gowda, *West Lafayette*
Anton B Guliaev, *San Francisco*
Zong Sheng Guo, *Pittsburgh*

James P Hardwick, *Rootstown*
Diane M Harper, *Kansas*
Odette A Harris, *Stanford*
Rod Havriluk, *Tallahassee*
Moonseong Heo, *Bronx*
Guoyuan Huang, *Evansville*
Michael Huncharek, *Columbia*
Reinhold J Hutz, *Milwaukee*
Bankole A Johnson, *Charlottesville*
Joseph M Kaminski, *Silver Spring*
Yong S Kim, *Bethesda*
Mark S Kindy, *Charleston*
Jennifer Kisamore, *Tulsa*
Georgios D Kitsios, *Boston*
Ronald Klein, *Shreveport*
Heidemarie Kremer, *Miami*
S Lakshmivarahan, *Norman*
Dawei Li, *New Haven*
Kenneth Maiese, *Newark*
Sameer Malhotra, *New York*
JL Mehta, *Little Rock*
Ray M Merrill, *Provo*
M Mimeault, *Nebraska*
Ron B Mitchell, *St Louis*
Anirban P Mitra, *Los Angeles*
Walter P Murphy, *Evanston*
Marja Tuuli Nevalainen, *Philadelphia*
Yan Peng, *Dallas*
George Perry, *San Antonio*
Ilona Petrikovics, *Huntsville*
Shengping Qin, *Davis*
Peter J Quesenberry, *Providence*
P Hemachandra Reddy, *Beaverton*
James V Rogers, *Columbus*
Troy Rohn, *Boise*
Paul R Sanberg, *Tampa*
Tor C Savidge, *Galveston*
Dong-Chul Seo, *Bloomington*
Igor Sevostianov, *Las Cruces*
Judy Y Tan, *Hillside*
Weihong Tan, *Gainesville*
Guangwen Tang, *Boston*
Paul D Terry, *Knoxville*
Guochuan Emil Tsai, *Torrance*
Catherine E Ulbricht, *Somerville*
Thomas TH Wan, *Orlando*
Xiao-Jing Wang, *Aurora*
Jang-Yen Wu, *Boca Raton*
Qing Wu, *Scottsdale*
Eleftherios S Xenos, *Lexington*
Lijun Xia, *Oklahoma City*
Xiong Xu, *New Orleans*
Li-Jun Yang, *Gainesville*
Wancai Yang, *Chicago*
Xinan Yang, *Chicago*
Fahd A Zarrouf, *Anderson*
Henry Zeringue, *Pittsburgh*
Jingbo Zhang, *New York*



Uruguay

Matias Victoria, *Salto*



REVIEW

- 1 Measurement of body composition as a surrogate evaluation of energy balance in obese patients

Rotella CM, Dicembrini I

ABOUT COVER

Editorial Board Member of *World Journal of Methodology*, Jan-Christoffer Luers, MD, PhD, Associate Professor, Department of Otorhinolaryngology, Head and Neck Surgery, University of Cologne, Cologne 50924, Germany

AIM AND SCOPE

World Journal of Methodology (*World J Methodol*, *WJM*, online ISSN 2222-0682, DOI: 10.5662) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

The primary task of *WJM* is to rapidly publish high-quality original articles, reviews, and commentaries that deal with the methodology to develop, validate, modify and promote diagnostic and therapeutic modalities and techniques in preclinical and clinical applications. *WJM* covers topics concerning the subspecialties including but not exclusively anesthesiology, cardiac medicine, clinical genetics, clinical neurology, critical care, dentistry, dermatology, emergency medicine, endocrinology, family medicine, gastroenterology and hepatology, geriatrics and gerontology, hematology, immunology, infectious diseases, internal medicine, obstetrics and gynecology, oncology, ophthalmology, orthopedics, otolaryngology, radiology, serology, pathology, pediatrics, peripheral vascular disease, psychiatry, radiology, rehabilitation, respiratory medicine, rheumatology, surgery, toxicology, transplantation, and urology and nephrology.

INDEXING/ABSTRACTING

World Journal of Methodology is now indexed in PubMed Central, PubMed, Digital Object Identifier.

FLYLEAF

I-III Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Yue-Li Tian*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Methodology

ISSN
ISSN 2222-0682 (online)

LAUNCH DATE
September 26, 2011

FREQUENCY
Quarterly

EDITOR-IN-CHIEF
Yicheng Ni, MD, PhD, Professor, Department of Radiology, University Hospitals, KU, Leuven, Herestraat 49, B-3000, Leuven, Belgium

EDITORIAL OFFICE
Jin-Lei Wang, Director
Xiu-Xia Song, Vice Director

World Journal of Methodology
Room 903, Building D, Ocean International Center,
No. 62 Dongsihuan Zhonglu, Chaoyang District,
Beijing 100025, China
Telephone: +86-10-85381891
Fax: +86-10-85381893
E-mail: editorialoffice@wjnet.com
Help Desk: <http://www.wjnet.com/esps/helpdesk.aspx>
<http://www.wjnet.com>

PUBLISHER
Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjnet.com
Help Desk: <http://www.wjnet.com/esps/helpdesk.aspx>
<http://www.wjnet.com>

PUBLICATION DATE
March 26, 2015

COPYRIGHT

© 2015 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

Full instructions are available online at http://www.wjnet.com/2222-0682/g_info_20100722180909.htm

ONLINE SUBMISSION

<http://www.wjnet.com/esps/>

Measurement of body composition as a surrogate evaluation of energy balance in obese patients

Carlo Maria Rotella, Ilaria Dicembrini

Carlo Maria Rotella, Ilaria Dicembrini, Section of Endocrinology, Department of Biomedical, Experimental and Clinical Sciences, University of Florence Medical School, 50121 Firenze, Italy
Carlo Maria Rotella, Ilaria Dicembrini, Obesity Agency, Diabetes Agency, Careggi University Hospital, 50127 Florence, Italy
Ilaria Dicembrini, Diabetes Agency, Careggi University Hospital, 50127 Florence, Italy

Author contributions: Rotella CM and Dicembrini I contributed equally to this work.

Conflict-of-interest: No.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Carlo Maria Rotella, Professor, Section of Endocrinology, Department of Biomedical, Experimental and Clinical Sciences, University of Florence Medical School, viale Pieraccini 6, 50127 Florence, Italy. c.rotella@dfc.unifi.it
Telephone: +39-055-4271427

Fax: +39-055-4271474

Received: November 26, 2013

Peer-review started: November 26, 2013

First decision: January 8, 2014

Revised: February 18, 2015

Accepted: March 16, 2015

Article in press: March 18, 2015

Published online: March 26, 2015

as free-fat mass, muscle mass, and fat mass. Among the numerous techniques actually available, bioelectrical impedance analysis seems to be the most suitable in a clinical setting because it is simple, inexpensive, noninvasive, and highly reproducible. To date, there is no consensus concerning the use of one preferred equation for the resting energy expenditure in overweight and/or obese population. Energy restriction alone is an effective strategy to achieve an early and significant weight loss, however it results in a reduction of both fat and lean mass therefore promoting or aggravating an unfavourable body composition (as sarcobesity) in terms of mortality and comorbidities. Therefore the implementation of daily levels of physical activity should be simultaneously promoted. The major role of muscle mass in the energy balance has been recently established by the rising prevalence of the combination of two condition as sarcopenia and obesity. Physical exercise stimulates energy expenditure, thereby directly improving energy balance, and also promotes adaptations such as fiber type, mitochondrial biogenesis, improvement of insulin resistance, and release of myokines, which may influence different tissues, including muscle.

Key words: Obesity; Body composition; Bioelectrical impedance analysis; Energy expenditure; Sarcobesity

© **The Author(s) 2015.** Published by Baishideng Publishing Group Inc. All rights reserved.

Abstract

In clinical practice obesity is primarily diagnosed through the body mass index. In order to characterize patients affected by obesity the use of traditional anthropometric measures appears misleading. Beyond the body mass index, there are overwhelming evidences towards the relevance of a more detailed description of the individual phenotype by characterizing the main body components

Core tip: There are overwhelming evidences towards the relevance of a more detailed description of the individual phenotype by characterizing the main body components as free-fat mass, muscle mass, and fat mass. Among the numerous techniques actually available, bioelectrical impedance analysis seems to be the most suitable in a clinical setting because it is simple, inexpensive, noninvasive, and highly reproducible. To date, there is no consensus concerning the use of one preferred equation for the resting energy expenditure in overweight and/or obese population.

Rotella CM, Dicembrini I. Measurement of body composition as a surrogate evaluation of energy balance in obese patients. *World J Methodol* 2015; 5(1): 1-9 Available from: URL: <http://www.wjgnet.com/2222-0682/full/v5/i1/1.htm> DOI: <http://dx.doi.org/10.5662/wjm.v5.i1.1>

INTRODUCTION

Obesity, defined by excessive adipose tissue, has been shown several deleterious effects on many body organs through thrombogenic, atherogenic, oncogenic, hemodynamic and neurohumoral pathways and has been linked to several chronic diseases, as diabetes, ischemic heart, and musculo-skeletal disorders, together with malignancies. Overweight and obesity represent the fifth leading risk for global death by The World Health Organization on March 2013^[1]. At least 2.8 million adults die each year as a result of overweight/obesity. Moreover the 44% of diabetes and 23% of ischemic heart disease burden can be attributable to an abnormal adipose tissue accumulation^[1].

Obesity is diagnosed accordingly to the body mass index (BMI). This index has been strongly recommended for use in clinical practice^[2,3]. Multiple studies have shown a U or J-shaped relation between BMI, all causes and/or cardiovascular mortality, thus identifying the best survival rate for BMI values in overweight (25-27 kg/m²) followed by a dramatic increase in risk profile out of these values^[4-9].

In a cross-sectional study enrolling 13601 subjects (45.5 ± 17 years; 48% men) from the Third National Health and Nutrition Examination Survey (NHANES)^[10], the BMI cut-off of ≥ 30 kg/m² shown an overall low sensitivity and an high specificity to identify obesity such as defined by an excessive body fat percentage at bioelectrical impedance analysis (> 25% in men and > 35% in women, respectively, thus according to the gold standard definition proposed by the World Health Organization^[2]). The BMI-based definition of obesity has important limitations on diagnostic performance, particularly in men and elderly, missing more than 50% of people with excessive fat mass. In men, BMI showed a more reliable association with lean mass than with total body fat. In contrast, in women BMI appears to be more accurate to estimate fat-mass, thus explaining why overweight women have been reported as more consistently related to increased mortality compared to overweight male individuals. Furthermore, there is significant inter-subject variability in body fat percentage in individuals with similar BMI values. About 30% of obese subjects do not show any metabolic complications such as diabetes, hypertension, dyslipidemia, etc.), then they can be defined metabolically healthy obese patients^[11]. On the other hand, in subgroups of BMI-based normal weight individuals an increased cardiometabolic risk profile and an insulin resistant condition could be described

and strictly related to fat mass accumulation^[12,13].

Further studies have pointed out the association between visceral adiposity, better than BMI, mortality and cardiovascular disease. Waist circumference as a marker of abdominal adipose tissue, has been placed as one of the main contributors to the Metabolic Syndrome by the National Cholesterol Education Program Adult Treatment Panel III in 2001^[14], then as the core feature of the diagnostic criteria proposed by the International Diabetes Federation in 2005.

In the International Day for Evaluation of Abdominal adiposity cross-sectional study, enrolling 168.000 patients in 63 countries, waist circumference showed a higher adjusted odds ratio (95%CI) for diabetes than BMI in overall sample population [1.35 (1.30-1.40) vs 1.23 (1.19-1.27) for men and 1.55 (1.50-1.60) vs 1.23 (1.19-1.27) for women, respectively) such as for cardiovascular disease [1.24 (1.19-1.28) vs 1.13 (1.09-1.17) for men and 1.21 (1.17-1.25) vs 1.20 (1.16-1.24) for women, respectively]^[15].

In the NHANES I and NHANES II longitudinal prospective cohort studies, performed in 10169 male subjects a continuous positive relationship between fat-mass and all-cause mortality rate (HR = 1.033, 95%CI: 1.005-1.063, *P* = 0.0213) have been reported, together with a preponderant significantly negative relationship between free-fat mass and all-cause mortality (*i.e.*, protective; HR = 0.923, 95%CI: 0.906-0.941, *P* < 0.0001)^[16].

Furthermore, BMI and weight loss rate may not be linked to clinical improvement of health-related outcomes, in comparison to different body composition measures by simple and non-invasive methods, which demonstrated a strong relationship between mortality, body lean mass and adipose tissue^[17-24].

In conclusion beyond the BMI, there are overwhelming evidences towards a more detailed description of the individual phenotype by characterizing the main body components as free-fat mass (FFM), muscle mass (MM), and fat mass (FM).

BODY COMPOSITION MEASUREMENTS: TECHNOLOGICAL ADVANCES AND CURRENTLY AVAILABLE NON INVASIVE TECHNIQUES

Numerous techniques for body composition analysis are currently available: anthropometry, including the 4-skinfold method, hydrostatic weighing, *in vivo* neutron activation analysis, anthropogammametry from total body ⁴⁰K, nuclear magnetic resonance, dual-energy X-ray absorptiometry (DEXA), computerized tomography (CT), and bioelectrical impedance analysis (BIA). Evaluating complexity, invasiveness, and cost only DEXA, CT, and BIA may represent the methods of choice to assess body composition in clinical practice whereas the other techniques are limited to scientific

purposes.

DEXA estimates body fat and lean mass percentage through a tissue-specific model (fat, lean tissues, and bone) based on X-ray (dose 1-3 mrad) tissue-dependent attenuation^[25]. The largest sample ($n = 22000$ participants) have been analyzed by the NHANES^[26]. DEXA systems currently available for scanning whole-body tissue composition are capable to analyze a wide range of weights including severe obese subjects (> 150 kg). DEXA scans can be subdivided into different body regions, *i.e.*, trunk, arms, and legs, thus identifying and estimating both android and gynoid fat distribution. DEXA can detect only abdominal adipose tissue accumulation without any distinction between visceral and subcutaneous fat because of their similar X-ray attenuation properties and tissue overlapping^[27]. DEXA is the gold standard technique for the evaluation of body composition in clinical research^[28-30], although limited in clinical practice by the radiation exposure, availability and cost. The use of the same DEXA instruments and analysis software are of relevant importance for longitudinal studies, since they could influence body composition measurements^[31].

In patients affected by malignancies, the analysis at the level of the 3rd lumbar vertebra by CT strongly predicted whole body fat and FFM as compared with DEXA^[32]. CT provided an accurate evaluation of body composition, not provided by DEXA or BIA and a X-ray exposition similar to a chest radiography. Although not validated, also CT images of the right thigh halfway showed to be significantly related to overall mortality rate in chronic obstructive pulmonary disease patients^[33]. Despite its wide use at diagnosis and follow-up in severely ill patients, CT scan at the 3rd lumbar vertebra cannot be considered a feasible method to assess body composition in obese population requiring expensive equipment, trained operators and exposure to ionizing radiation.

BIA is based on the capacity of hydrated tissues to conduct electrical energy. The BIA methodology have been described in two ESPEN position papers^[34,35]. The analysis of total body impedance is based on the estimation of total body water. From total body water, prediction equations allow the calculation of FFM and FM. BIA equations have been validated for chronic obstructive pulmonary disease, AIDS, transplant patients and elderly individuals^[36]. A prediction equation for FFM estimation by BIA in adults (age: 20-94 years; BMI: 17.0-33.8 kg/m²) has been proposed^[37] as well as reference values of FFM and FM for a Caucasian population^[38]. Previously reported data showed no significant differences between DEXA and BIA for the assessment of fat mass in overweight and/or obese patients with a significantly good linear correlation^[39,40]. However several factors could limit the validity of BIA in severe obesity^[41].

ENERGY BALANCE IMPLICATIONS FOR ADDRESSING OBESITY: AVAILABLE ENERGY EXPENDITURE MEASUREMENTS

Energy balance is constituted by three major components including energy intake, energy expenditure (EE) and energy storage. Energy expenditure is expressed by resting energy expenditure (REE), as the amount of energy required for the endogenous metabolic activity separated from the metabolic effects of food and physical activity, the food-related thermic effects (TEF), as energy need to absorb and metabolize ingested food, and energy expenditure associated to physical activity (EE_{PA}). EE_{PA} is the most inter-individual variable component of energy expenditure and consists of the amount of physical activity performed multiplied the energy requirement of that activity. The REE is proportional to body mass, particularly to the FFM^[42]. The REE prediction equations evaluate the energy expenditure of 2 different body compartments as the adipose tissue or FM and the lean mass or FFM. FFM is the main source for REE and is commonly considered as a surrogate measure of metabolically active tissues. Brain, liver, heart, and kidney account for approximately the 60% of REE, despite a combined weight $< 7\%$ of FFM. In comparison to the skeletal muscle, the metabolic rate of heart and kidney is approximately 30-fold higher, and the later approximately 2-fold higher in comparison to brain and liver. The skeletal muscle represents the major contributor to the FFM (50%), but accounts for only the 21% of the REE. Adipose tissue has a low energy expenditure, however its body rate varies significantly according to the dramatic increase of overweight and obesity. Then it is notable that FM represent an important source for the REE in all prediction models. Among adults, REE is lower in elderly, almost in part explained by the change in body composition^[27].

Advantage and limitations of currently available non invasive techniques for the estimation of EE are summarized in Table 1.

Indirect calorimetry represents the gold standard technique to evaluate REE. This method used the rates of oxygen consumption (V_{O_2}) and/or carbon dioxide production (V_{CO_2}) to calculate EE according to the Weir predictive equation, derived from studies comparing indirect calorimetry with direct calorimetry^[43]. The complexity and cost, together with the requirement of the patient isolation for at least 24 h limit the feasibility of the direct calorimetry in humans.

The indirect calorimetry consists in a gas collector, a canopy and a system that measures the volume and concentrations of O₂ and CO₂ minute by minute. Through a unidirectional valve located in the ventilated canopy, the calorimeter collect and quantify the volume and concentration of O₂ inspired and of CO₂ expired. A systematic literature review aimed to determine the

Table 1 Advantages and limitations of available techniques the measurement of Energy Expenditure

Technique	Advantages	Limitations
Direct calorimetry	The gold standard method for measure EE in animal models	High complexity, high cost, need to confine the subject for almost 24 h
Indirect calorimetry	The gold standard to measure REE in humans. Non invasive, adequately accurate, highly reproducible	High cost, relatively complex, need of trained personnel
Bioelectrical impedance analysis	Non invasive, simple, adequately accurate for body composition analysis, relatively inexpensive	The estimation of EE is limited by the need of obesity-specific predictive equations
Multi-sensor device	Easy and practical to use	The estimation of EE is limited by the need of obesity-specific predictive equation

EE: Energy expenditure; REE: Resting energy expenditure.

optimal conditions for obtaining reliable measures of REE by indirect calorimetry recommends fasting for at least 6 h, avoiding caffeine during the night, nicotine and alcohol for at least 2 h, moderate physical activity for at least 2 h, and vigorous physical activity for 14 h before^[44]. Despite methodological, environmental and individual limitations indirect calorimetry represents a non-invasive and very accurate method to estimate REE^[45]. However this method is widely limited in most of clinical settings by the requirement of expensive equipment, and trained operators, then many efforts are spent to identify the most accurate predictive equation to determine REE in overweight and obesity.

Predictive equations have generally been calculated from experimental studies performed in healthy subjects on the basis of regression analysis of bodyweight, height, sex, and age as independent variables and REE by indirect calorimetry as a dependent variable. On the basis of a review of available evidences from Harris and Benedict^[46], FAO/WHO/UNU weight or weight and height equations^[47], and the equations of Mifflin *et al.*^[48] and Owen *et al.*^[49], Frankenfield *et al.*^[50] proposed the use of the Mifflin equation both for overweight and obese subjects. All the available predictive equations have been further validated with indirect calorimetry from adults aged 18-65 years with a BMI of 25 to 40 kg/m² in order to identify the most accurate and precise REE predictive equation in specific overweight and obese groups of United States and Dutch subjects: the results of this study were similar to the data of Frankenfield *et al.*^[50], then supporting the use of Mifflin equation in the United States. However for overweight and obese Dutch adults, there appears to be no accurate equation^[51]. This discrepancy for the Dutch than for the United States adults could be explained by the difference about weight and height values, even within sex and BMI subgroups, thus limiting the validity of this equation in similar taller populations. Numerous studies contributed to further evaluate the currently available predictive equations in overweight and obese subjects^[52-55] and/or in extremely obese subjects^[56-61]. There is some evidences supporting the Mifflin^[59], the FAOW^[56,58] and the Harris Benedict^[60], the Siervo equations^[61] in extremely obese subjects. More recently a new equation, using FFM, Horie-Waitzberg, and Gonzalez equations, have been validated and proposed in Brazilian severely obese

subjects^[62]. The most commonly proposed equation for the measurement of REE have reported in Table 2.

To date there is no consensus about the use of one REE equation compared to others, in overweight and/or obese population. This might be related to differences about group composition, methods, or statistical analysis, at least in part.

In order to estimate EE including EE_{PA}, a practical multi-sensor device, the SenseWear Armband (BodyMedia, Inc., Pittsburgh) has been recently developed. This device contains four sensors to detect heat flux, accelerometry, galvanic skin reaction, skin temperature. Despite excellent results by comparison with energy expenditure measured using the doubly labeled water in overweight/obese children^[63] and lactating women^[64], other evidences underline the need of obesity-specific equations^[65,66].

A meta-analysis of randomized clinical trial suggest that pedometer use is associated with a significant decrease of body weight and blood pressure^[67]. However a recent review assessing different accelerometers to measure daily physical activity as a surrogate of EE in comparison with the doubly labeled water only few available devices have been proven to adequately correlate with the reference method^[68].

BEYOND THE ADIPOSE TISSUE EXCESS: THE IMPORTANCE OF MUSCLE MASS IN OBESITY

Skeletal muscle plays a critical role in the glucose metabolism and peripheral insulin sensitivity as well as musculoskeletal performance. The importance of lean mass on energy balance has been widely recognized by the rising prevalence of the combination of two condition as sarcopenia and obesity^[69]. Two major consensus documents provide different definition of sarcopenia. The European consensus, the Working Group on Sarcopenia in Older People, defines sarcopenia as generalized loss of skeletal muscle mass and strength^[70]. The International consensus, the International Working Group on Sarcopenia, requires a decline in muscle mass and walking speed to diagnose sarcopenia^[71]. Although sarcopenic obesity is commonly used to define the coexistence of diminished muscle mass and increased adipose tissue, a standard

Table 2 Most commonly proposed predictive equations for the estimation of the resting energy expenditure

	Age	Sex	Equation
Harris and Benedict (kcal/d)	15-74	Male	$66.4730 + 13.7516 (W) + 5.0033 (H) - 6.7550 (A)$
	15-74	Female	$655.0955 + 9.5634 (W) + 1.8496 (H) - 4.6756 (A)$
Schofield (MJ/die)	10-17	Male	$0.074 (W) + 2.754$
	10-17	Female	$0.056 (W) + 2.898$
	18-29	Male	$0.063 (W) + 2.896$
	18-29	Female	$0.062 (W) + 2.036$
	30-59	Male	$0.048 (W) + 3.653$
	30-59	Female	$0.034 (W) + 3.538$
	≥ 60	Male	$0.049 (W) + 2.459$
	≥ 60	Female	$0.038 (W) + 2.755$
FAO/ WHO/ UNU (MJ/d)	10-17	Male	$0.0732 (W) + 2.72$
	10-17	Female	$0.0510 (W) + 3.12$
	18-29	Male	$0.0640 (W) + 2.84$
	18-29	Female	$0.0615 (W) + 2.08$
	30-60	Male	$0.0485 (W) + 3.67$
	30-60	Female	$0.0364 (W) + 3.47$
	> 60	Male	$0.0565 (W) + 2.04$
	> 60	Female	$0.0439 (W) + 2.49$
Mifflin-St Jeor (kcal/d)	19-78	Male	$10 \times W + 6.25 \times H - 5 \times A + 5$
	19-78	Female	$10 \times W + 6.25 \times H - 5 \times A - 161$
Owen (kcal/d)	18-65	Male	$879 + 10.2 \times W$
	18-65	Female	$795 + 7.18 \times W$

To convert MJ into kcal, multiply the result by 239. W: Body weight (kg); H: Height (cm); A: Age (years).

definition of sarcopenic obesity is still lacking. Several clinical studies have indicated that obesity and/or insulin resistance may underlie the development of sarcopenia^[72]. A possible role of vitamin D in sarcopenia has been postulated in two studies demonstrating that serum 25-hydroxy vitamin D was negatively correlated with appendicular (legs and arms) fat mass and positively associated with appendicular muscle mass, both evaluated through DEXA analysis^[73,74].

Fat mass excess may induce and/or worsen sarcopenia because the increase of lipid depot reduces both amino acid utilization and protein synthesis in muscle fibers. Evaluating 3132 elderly male subjects without diabetes the highest quartile of homeostasis model assessment of insulin resistance showed the highest risk for a decrease in lean body mass and appendicular mass^[75]. On the counterpart, skeletal muscle is the main target of insulin, then loss of muscle mass may cause insulin resistance. A previous study identified sarcopenia as a risk factor for exacerbating insulin resistance in obese subject with dysglycemia^[76].

Skeletal muscle fibers can be classified into different categories: from slow (type I) fibers with low contractile abilities, numerous mitochondria and high oxidative energy metabolism, to type IIa and type II d/x, and eventually fast (type II b) fibers able to contract rapidly and predisposed toward glycolytic processes. In obese and diabetic population skeletal muscle commonly contains more type II b fibers, while slow muscle fiber percentage and skeletal muscle glucose transport are significantly reduced. Sarcopenic individuals have been shown to be associated with

a further impairment of muscle fiber content, predominantly of type I^[77]. Recent studies have shown that exercise is able to increase the content of type I and type II a fibers^[78]. Furthermore, in skeletal muscle with ageing both the number and morphology of mitochondria are changed with an impaired function and an associated reduced oxidative capacity^[79]. In particular since the primary role of mitochondria is to maintain the cellular energy balance, shifts in mitochondria respiratory activity can lead to a reduced maximal capacity of the tricarboxylic acid cycle and the electron transport chain together with an incomplete lipid-induced upregulation of β -oxidation rates, thus inducing a significant accumulation of intramyocellular lipids and further impairing whole body insulin resistance^[79]. Physical exercise has been demonstrated to induce mitochondria biogenesis, upregulates skeletal muscle gene expression and protein synthesis, and increases skeletal muscle oxidative capacity both in older and sedentary populations^[80].

Moreover, recent evidences promote the hypothesis of cross talk between muscle and different tissues mediated by cytokines and other peptides called myokines^[81] which are involved both in acute exercise-induced metabolic reactions, as well as in the long-term metabolic benefits induced by exercise^[82]. Irisin is one of the most recently identified myokines. Following regular physical activity, Irisin increases two-fold its circulating levels and promotes the shift of white adipocytes into "brite" cells: white adipocytes with a brown-fat-like phenotype. Brown adipocytes activate thermogenesis *via* the mitochondria uncoupling protein UCP-1 (Figure 1). Overexpression of irisin




Figure 1 Physical activity increases the intramuscular expression of the membrane protein fibronectin type III domain containing 5 (FNDC5) which is cleaved to form Irisin, a myokine. Irisin may stimulate the transformation of white adipocytes into brite cells, thus suggested by a marked increase in the expression of uncoupling protein 1 (UCP-1). Adapted from Pedersen^[82].

determined through a gene therapy approach mediated by adenoviral particles, and leading to a modest approximately 3 fold increase in circulating levels, induced browning of subcutaneous white adipose tissue, stimulating a 10-20 fold increase in UCP1, thus increasing energy expenditure and improving glucose tolerance of high fat fed mice^[83]. Despite more recent *in vitro* evidences against the possible translation of the beneficial effects observed in mice to humans, the major role of muscle mass in the energy balance has been definitively established.

In conclusion, physical exercise increases EE and stimulates muscle adaptation mechanisms with potential benefits on different tissues.

ENERGY BALANCE, PHYSICAL ACTIVITY AND DIETARY RESTRICTION

Analysis from the NHANES database describes an average daily intake increase of 168 kcal/d for men and of 335 kcal/d for women, from 1971 to 2000^[84]. On the counterpart, examining in 2004, the physical activity patterns in a typical agrarian population in the United States through pedometers an average of about 18000 steps per day in men and of about 14000 steps per day have been reported^[85], whereas an average American adult walked about 5000 steps per day^[86].

An energy balance flipping point between energy intake and energy expenditure could be recognized in United States around 60' years, thereafter followed by energy intake continuously driving energy expenditure, together with a worldwide burden in the incidence of obesity and type 2 diabetes^[87].

The most frequently proposed strategy for treating obesity is food restriction. Energy restriction alone is an effective strategy to achieve a rapid and substantial weight loss, however it results in a reduction of both fat and muscle mass therefore promoting or aggravating an detrimental body composition (as sarcobesity) in terms of mortality and comorbidities. Specifically, approximately

25% of weight loss obtained through short-term low energy diets is lean muscle mass^[88-92]. Moreover, the lack of success in weight loss maintenance after low-energy regimen and the subsequent weight regained comprising of up to 80% fat mass compound a further detrimental body composition impairment^[93,94]. Weight loss should not be considered the sole focus of therapeutic approach aimed to decrease obesity-related disease risk profile. A systematic review by Chaston *et al.*^[95] demonstrates that very low calories regimens (VLCD) result in significant greater loss of FFM (lean) in comparison to low-calories diets^[95]. Current evidences sustained that the most beneficial long-term outcomes are achieved with a modest energy deficit (2000-4000 kJ/d)^[96]. Therefore increased levels of daily physical activity should be simultaneously promoted. Although resistance training retains or improves the relative percentage of lean mass to total body, current evidence failed to detect a loss of fat mass of a similar magnitude to aerobic training which promotes fat mass loss together with beneficial changes in muscle tissue^[97-99].

Studies evaluating efficacy of dietary restriction together with exercise program showed more favorable results (as weight loss rate and body composition improvement) compared to diet or exercise prescription alone^[69]. Emerging data suggest that the combination of resistance and aerobic exercise with a modest energy restriction was successful for preserving skeletal muscle concomitantly with a significant decrease of fat mass^[100].

CONCLUSION

In order to characterize patients affected by obesity the use of traditional anthropometric measures appears misleading. A systematic and deep phenotyping of these patients, thus integrating data from body composition analysis and energy expenditure should be used in a dynamic rather than only basal approach to define and periodically verify the efficacy of the therapeutic regimen proposed. Among the numerous techniques evaluated in this paper, BIA seems to be the most suitable in a clinical setting because it is simple, inexpensive, noninvasive, and highly reproducible. Moreover, a recent paper by our research group showed that the FM and MM percentage estimated by BIA at baseline should be considered as predictors of success (weight loss > 5% at 6 mo from baseline) in a individual cognitive-behavioral program^[40].

REFERENCES

- 1 **World Health Organization.** Fact Sheet N° 311. Available from: URL: <http://www.who.int/mediacentre/factsheets/fs311/en/index.html>.
- 2 **World Health Organization.** Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organ Tech Rep Ser* 1995; **854**: 1-452 [PMID: 8594834]
- 3 **Seidell JC, Kahn HS, Williamson DF, Lissner L, Valdez R.** Report

- from a Centers for Disease Control and Prevention Workshop on use of adult anthropometry for public health and primary health care. *Am J Clin Nutr* 2001; **73**: 123-126 [PMID: 11124761]
- 4 **Romero-Corral A**, Montori VM, Somers VK, Korinek J, Thomas RJ, Allison TG, Mookadam F, Lopez-Jimenez F. Association of bodyweight with total mortality and with cardiovascular events in coronary artery disease: a systematic review of cohort studies. *Lancet* 2006; **368**: 666-678 [PMID: 16920472]
- 5 **Franzosi MG**. Should we continue to use BMI as a cardiovascular risk factor? *Lancet* 2006; **368**: 624-625 [PMID: 16920449]
- 6 **McGee DL**. Body mass index and mortality: a meta-analysis based on person-level data from twenty-six observational studies. *Ann Epidemiol* 2005; **15**: 87-97 [PMID: 15652713]
- 7 **Flegal KM**, Graubard BI, Williamson DF, Gail MH. Cause-specific excess deaths associated with underweight, overweight, and obesity. *JAMA* 2007; **298**: 2028-2037 [PMID: 17986696]
- 8 **Yusuf S**, Hawken S, Ounpuu S, Bautista L, Franzosi MG, Commerford P, Lang CC, Rumboldt Z, Onen CL, Lisheng L, Tanomsup S, Wangai P, Razak F, Sharma AM, Anand SS. Obesity and the risk of myocardial infarction in 27,000 participants from 52 countries: a case-control study. *Lancet* 2005; **366**: 1640-1649 [PMID: 16271645 DOI: 10.1371/journal.pone.0063999]
- 9 **Price GM**, Uauy R, Breeze E, Bulpitt CJ, Fletcher AE. Weight, shape, and mortality risk in older persons: elevated waist-hip ratio, not high body mass index, is associated with a greater risk of death. *Am J Clin Nutr* 2006; **84**: 449-460 [PMID: 16895897]
- 10 **Romero-Corral A**, Somers VK, Sierra-Johnson J, Thomas RJ, Collazo-Clavell ML, Korinek J, Allison TG, Batsis JA, Sert-Kuniyoshi FH, Lopez-Jimenez F. Accuracy of body mass index in diagnosing obesity in the adult general population. *Int J Obes (Lond)* 2008; **32**: 959-966 [PMID: 18283284]
- 11 **Wildman RP**, Muntner P, Reynolds K, McGinn AP, Rajpathak S, Wyllie-Rosett J, Sowers MR. The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: prevalence and correlates of 2 phenotypes among the US population (NHANES 1999-2004). *Arch Intern Med* 2008; **168**: 1617-1624 [PMID: 18695075 DOI: 10.1001/archinte.168.15.1617]
- 12 **Thomas EL**, Saeed N, Hajnal JV, Brynes A, Goldstone AP, Frost G, Bell JD. Magnetic resonance imaging of total body fat. *J Appl Physiol* (1985) 1998; **85**: 1778-1785 [PMID: 9804581]
- 13 **Thomas EL**, Parkinson JR, Frost GS, Goldstone AP, Doré CJ, McCarthy JP, Collins AL, Fitzpatrick JA, Duriguel G, Taylor-Robinson SD, Bell JD. The missing risk: MRI and MRS phenotyping of abdominal adiposity and ectopic fat. *Obesity* (Silver Spring) 2012; **20**: 76-87 [PMID: 21660078 DOI: 10.1038/oby.2011.142]
- 14 Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002; **106**: 3143-3421 [PMID: 12485966 DOI: 10.1161/01.CIR.0000048067.86569.E1]
- 15 **Balkau B**, Deanfield JE, Després JP, Bassand JP, Fox KA, Smith SC, Barter P, Tan CE, Van Gaal L, Wittchen HU, Massien C, Haffner SM. International Day for the Evaluation of Abdominal Obesity (IDEA): a study of waist circumference, cardiovascular disease, and diabetes mellitus in 168,000 primary care patients in 63 countries. *Circulation* 2007; **116**: 1942-1951 [PMID: 17965405]
- 16 **Allison DB**, Zhu SK, Plankey M, Faith MS, Heo M. Differential associations of body mass index and adiposity with all-cause mortality among men in the first and second National Health and Nutrition Examination Surveys (NHANES I and NHANES II) follow-up studies. *Int J Obes Relat Metab Disord* 2002; **26**: 410-416 [PMID: 11896498]
- 17 **Schols AM**, Broekhuizen R, Weling-Scheepers CA, Wouters EF. Body composition and mortality in chronic obstructive pulmonary disease. *Am J Clin Nutr* 2005; **82**: 53-59 [PMID: 16002800]
- 18 **Slinde F**, Grönberg A, Engström CP, Rossander-Hulthén L, Larsson S. Body composition by bioelectrical impedance predicts mortality in chronic obstructive pulmonary disease patients. *Respir Med* 2005; **99**: 1004-1009 [PMID: 15950141]
- 19 **Futter JE**, Cleland JG, Clark AL. Body mass indices and outcome in patients with chronic heart failure. *Eur J Heart Fail* 2011; **13**: 207-213 [PMID: 21138908]
- 20 **Segall L**, Mardare NG, Ungureanu S, Busuioc M, Nistor I, Enache R, Marian S, Covic A. Nutritional status evaluation and survival in haemodialysis patients in one centre from Romania. *Nephrol Dial Transplant* 2009; **24**: 2536-2540 [PMID: 19297358]
- 21 **Beddhu S**, Pappas LM, Ramkumar N, Samore M. Effects of body size and body composition on survival in hemodialysis patients. *J Am Soc Nephrol* 2003; **14**: 2366-2372 [PMID: 12937315]
- 22 **Marin B**, Desport JC, Kajeu P, Jesus P, Nicolaud B, Nicol M, Preux PM, Couratier P. Alteration of nutritional status at diagnosis is a prognostic factor for survival of amyotrophic lateral sclerosis patients. *J Neurol Neurosurg Psychiatry* 2011; **82**: 628-634 [PMID: 21097551 DOI: 10.1136/jnnp.2010.211474]
- 23 **Prado CM**, Lieffers JR, McCargar LJ, Reiman T, Sawyer MB, Martin L, Baracos VE. Prevalence and clinical implications of sarcopenic obesity in patients with solid tumours of the respiratory and gastrointestinal tracts: a population-based study. *Lancet Oncol* 2008; **9**: 629-635 [PMID: 18539529 DOI: 10.1016/S1470-2045(08)70153-0]
- 24 **Kimyagarov S**, Klid R, Levenkrohn S, Fleissig Y, Kopel B, Arad M, Adunsky A. Body mass index (BMI), body composition and mortality of nursing home elderly residents. *Arch Gerontol Geriatr* 2010; **51**: 227-230 [PMID: 19939476 DOI: 10.1016/j.archger.2009.10.013]
- 25 **Leonard CM**, Roza MA, Barr RD, Webber CE. Reproducibility of DXA measurements of bone mineral density and body composition in children. *Pediatr Radiol* 2009; **39**: 148-154 [PMID: 19052738 DOI: 10.1007/s00247-008-1067-7]
- 26 **Kelly TL**, Wilson KE, Heymsfield SB. Dual energy X-Ray absorptiometry body composition reference values from NHANES. *PLoS One* 2009; **4**: e7038 [PMID: 19753111 DOI: 10.1371/journal.pone.0007038]
- 27 **Baracos V**, Caserotti P, Earthman CP, Fields D, Gallagher D, Hall KD, Heymsfield SB, Müller MJ, Rosen AN, Pichard C, Redman LM, Shen W, Shepherd JA, Thomas D. Advances in the science and application of body composition measurement. *JPEN J Parenter Enteral Nutr* 2012; **36**: 96-107 [PMID: 22235108 DOI: 10.1177/0148607111417448]
- 28 **Slosman DO**, Casez JP, Pichard C, Rochat T, Fery F, Rizzoli R, Bonjour JP, Morabia A, Donath A. Assessment of whole-body composition with dual-energy x-ray absorptiometry. *Radiology* 1992; **185**: 593-598 [PMID: 1410379]
- 29 **Steiner MC**, Barton RL, Singh SJ, Morgan MD. Bedside methods versus dual energy X-ray absorptiometry for body composition measurement in COPD. *Eur Respir J* 2002; **19**: 626-631 [PMID: 11998990]
- 30 **Ellegård LH**, Ahlén M, Körner U, Lundholm KG, Plank LD, Bosaeus IG. Bioelectric impedance spectroscopy underestimates fat-free mass compared to dual energy X-ray absorptiometry in incurable cancer patients. *Eur J Clin Nutr* 2009; **63**: 794-801 [PMID: 18478025 DOI: 10.1038/ejcn.2008.35]
- 31 **Genton L**, Hans D, Kyle UG, Pichard C. Dual-energy X-ray absorptiometry and body composition: differences between devices and comparison with reference methods. *Nutrition* 2002; **18**: 66-70 [PMID: 11827768]
- 32 **Mourtzakis M**, Prado CM, Lieffers JR, Reiman T, McCargar LJ, Baracos VE. A practical and precise approach to quantification of body composition in cancer patients using computed tomography images acquired during routine care. *Appl Physiol Nutr Metab* 2008; **33**: 997-1006 [PMID: 18923576 DOI: 10.1139/H08-075]
- 33 **Marquis K**, Debigaré R, Lacasse Y, LeBlanc P, Jobin J, Carrier G, Maltais F. Midthigh muscle cross-sectional area is a better predictor of mortality than body mass index in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2002; **166**: 809-813 [PMID: 12231489]
- 34 **Kyle UG**, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Gómez JM, Heitmann BL, Kent-Smith L, Melchior JC, Pirllich

- M, Scharfetter H, Schols AM, Pichard C. Bioelectrical impedance analysis--part I: review of principles and methods. *Clin Nutr* 2004; **23**: 1226-1243 [PMID: 15380917]
- 35 **Kyle UG**, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Manuel Gómez J, Lilienthal Heitmann B, Kent-Smith L, Melchior JC, Pirlich M, Scharfetter H, M W J Schols A, Pichard C. Bioelectrical impedance analysis-part II: utilization in clinical practice. *Clin Nutr* 2004; **23**: 1430-1453 [PMID: 15556267]
- 36 **Thibault R**, Pichard C. The evaluation of body composition: a useful tool for clinical practice. *Ann Nutr Metab* 2012; **60**: 6-16 [PMID: 22179189 DOI: 10.1159/000334879]
- 37 **Kyle UG**, Genton L, Karsegard L, Slosman DO, Pichard C. Single prediction equation for bioelectrical impedance analysis in adults aged 20--94 years. *Nutrition* 2001; **17**: 248-253 [PMID: 11312069]
- 38 **Kyle UG**, Genton L, Slosman DO, Pichard C. Fat-free and fat mass percentiles in 5225 healthy subjects aged 15 to 98 years. *Nutrition* 2001; **17**: 534-541 [PMID: 11448570]
- 39 **Donini LM**, Poggiogalle E, Del Balzo V, Lubrano C, Faliva M, Opizzi A, Perna S, Pinto A, Rondanelli M. How to estimate fat mass in overweight and obese subjects. *Int J Endocrinol* 2013; **2013**: 285680 [PMID: 23662101 DOI: 10.1155/2013/285680]
- 40 **Dicembrini I**, Pala L, Cresci B, Cremasco F, Rotella CM. Predictors of weight loss in the clinical management of obese patients: the relevance of body composition. *Obesity and Metabolism* 2010; **6**: 29-33
- 41 **Deurenberg P**. Limitations of the bioelectrical impedance method for the assessment of body fat in severe obesity. *Am J Clin Nutr* 1996; **64**: 449S-452S [PMID: 8780361]
- 42 **Hill JO**, Wyatt HR, Peters JC. Energy balance and obesity. *Circulation* 2012; **126**: 126-132 [PMID: 22753534 DOI: 10.1161/CIRCULATIONAHA.111.087213]
- 43 **Haugen HA**, Chan LN, Li F. Indirect calorimetry: a practical guide for clinicians. *Nutr Clin Pract* 2007; **22**: 377-388 [PMID: 17644692]
- 44 **Compher C**, Frankenfield D, Keim N, Roth-Yousey L. Best practice methods to apply to measurement of resting metabolic rate in adults: a systematic review. *J Am Diet Assoc* 2006; **106**: 881-903 [PMID: 16720129]
- 45 **Kee AL**, Isenring E, Hickman I, Vivanti A. Resting energy expenditure of morbidly obese patients using indirect calorimetry: a systematic review. *Obes Rev* 2012; **13**: 753-765 [PMID: 22568725 DOI: 10.1111/j.1467-789X.2012.01000.x]
- 46 **Harris JA**, Benedict FG. A Biometric Study of Human Basal Metabolism. *Proc Natl Acad Sci USA* 1918; **4**: 370-373 [PMID: 16576330]
- 47 **FAO/WHO/UNU**. Energy and protein requirements. Geneva, Switzerland: World Health Organ Tech Rep Ser, 1985
- 48 **Mifflin MD**, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive equation for resting energy expenditure in healthy individuals. *Am J Clin Nutr* 1990; **51**: 241-247 [PMID: 2305711]
- 49 **Owen OE**, Holup JL, D'Alessio DA, Craig ES, Polansky M, Smalley KJ, Kavle EC, Bushman MC, Owen LR, Mozzoli MA. A reappraisal of the caloric requirements of men. *Am J Clin Nutr* 1987; **46**: 875-885 [PMID: 3687821]
- 50 **Frankenfield D**, Roth-Yousey L, Compher C. Comparison of predictive equations for resting metabolic rate in healthy nonobese and obese adults: a systematic review. *J Am Diet Assoc* 2005; **105**: 775-789 [PMID: 15883556]
- 51 **Weijls PJ**. Validity of predictive equations for resting energy expenditure in US and Dutch overweight and obese class I and II adults aged 18-65 y. *Am J Clin Nutr* 2008; **88**: 959-970 [PMID: 18842782]
- 52 **Frankenfield DC**, Rowe WA, Smith JS, Cooney RN. Validation of several established equations for resting metabolic rate in obese and nonobese people. *J Am Diet Assoc* 2003; **103**: 1152-1159 [PMID: 12963943]
- 53 **Siervo M**, Boschi V, Falconi C. Which REE prediction equation should we use in normal-weight, overweight and obese women? *Clin Nutr* 2003; **22**: 193-204 [PMID: 12706138]
- 54 **Müller MJ**, Bosy-Westphal A, Klaus S, Kreyman G, Lührmann PM, Neuhäuser-Berthold M, Noack R, Pirke KM, Platte P, Selberg O, Steiniger J. World Health Organization equations have shortcomings for predicting resting energy expenditure in persons from a modern, affluent population: generation of a new reference standard from a retrospective analysis of a German database of resting energy expenditure. *Am J Clin Nutr* 2004; **80**: 1379-1390 [PMID: 15531690]
- 55 **de Luis DA**, Aller R, Izaola O, Romero E. Prediction equation of resting energy expenditure in an adult Spanish population of obese adult population. *Ann Nutr Metab* 2006; **50**: 193-196 [PMID: 16407645]
- 56 **Das SK**, Saltzman E, McCrory MA, Hsu LK, Shikora SA, Dolnikowski G, Kehayias JJ, Roberts SB. Energy expenditure is very high in extremely obese women. *J Nutr* 2004; **134**: 1412-1416 [PMID: 15173405]
- 57 **Huang KC**, Kormas N, Steinbeck K, Loughnan G, Caterson ID. Resting metabolic rate in severely obese diabetic and nondiabetic subjects. *Obes Res* 2004; **12**: 840-845 [PMID: 15166305]
- 58 **Lazzer S**, Agosti F, Resnik M, Marazzi N, Mornati D, Sartorio A. Prediction of resting energy expenditure in severely obese Italian males. *J Endocrinol Invest* 2007; **30**: 754-761 [PMID: 17993767]
- 59 **Dobratz JR**, Sibley SD, Beckman TR, Valentine BJ, Kellogg TA, Ikramuddin S, Earthman CP. Predicting energy expenditure in extremely obese women. *JPEN J Parenter Enteral Nutr* 2007; **31**: 217-227 [PMID: 17463148]
- 60 **Boullata J**, Williams J, Cottrell F, Hudson L, Compher C. Accurate determination of energy needs in hospitalized patients. *J Am Diet Assoc* 2007; **107**: 393-401 [PMID: 17324656]
- 61 **Weijls PJ**, Vansant GA. Validity of predictive equations for resting energy expenditure in Belgian normal weight to morbid obese women. *Clin Nutr* 2010; **29**: 347-351 [PMID: 19853980 DOI: 10.1016/j.clnu.2009.09.009]
- 62 **Horie LM**, Gonzalez MC, Torrinhas RS, Cecconello I, Waitzberg DL. New specific equation to estimate resting energy expenditure in severely obese patients. *Obesity* (Silver Spring) 2011; **19**: 1090-1094 [PMID: 21233808 DOI: 10.1038/oby.2010.326]
- 63 **Bäcklund K**, Sundelin C, Larsson C. Validity of armband measuring energy expenditure in overweight and obese children. *Med Sci Sports Exerc* 2010; **42**: 1154-1161 [PMID: 19997020]
- 64 **Slinde F**, Bertz F, Winkvist A, Ellegård L, Olausson H, Brekke HK. Energy expenditure by multisensor armband in overweight and obese lactating women validated by doubly labeled water. *Obesity* (Silver Spring) 2013; **21**: 2231-2235 [PMID: 23512821 DOI: 10.1002/oby.20363]
- 65 **Papazoglou D**, Augello G, Tagliaferri M, Savia G, Marzullo P, Maltezos E, Liuzzi A. Evaluation of a multisensor armband in estimating energy expenditure in obese individuals. *Obesity* (Silver Spring) 2006; **14**: 2217-2223 [PMID: 17189549]
- 66 **Bertoli S**, Posata A, Battezzati A, Spadafranca A, Testolin G, Bedogni G. Poor agreement between a portable armband and indirect calorimetry in the assessment of resting energy expenditure. *Clin Nutr* 2008; **27**: 307-310 [PMID: 18276043 DOI: 10.1016/j.clnu.2007.11.005]
- 67 **Bravata DM**, Smith-Spangler C, Sundaram V, Gienger AL, Lin N, Lewis R, Stave CD, Olkin I, Sirard JR. Using pedometers to increase physical activity and improve health: a systematic review. *JAMA* 2007; **298**: 2296-2304 [PMID: 18029834]
- 68 **Plasqui G**, Westerterp KR. Physical activity assessment with accelerometers: an evaluation against doubly labeled water. *Obesity* (Silver Spring) 2007; **15**: 2371-2379 [PMID: 17925461]
- 69 **Parr EB**, Coffey VG, Hawley JA. 'Sarcobesity': a metabolic conundrum. *Maturitas* 2013; **74**: 109-113 [PMID: 23201324]
- 70 **Cruz-Jentoft AJ**, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel JP, Rolland Y, Schneider SM, Topinková E, Vandewoude M, Zamboni M. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 2010; **39**: 412-423 [PMID: 20392703 DOI: 10.1093/ageing/afq034]
- 71 **Fielding RA**, Vellas B, Evans WJ, Bhasin S, Morley JE, Newman AB, Abellan van Kan G, Andrieu S, Bauer J, Breuille D, Cederholm T, Chandler J, De Meynard C, Donini L, Harris T, Kannt A, Keime

- Guibert F, Onder G, Papanicolaou D, Rolland Y, Rooks D, Sieber C, Souhami E, Verlaan S, Zamboni M. Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc* 2011; **12**: 249-256 [PMID: 21527165 DOI: 10.1016/j.jamda.2011.01.003]
- 72 Stenholm S, Harris TB, Rantanen T, Visser M, Kritchevsky SB, Ferrucci L. Sarcopenic obesity: definition, cause and consequences. *Curr Opin Clin Nutr Metab Care* 2008; **11**: 693-700 [PMID: 18827572 DOI: 10.1097/MCO.0b013e328312c37d]
- 73 Kim MK, Baik KH, Song KH, Il Kang M, Park CY, Lee WY, Oh KW. Vitamin D deficiency is associated with sarcopenia in older Koreans, regardless of obesity: the Fourth Korea National Health and Nutrition Examination Surveys (KNHANES IV) 2009. *J Clin Endocrinol Metab* 2011; **96**: 3250-3256 [PMID: 21832109 DOI: 10.1210/jc.2011-1602]
- 74 Seo JA, Cho H, Eun CR, Yoo HJ, Kim SG, Choi KM, Baik SH, Choi DS, Park MH, Han C, Kim NH. Association between visceral obesity and sarcopenia and vitamin D deficiency in older Koreans: the Ansan Geriatric Study. *J Am Geriatr Soc* 2012; **60**: 700-706 [PMID: 22316299 DOI: 10.1111/j.1532-5415.2012.03887.x]
- 75 Lee CG, Boyko EJ, Strotmeyer ES, Lewis CE, Cawthon PM, Hoffman AR, Everson-Rose SA, Barrett-Connor E, Orwoll ES. Association between insulin resistance and lean mass loss and fat mass gain in older men without diabetes mellitus. *J Am Geriatr Soc* 2011; **59**: 1217-1224 [PMID: 21718263 DOI: 10.1111/j.1532-5415.2011.03472.x]
- 76 Srikanthan P, Hevener AL, Karlamangla AS. Sarcopenia exacerbates obesity-associated insulin resistance and dysglycemia: findings from the National Health and Nutrition Examination Survey III. *PLoS One* 2010; **5**: e10805 [PMID: 22421977 DOI: 10.1371/journal.pone.0010805]
- 77 Harrison BC, Leinwand LA. Fighting fat with muscle: bulking up to slim down. *Cell Metab* 2008; **7**: 97-98 [PMID: 18249167 DOI: 10.1016/j.cmet.2008.01.003]
- 78 Eckardt K, Taube A, Eckel J. Obesity-associated insulin resistance in skeletal muscle: role of lipid accumulation and physical inactivity. *Rev Endocr Metab Disord* 2011; **12**: 163-172 [PMID: 21336841 DOI: 10.1007/s11154-011-9168-2]
- 79 Kohara K. Sarcopenic obesity in aging population: current status and future directions for research. *Endocrine* 2014; **45**: 15-25 [PMID: 23821364]
- 80 Peterson CM, Johannsen DL, Ravussin E. Skeletal muscle mitochondria and aging: a review. *J Aging Res* 2012; **2012**: 194821 [PMID: 22888430 DOI: 10.1155/2012/194821]
- 81 Pedersen L, Hojman P. Muscle-to-organ cross talk mediated by myokines. *Adipocyte* 2012; **1**: 164-167 [PMID: 23700527 DOI: 10.4161/adip.20344]
- 82 Pedersen BK. Muscle as a secretory organ. *Compr Physiol* 2013; **3**: 1337-1362 [PMID: 23897689]
- 83 Boström P, Wu J, Jedrychowski MP, Korde A, Ye L, Lo JC, Rasbach KA, Boström EA, Choi JH, Long JZ, Kajimura S, Zingaretti MC, Vind BF, Tu H, Cinti S, Højlund K, Gygi SP, Spiegelman BM. A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* 2012; **481**: 463-468 [PMID: 22237023 DOI: 10.1038/nature10777]
- 84 Centers for Disease Control and Prevention. National Health and Nutrition Examination Survey. Available from: URL: <http://www.cdc.gov/nchs/nhanes.htm>
- 85 Bassett DR, Schneider PL, Huntington GE. Physical activity in an Old Order Amish community. *Med Sci Sports Exerc* 2004; **36**: 79-85 [PMID: 14707772]
- 86 Bassett DR, Wyatt HR, Thompson H, Peters JC, Hill JO. Pedometer-measured physical activity and health behaviors in U.S. adults. *Med Sci Sports Exerc* 2010; **42**: 1819-1825 [PMID: 20305579 DOI: 10.1249/MSS.0b013e3181dc2e54]
- 87 Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, Gortmaker SL. The global obesity pandemic: shaped by global drivers and local environments. *Lancet* 2011; **378**: 804-814 [PMID: 21872749 DOI: 10.1016/S0140-6736(11)60813-1]
- 88 Nicklas BJ, Wang X, You T, Lyles MF, Demons J, Easter L, Berry MJ, Lenchik L, Carr JJ. Effect of exercise intensity on abdominal fat loss during calorie restriction in overweight and obese postmenopausal women: a randomized, controlled trial. *Am J Clin Nutr* 2009; **89**: 1043-1052 [PMID: 19211823 DOI: 10.3945/ajcn.2008.26938]
- 89 Hoie LH, Bruusgaard D, Thom E. Reduction of body mass and change in body composition on a very low calorie diet. *Int J Obes Relat Metab Disord* 1993; **17**: 17-20 [PMID: 8383636]
- 90 Ballor DL, Katch VL, Becque MD, Marks CR. Resistance weight training during caloric restriction enhances lean body weight maintenance. *Am J Clin Nutr* 1988; **47**: 19-25 [PMID: 3337037]
- 91 Campbell WW, Haub MD, Wolfe RR, Ferrando AA, Sullivan DH, Apolzan JW, Iglay HB. Resistance training preserves fat-free mass without impacting changes in protein metabolism after weight loss in older women. *Obesity* (Silver Spring) 2009; **17**: 1332-1339 [PMID: 19247271 DOI: 10.1038/oby.2009.2]
- 92 Foster-Schubert KE, Alfano CM, Duggan CR, Xiao L, Campbell KL, Kong A, Bain CE, Wang CY, Blackburn GL, McTiernan A. Effect of diet and exercise, alone or combined, on weight and body composition in overweight-to-obese postmenopausal women. *Obesity* (Silver Spring) 2012; **20**: 1628-1638 [PMID: 21494229 DOI: 10.1038/oby.2011.76]
- 93 Newman AB, Lee JS, Visser M, Goodpaster BH, Kritchevsky SB, Tykavsky FA, Nevitt M, Harris TB. Weight change and the conservation of lean mass in old age: the Health, Aging and Body Composition Study. *Am J Clin Nutr* 2005; **82**: 872-878; quiz 915-916 [PMID: 16210719]
- 94 Beavers KM, Lyles MF, Davis CC, Wang X, Beavers DP, Nicklas BJ. Is lost lean mass from intentional weight loss recovered during weight regain in postmenopausal women? *Am J Clin Nutr* 2011; **94**: 767-774 [PMID: 21795437 DOI: 10.3945/ajcn.110.004895]
- 95 Chaston TB, Dixon JB, O'Brien PE. Changes in fat-free mass during significant weight loss: a systematic review. *Int J Obes (Lond)* 2007; **31**: 743-750 [PMID: 17075583]
- 96 Strychar I. Diet in the management of weight loss. *CMAJ* 2006; **174**: 56-63 [PMID: 16389240]
- 97 Willis LH, Slentz CA, Bateman LA, Shields AT, Piner LW, Bales CW, Houmard JA, Kraus WE. Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults. *J Appl Physiol* (1985) 2012; **113**: 1831-1837 [PMID: 23019316 DOI: 10.1152/jappphysiol.01370.2011]
- 98 Ismail I, Keating SE, Baker MK, Johnson NA. A systematic review and meta-analysis of the effect of aerobic vs. resistance exercise training on visceral fat. *Obes Rev* 2012; **13**: 68-91 [PMID: 21951360 DOI: 10.1111/j.1467-789X.2011.00931.x]
- 99 Rolland Y, Czerwinski S, Abellan Van Kan G, Morley JE, Cesari M, Onder G, Woo J, Baumgartner R, Pillard F, Boirie Y, Chumlea WM, Vellas B. Sarcopenia: its assessment, etiology, pathogenesis, consequences and future perspectives. *J Nutr Health Aging* 2008; **12**: 433-450 [PMID: 18615225]
- 100 Frimel TN, Sinacore DR, Villareal DT. Exercise attenuates the weight-loss-induced reduction in muscle mass in frail obese older adults. *Med Sci Sports Exerc* 2008; **40**: 1213-1219 [PMID: 18580399 DOI: 10.1249/MSS.0b013e31816a85ce]

P- Reviewer: Bertoli S, Niculescu M, Verduci E S- Editor: Ji FF
L- Editor: A E- Editor: Lu YJ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

